

The rectilinear local crossing number of K_n

Bernardo M. Ábrego

California State University, Northridge
bernardo.abrego@csun.edu

Silvia Fernández-Merchant *

California State University, Northridge
silvia.fernandez@csun.edu

May 20, 2024

Abstract

The *local crossing number* of a drawing of a graph is the largest number of crossings on any edge of the drawing. In a *rectilinear drawing* of a graph, the vertices are points in the plane in general position and the edges are straight-line segments. The *rectilinear local crossing number* of the complete graph K_n , denoted by $\overline{\text{lcr}}(K_n)$, is the minimum local crossing number over all rectilinear drawings of K_n .

We determine $\overline{\text{lcr}}(K_n)$. More precisely, for every $n \notin \{8, 14\}$,

$$\overline{\text{lcr}}(K_n) = \left\lceil \frac{1}{2} \left(n - 3 - \left\lceil \frac{n-3}{3} \right\rceil \right) \left\lceil \frac{n-3}{3} \right\rceil \right\rceil,$$

$\overline{\text{lcr}}(K_8) = 4$, and $\overline{\text{lcr}}(K_{14}) = 15$.

Keywords: local crossing number, complete graph, geometric drawing, rectilinear drawing, rectilinear crossing number, balanced partition.

MSC2010: 05C10, 05C62, 52C10, 68R10.

1 Introduction

We are concerned with rectilinear drawings of the complete graph K_n . That is, drawings where each of the n vertices is a point in the plane, and every edge is drawn as a straight-line segment among these n vertices. We assume that the set of points is in general position; that is, there are no three collinear points.

According to Guy et al. [10] and Kainen [11], Ringel defined the local crossing number of a graph as follows: In a *drawing* of a graph, each vertex is represented by a point and each edge is represented by a simple continuous arc not passing through any vertex other than its endpoints. The *local crossing number* of a drawing D of a graph G , denoted $\text{lcr}(D)$, is the largest number of crossings on any edge of D . The *local crossing number* of G , denoted $\text{lcr}(G)$, is the minimum of $\text{lcr}(D)$ over all drawings D of G . Other authors (like Thomassen [15]) have called it the *cross-index*, but we follow Schaefer [14] who strongly encourages

*Supported by NSF grant DMS-1400653

the use of local crossing number. The equivalent definition for rectilinear drawings is the *rectilinear local crossing number* of G , denoted $\overline{\text{lcr}}(G)$, as the minimum of $\text{lcr}(D)$ over all rectilinear drawings D of G .

Recently, Lara [12] posed the problem of determining $\overline{\text{lcr}}(K_n)$. Together with Rubio-Montiel, and Zaragoza, they claimed that

$$\frac{1}{18}n^2 + \Theta(n) \leq \overline{\text{lcr}}(K_n) \leq \frac{1}{9}n^2 + \Theta(n),$$

where the upper bound appears in [13].

In the 1960s, Guy, Jenkyns, and Schaer [10] worked on the problem of determining the local crossing number of K_n drawn on the surface of a torus. They determined the exact values when $n \leq 9$, and they provided asymptotic estimates based on their bounds of the toroidal crossing number. In particular, they used an argument in the torus equivalent to the following: The sum of the number of crossings of every edge over all edges of a graph G counts precisely twice the number of crossings of G . It follows that

$$\overline{\text{lcr}}(G) \geq \frac{2 \overline{\text{cr}}(G)}{\binom{n}{2}},$$

where $\overline{\text{cr}}(K_n)$ denotes the *rectilinear crossing number* of K_n ; that is the smallest number of crossings among all rectilinear drawings of K_n . The current best lower bound by Ábrego et al. [3], $\overline{\text{cr}}(K_n) \geq \frac{277}{729}\binom{n}{4} + \Theta(n^3)$, yields

$$\overline{\text{lcr}}(K_n) \geq \frac{277}{4374}n^2 + \Theta(n) > \frac{1}{15.8}n^2 + \Theta(n),$$

which is already an improvement over the lower bound by Lara et al. Note that this approach also gives a bound for the local crossing number of K_n when the drawings are not necessarily rectilinear. The current best bound by De Klerk et al. [8], $\text{cr}(K_n) \geq (0.8594/64)n^4 + \Theta(n^3)$, yields

$$\text{lcr}(K_n) \geq \frac{0.8594}{16}n^2 + \Theta(n) > \frac{1}{18.62}n^2 + \Theta(n).$$

In this paper, we determine $\overline{\text{lcr}}(K_n)$ precisely. Our main result is the following.

Theorem 1. *If n is a positive integer, then*

$$\begin{aligned} \overline{\text{lcr}}(K_n) &= \left\lceil \frac{1}{2} \left(n - 3 - \left\lfloor \frac{n-3}{3} \right\rfloor \right) \left\lceil \frac{n-3}{3} \right\rceil \right\rceil \text{ if } n \notin \{8, 14\}, \text{ that is,} \\ \overline{\text{lcr}}(K_n) &= \begin{cases} \frac{1}{9}(n-3)^2 & \text{if } n \equiv 0 \pmod{3}, \\ \frac{1}{9}(n-1)(n-4) & \text{if } n \equiv 1 \pmod{3}, \\ \frac{1}{9}(n-2)^2 - \left\lfloor \frac{n-2}{6} \right\rfloor & \text{if } n \equiv 2 \pmod{3}, n \notin \{8, 14\}. \end{cases} \end{aligned}$$

In addition, $\overline{\text{lcr}}(K_8) = 4$ and $\overline{\text{lcr}}(K_{14}) = 15$.

To prove this theorem, we employ a separation lemma for sets P with n points (Lemma 2). We show that there is either an edge or a path of length 2 whose endpoints are vertices

of the convex hull of P that separates the rest of the set into parts that differ by at most $(n - 2)/3$ or $(n - 3)/3$ points, respectively. This lemma may have applications to other separation or crossing problems. To match the lower bound obtained from Lemma 2, we present a couple of constructions: A simple one that works when $n \not\equiv 2 \pmod{3}$ and a more elaborate construction when $n \equiv 2 \pmod{3}$.

In Section 2, we present the lower bound needed for the proof of Theorem 1. In Sections 3 and 4 we present constructions that match the lower bound, except for a few exceptional cases. In Section 5 we complete the proof of the Theorem, and in Section 6 we present some remarks and open problems. We also note that the remaining special case of $\overline{\text{lcr}}(K_{14}) = 15$ was settled by Aichholzer [5].

2 Lower Bound

Let P be a set of points in the plane. For every pair of distinct points p and q in the plane, denote by $H^+(pq)$ the set of points of P that are on the right side of the oriented line pq . Similarly, define $H^-(pq)$ as the set of points of P on the left side of the oriented line pq . If p , q , and r are three points in the plane, let $S(prq)$ be the set of points of P in the open sector defined by the oriented angle $\angle prq$; that is the sector obtained by rotating a ray counterclockwise around r from rp to rq (excluding both of the rays rp and rq). Finally, let $\triangle prq$ denote the interior of triangle pqr . We first prove the following separation lemma.

Lemma 2. *If P is a set of $n \geq 3$ points in the plane, then one of the following occurs:*

(a) *There are two vertices p and q of the convex hull of P such that*

$$\left| |H^+(pq)| - |H^-(pq)| \right| \leq \frac{n-2}{3}.$$

(b) *There are three points p , q , and r in P , such that p and q are vertices of the convex hull of P and*

$$\left| |S(prq)| - |S(qrp)| \right| \leq \left\lceil \frac{n-3}{3} \right\rceil.$$

Proof. The desired situation is illustrated in Figure 1. Suppose that p_1, p_2, \dots, p_k are the vertices of the convex hull of P in counterclockwise order. We first prove that either (a) occurs or else there are q_1 , q_2 , and q_3 vertices of the convex hull of P such that $\triangle q_1q_2q_3$ contains at least $(n - 3)/3$ points of P .

If $(n - 2)/3 \leq |H^+(p_1p_i)| \leq 2(n - 2)/3$ for some $2 \leq i \leq k$, then

$$\left| |H^+(p_1p_i)| - |H^-(p_1p_i)| \right| \leq \frac{2(n-2)}{3} - \frac{(n-2)}{3} = \frac{n-2}{3},$$

which is condition (a). So we may assume that for each $2 \leq i \leq k$ either $|H^+(p_1p_i)| < (n - 2)/3$ or $|H^+(p_1p_i)| > 2(n - 2)/3$. Because

$$0 = |H^+(p_1p_2)| \leq |H^+(p_1p_3)| \leq \dots \leq |H^+(p_1p_k)| = n - 2,$$




Figure 1: The absolute difference of points between the shaded and not shaded regions is at most $(n-2)/3$ (left) or $\lceil (n-3)/3 \rceil$ (right).

it follows that there is j such that $2 < j \leq k$, $|H^+(p_1 p_{j-1})| < (n-2)/3$, and $|H^+(p_1 p_j)| > 2(n-2)/3$. Then

$$\begin{aligned} |P \cap \triangle p_1 p_{j-1} p_j| &= |H^+(p_1 p_j)| - |H^+(p_1 p_{j-1})| - 1 \\ &\geq \left(\frac{2(n-2)}{3} + \frac{1}{3} \right) - \left(\frac{n-2}{3} - \frac{1}{3} \right) - 1 = \frac{n-3}{3}. \end{aligned}$$

Thus $\triangle p_1 p_{j-1} p_j$ has at least $(n-3)/3$ points of P , which gives the desired triple q_1, q_2, q_3 . We assume that the triangle $q_1 q_2 q_3$ is positively oriented (counterclockwise).

Suppose by contradiction that

$$|S(q_i r q_j)| - |S(q_j r q_i)| > \left\lceil \frac{n-3}{3} \right\rceil,$$

for every point $r \in P \setminus \{q_1, q_2, q_3\}$ and every $i, j \in \{1, 2, 3\}$ with $i \neq j$. It follows that, for every point $r \in P \setminus \{q_1, q_2, q_3\}$, two of the following are less than $(n-3)/3$ and the other is greater than $2(n-3)/3$: $|S(q_1 r q_2)|$, $|S(q_2 r q_3)|$, and $|S(q_3 r q_1)|$.

Let $f(r)$ be the pair (q_i, q_j) corresponding to the sector greater than $2(n-3)/3$. One of the pairs (q_1, q_2) , (q_2, q_3) , or (q_3, q_1) must be the image of at least one third of the points in $P \setminus \{q_1, q_2, q_3\}$. Suppose without loss of generality, that at least $(n-3)/3$ points r have $f(r) = (q_1, q_2)$. Among all those points r , let r_0 be one such that $|S(q_1 r_0 q_2)|$ is minimum. Note that r_0 is in $H^-(q_1 q_2)$, otherwise $S(q_1 r_0 q_2)$ and $\triangle q_1 q_2 q_3$ would be disjoint (Figure 2a) and together would contain more than $2(n-3)/3 + (n-3)/3 = n-3$ points of $P \setminus \{q_1, q_2, q_3\}$. Let $p \in S(q_1 r_0 q_2)$ (Figure 2bc). Then $S(q_1 p q_2)$ is a proper subset of $S(q_1 r_0 q_2)$, and by minimality of r_0 it follows that $|S(q_1 p q_2)| \leq 2(n-3)/3$. Thus $f(p) \neq (q_1, q_2)$. Therefore none of the points p in $S(q_1 r_0 q_2)$ has $f(p) = (q_1, q_2)$; that is, there are less than $(n-4) - 2(n-3)/3 = (n-6)/3$ points r such that $f(r) = (q_1, q_2)$. This is a contradiction. Hence there is a point r and a pair $i, j \in \{1, 2, 3\}$ with $i \neq j$, such that

$$|S(q_i r q_j)| - |S(q_j r q_i)| \leq \left\lceil \frac{n-3}{3} \right\rceil,$$

and condition (b) is satisfied. \square




Figure 2: (a) $\triangle q_1 q_2 q_3 \cup S(q_1 r_0 q_2)$ has more than $n - 3$ points of $P \setminus \{q_1, q_2, q_3\}$. (bc) $S(q_1 p q_2)$ is a proper subset of $S(q_1 r_0 q_2)$.

Now we prove the lower bound.

Theorem 3. *If $n \geq 3$, then*

$$\overline{\text{lcr}}(K_n) \geq \begin{cases} \frac{1}{9}(n-3)^2 & \text{if } n \equiv 0 \\ \frac{1}{9}(n-1)(n-4) & \text{if } n \equiv 1 \pmod{3} \\ \frac{1}{9}(n-2)^2 - \lfloor \frac{n-2}{6} \rfloor & \text{if } n \equiv 2. \end{cases}$$

Proof. The result is trivial for $n = 3$ and 4. Let $n \geq 5$ and consider a set P of n points in the plane. We use the previous lemma. First suppose that there are two vertices p and q of the convex hull of P such that

$$\left| |H^+(pq)| - |H^-(pq)| \right| \leq \frac{n-2}{3}.$$

Every edge xy with $x \in H^+(pq)$ and $y \in H^-(pq)$ intersects the edge pq . Thus pq is crossed at least $|H^+(pq)| \cdot |H^-(pq)|$ times. Because the absolute difference of the factors is at most $(n-2)/3$ and their sum is $n-2$, it follows that for $n \geq 5$

$$|H^+(pq)| \cdot |H^-(pq)| \geq \left\lceil \frac{n-2}{3} \right\rceil \left(n-2 - \left\lceil \frac{n-2}{3} \right\rceil \right) \geq \frac{2}{9}(n-2)^2,$$

which is larger than the desired bound for any congruence class.

Now, suppose that there are three points p , q , and r in P such that p and q are vertices of the convex hull of P and

$$\left| |S(prq)| - |S(qrp)| \right| \leq \left\lceil \frac{n-3}{3} \right\rceil.$$

Every edge xy with $x \in S(prq)$ and $y \in S(qrp)$ intersects the path $pr \cup rq$. Thus the union of the segments pr and rq is crossed at least $|S(prq)| \cdot |S(qrp)|$ times. Because the absolute difference of the factors is at most $\lceil (n-3)/3 \rceil$ and their sum is $n-3$, it follows that

$$|S(prq)| \cdot |S(qrp)| \geq \left\lceil \frac{n-3}{3} \right\rceil \left(n-3 - \left\lceil \frac{n-3}{3} \right\rceil \right).$$

Therefore one of pr or rq must be crossed at least $\lceil \frac{1}{2} \lceil (n-3)/3 \rceil (n-3 - \lceil (n-3)/3 \rceil) \rceil$ times. Hence

$$\overline{\text{lcr}}(K_n) \geq \begin{cases} \frac{1}{9}(n-3)^2 & \text{if } n \equiv 0 \\ \frac{1}{9}(n-1)(n-4) & \text{if } n \equiv 1 \pmod{3} \\ \lceil \frac{1}{18}(n-2)(2n-7) \rceil = \frac{1}{9}(n-2)^2 - \lfloor \frac{n-2}{6} \rfloor & \text{if } n \equiv 2. \end{cases} \quad \square$$

3 The First Construction

To complete the proof of Theorem 1, we present constructions that match the lower bound. The first construction was originally presented by Lara et al. [13], it is based on the currently best known constructions for the rectilinear crossing number of K_n by Ábrego et al. [2, 4] and the conjectured structure of the optimal sets (see [4]). Lara et al. [13] asymptotically calculated $\overline{\text{lcr}}(D)$ for the construction that follows; however, for the purpose of obtaining an exact formula, we have to carefully calculate the exact local crossing number of these drawings. Theorem 4 is tight for $n \not\equiv 2 \pmod{3}$. In the next section, we present a slightly better construction that matches the lower bound for the class $n \equiv 2 \pmod{3}$.

Theorem 4. *If $n \geq 3$, then*

$$\overline{\text{lcr}}(K_n) \leq \begin{cases} \frac{1}{9}(n-3)^2 & \text{if } n \equiv 0 \\ \frac{1}{9}(n-1)(n-4) & \text{if } n \equiv 1 \pmod{3} \\ \frac{1}{9}(n-2)^2 & \text{if } n \equiv 2. \end{cases}$$

Proof. Consider an arc C_0 of circle passing through the points with coordinates $(1, 0)$, $(3, 0)$, and $(2, \varepsilon)$, where ε is a very small positive real. Let C_1 and C_2 be the rotations of C_0 with center at the origin and angles $2\pi/3$ and $4\pi/3$, respectively. We set ε small enough so that any secant line to the arc C_0 separates C_1 from C_2 .

Thus, if $x_0, y_0 \in C_0$, $x_1 \in C_1$, and $x_2 \in C_2$, then the segments x_0x_1 and y_0x_2 do not cross, and the same occurs for the corresponding rotations.

The point set P consists of $n_0 = \lfloor n/3 \rfloor$ points in C_0 , $n_1 = \lfloor (n+1)/3 \rfloor$ points in C_1 , and $n_2 = \lfloor (n+2)/3 \rfloor$ points in C_2 .

Now we determine $\overline{\text{lcr}}(P)$. First suppose that xy is an edge with both endpoints in C_i . Moreover, suppose that there are a points of P in the arc C_i between x and y (see Figure 3(a)). The edges obtained from connecting each of these a points to the remaining $n_i - a - 2$ points on C_i , or to any of the points in C_{i-1} (where the indices are taken modulo 3) all cross xy . By convexity of C_i none of the edges from C_i to C_{i+1} crosses xy , the same is true about edges from C_{i-1} to C_{i+1} . Thus xy is crossed exactly $a(n_{i-1} + n_i - a - 2)$ times. The maximum occurs when $a = n_i - 2$ and it equals $(n_i - 2)n_{i-1}$ which is at most $(n_2 - 2)n_1$.

Now, suppose that xy is an edge with $x \in C_i$ and $y \in C_{i+1}$. Moreover, suppose that there are a points of P before x in C_i and $n_i - 1 - a$ after x , and b points of P before y in C_{i+1} and $n_{i+1} - 1 - b$ after y , as shown in Figure 3(b). Each edge uv with u before x in C_i and v after y in C_{i+1} crosses xy . The same occurs for the edges uv with u after x and v before y . In addition, if uv is an edge in C_{i+1} with u before y and v after y , then uv crosses xy . Edges




Figure 3: The number of crossings of edges in the same C_i or between C_i and C_{i+1}

with both endpoints in C_i or both in C_{i-1} do not cross xy . The same is true for edges with one endpoint in C_{i-1} and the other in C_i or C_{i+1} . Thus xy is crossed exactly

$$b(n_{i+1} - 1 - b) + b(n_i - 1 - a) + a(n_{i+1} - 1 - b)$$

times. This expression factors as

$$(n_{i+1} - 1)(n_i - 1) - a(n_i - 1 - a) - (a + b + 1 - n_{i+1})(a + b + 1 - n_i).$$

Because $0 \leq a \leq n_i - 1$, the second term is nonpositive. The third term is nonpositive because the two factors are either equal or differ by one. Therefore, the maximum number of crossings of xy is $(n_{i+1} - 1)(n_i - 1)$ and it occurs when $a = 0$ and $b = n_{i+1} - 1$ or $b = n_i - 1 = n_{i+1} - 2$ (only if $n_i < n_{i+1}$); or when $a = n_i - 1$ and $b = 0$ or $b = n_{i+1} - n_i = 1$ (only if $n_i < n_{i+1}$). In any case, $(n_{i+1} - 1)(n_i - 1) \leq (n_2 - 1)(n_1 - 1)$, and comparing to the first case, $(n_2 - 2)n_1 \leq (n_2 - 1)(n_1 - 1)$. Therefore,

$$\overline{\text{lcr}}(P) = (n_2 - 1)(n_1 - 1) = \begin{cases} \frac{1}{9}(n - 3)^2 & \text{if } n \equiv 0 \\ \frac{1}{9}(n - 1)(n - 4) & \text{if } n \equiv 1 \\ \frac{1}{9}(n - 2)^2 & \text{if } n \equiv 2. \end{cases} \pmod{3} \quad \square$$

4 The Second Construction

In this section, we present a construction that matches the lower bound of $\overline{\text{lcr}}(K_n)$ when $n \equiv 2 \pmod{3}$. Set $n = 3k + 8$ for some positive integer $k \geq 4$.

Theorem 5. *If $n = 3k + 8$ with $k \geq 4$, then*

$$\overline{\text{lcr}}(K_n) \leq \frac{1}{9}(n - 2)^2 - \left\lfloor \frac{n - 2}{6} \right\rfloor = k^2 + 4k + 3 - \left\lfloor \frac{k}{2} \right\rfloor.$$




Figure 4: The construction when $n = 3k + 8$. Each of the five thick dashed lines joins the endpoints of the corresponding arc.

Proof. The construction consists of five parts with a linear number of points and two additional isolated points. Specifically, let $\varepsilon > 0$ be a small constant to be determined. Let A be a set of $k + 2$ points on the arc of circle with endpoints $(0, 24)$ and $(0, 20)$ that passes through $(\varepsilon, 22)$. Similarly, let B_1 be a set of $k + 2 - \lfloor k/2 \rfloor$ points on the arc of circle with endpoints $(-24, -12)$ and $(-20, -10)$ that passes through $(-22 - \varepsilon, -11 + 2\varepsilon)$; and let C_1 be a set of $\lfloor k/2 \rfloor$ points on the arc of circle with endpoints $(-5, -1)$ and $(-4, 0)$ that passes through $(-4.5 - \varepsilon, -0.5 + \varepsilon)$. Let $x_1 = (-11, 1)$, and set B_2 , C_2 , and x_2 as the reflections with respect to the y -axis of B_1 , C_1 , and x_1 , respectively. Choose ε small enough so that none of the lines tangent to the arcs of circle intersects any of the other arcs of circle. This choice ensures that every line joining points in the same part separates the other parts as shown in Figure 4 (any $\varepsilon < 0.03$ works). The set P consists of $A \cup B_1 \cup B_2 \cup C_1 \cup C_2 \cup \{x_1, x_2\}$ and it has $n = (k + 2) + 2(k + 2 - \lfloor k/2 \rfloor) + 2\lfloor k/2 \rfloor + 2 = 3k + 8$ points.

To determine $\overline{\text{lcr}}(P)$, we proceed as in Theorem 4; that is, we find the maximum number of crossings on a segment xy whose endpoints belong to a prescribed pair of parts. Accounting for symmetry and for the fact that A is on an arc of circle that is concave to the left, the cases that need to be analyzed are the following: (A, A) , $(A, \{x_1\})$, (A, B_1) , (A, C_1) , $(\{x_1\}, \{x_2\})$, $(\{x_1\}, B_1)$, $(\{x_1\}, B_2)$, $(\{x_1\}, C_1)$, $(\{x_1\}, C_2)$, (B_1, B_1) , (B_1, B_2) , (B_1, C_1) , (B_1, C_2) , (C_1, C_1) , and (C_1, C_2) ; where each pair (X, Y) indicates that $x \in X$ and $y \in Y$. In all the following cases with $X \neq Y$, if $p \in \{x, y\}$ belongs to $S \in \{A, B_1, B_2, C_1, C_2\}$, then we call *lower* S the set of points in S that have y -coordinate less than p , and we call *upper* S the set of points

in S that have y -coordinate greater than p .

- (A, A) Let $x, y \in A$. Suppose that there are a points of A between x and y , and $k - a$ outside. The only segments crossing xy are those connecting one of the a inside points in A to points in $C_1 \cup B_1 \cup \{x_1\}$ or to points in A outside xy . The total is

$$a \left(\left\lfloor \frac{k}{2} \right\rfloor + (k + 2 - \left\lfloor \frac{k}{2} \right\rfloor) + 1 + (k - a) \right) = a(2k + 3 - a).$$

This is maximized when $a = k$ and it gives $k^2 + 3k$.

- (A, $\{x_1\}$) Let $x = x_1$ and $y \in A$. Suppose that upper A and lower A have a and $k + 1 - a$ points of A , respectively. The only segments crossing xy are those connecting (upper A) to $C_1 \cup$ (lower A), or (lower A) to B_1 . The total is

$$\begin{aligned} a \left(\left\lfloor \frac{k}{2} \right\rfloor + (k + 1 - a) \right) + (k + 1 - a) \left(k + 2 - \left\lfloor \frac{k}{2} \right\rfloor \right) \\ = a(2 \left\lfloor \frac{k}{2} \right\rfloor - 1 - a) + (k + 1)(k + 2 - \left\lfloor \frac{k}{2} \right\rfloor). \end{aligned}$$

This is maximized when $a \in \{ \lfloor k/2 \rfloor, \lfloor k/2 \rfloor - 1 \}$ and it gives $k^2 + 3k + 2 - \lfloor k/2 \rfloor(k + 2 - \lfloor k/2 \rfloor) < k^2 + 3k + 2$.

- (A, B_1) Let $x \in B_1$ and $y \in A$. Suppose that upper A , lower A , lower B_1 , and upper B_1 have a , $k + 1 - a$, b , and $k + 1 - \lfloor k/2 \rfloor - b$ points, respectively. The only segments crossing xy are those connecting (upper A) to $C_1 \cup$ (lower A) \cup (upper B_1) $\cup \{x_1\}$, or (lower B_1) to (lower A) $\cup C_1 \cup \{x_1\}$. The total is

$$\begin{aligned} a \left(\left\lfloor \frac{k}{2} \right\rfloor + (k + 1 - a) + (k + 1 - \left\lfloor \frac{k}{2} \right\rfloor - b) + 1 \right) + b \left((k + 1 - a) + \left\lfloor \frac{k}{2} \right\rfloor + 1 \right) \\ = a(2k + 3 - 2b - a) + b(k + 2 + \left\lfloor \frac{k}{2} \right\rfloor). \end{aligned}$$

As a function of a , this is maximized when $a \in \{k + 1 - b, k + 2 - b\}$ and it gives

$$b(b - k - 1 + \left\lfloor \frac{k}{2} \right\rfloor) + (k^2 + 3k + 2).$$

This is maximized when $b \in \{0, k + 1 - \lfloor k/2 \rfloor\}$ and it gives $k^2 + 3k + 2$.

- (A, C_1) Let $x \in C_1$ and $y \in A$. Suppose that upper A , lower A , lower C_1 , and upper C_1 have a , $k + 1 - a$, c , and $\lfloor k/2 \rfloor - 1 - c$ points, respectively. The only segments crossing xy are those connecting (upper A) to (upper C_1) \cup (lower A), or (lower A) to $B_1 \cup$ (lower C_1) $\cup \{x_1\}$, or (upper C_1) to $B_1 \cup \{x_1\}$, or x_1 to $C_2 \cup \{x_2\}$. The total is

$$\begin{aligned} a \left(\left(\left\lfloor \frac{k}{2} \right\rfloor - 1 - c \right) + (k + 1 - a) \right) + (k + 1 - a) \left((k + 2 - \left\lfloor \frac{k}{2} \right\rfloor) + c + 1 \right) \\ + \left(\left\lfloor \frac{k}{2} \right\rfloor - 1 - c \right) (k + 2 - \left\lfloor \frac{k}{2} \right\rfloor + 1) + \left(\left\lfloor \frac{k}{2} \right\rfloor + 1 \right) \\ = a(2 \left\lfloor \frac{k}{2} \right\rfloor - 3 - 2c - a) + c \left(\left\lfloor \frac{k}{2} \right\rfloor - 2 \right) + k^2 + 3k + 1 + 4 \left\lfloor \frac{k}{2} \right\rfloor - \left\lfloor \frac{k}{2} \right\rfloor^2. \end{aligned}$$

As a function of a , this is maximized when $a \in \{ \lfloor k/2 \rfloor - c - 1, \lfloor k/2 \rfloor - c - 2 \}$ and it gives

$$c(c + 1 - \left\lfloor \frac{k}{2} \right\rfloor) + (k^2 + 3k + 3 + \left\lfloor \frac{k}{2} \right\rfloor).$$

This is maximized when $c \in \{0, \lfloor k/2 \rfloor - 1\}$ and it gives $k^2 + 3k + 3 + \lfloor k/2 \rfloor$.

$(\{x_1\}, \{x_2\})$ The only segments that cross x_1x_2 are those with one endpoint in A and the other in $C_1 \cup C_2$. The total is $(k+2)(2\lfloor k/2 \rfloor) \leq k^2 + 2k$.

$(\{x_1\}, B_1)$ Let $x = x_1$ and $y \in B_1$. Suppose that lower B_1 has b points and upper B_1 has $k+1 - \lfloor k/2 \rfloor - b$ points. The only segments crossing xy are those connecting lower B_1 to C_1 , or upper B_1 to A . The total is

$$b \lfloor \frac{k}{2} \rfloor + (k+1 - \lfloor \frac{k}{2} \rfloor - b)(k+2) = b(\lfloor \frac{k}{2} \rfloor - k - 2) + (k+2)(k+1 - \lfloor \frac{k}{2} \rfloor).$$

This is maximized when $b = 0$ and it gives $(k+2)(k+1 - \lfloor k/2 \rfloor) < k^2 + 3k + 2$.

$(\{x_1\}, B_2)$ Let $x = x_1$ and $y \in B_2$. Suppose that lower B_2 has b points and upper B_2 has $k+1 - \lfloor k/2 \rfloor - b$ points. The only segments crossing xy are those connecting B_1 to $C_1 \cup C_2 \cup (\text{upper } B_2) \cup \{x_2\}$, or (lower B_2) to $C_1 \cup (\text{upper } B_2)$. The total is

$$\begin{aligned} (k+2 - \lfloor \frac{k}{2} \rfloor) [\lfloor \frac{k}{2} \rfloor + \lfloor \frac{k}{2} \rfloor + (k+1 - \lfloor \frac{k}{2} \rfloor - b) + 1] + b [\lfloor \frac{k}{2} \rfloor + (k+1 - \lfloor \frac{k}{2} \rfloor - b)] \\ = b(\lfloor \frac{k}{2} \rfloor - 1 - b) + (k^2 + 4k + 4) - \lfloor \frac{k}{2} \rfloor^2 \\ < b(\lfloor \frac{k}{2} \rfloor - b) + (k^2 + 4k + 4) - \lfloor \frac{k}{2} \rfloor^2. \end{aligned}$$

This is maximized (for b real) when $b = \lfloor k/2 \rfloor / 2$ and it gives $k^2 + 4k + 4 - \frac{3}{4}\lfloor k/2 \rfloor^2$, which is at most $k^2 + 4k + 3 - \lfloor k/2 \rfloor$ for $k \geq 4$.

$(\{x_1\}, C_1)$ Let $x = x_1$ and $y \in C_1$. Suppose that lower C_1 has c points and upper C_1 has $\lfloor k/2 \rfloor - 1 - c$ points. The only segments crossing xy are those connecting upper C_1 to B_1 , or lower C_1 to A . The total is

$$(\lfloor \frac{k}{2} \rfloor - 1 - c)(k+2 - \lfloor \frac{k}{2} \rfloor) + c(k+2) = (\lfloor \frac{k}{2} \rfloor - 1)(k+2 - \lfloor \frac{k}{2} \rfloor) + c \lfloor \frac{k}{2} \rfloor.$$

This is maximized when $c = \lfloor k/2 \rfloor - 1$ and it gives $(k+2)(\lfloor k/2 \rfloor - 1) \leq k^2/2 - 2$.

$(\{x_1\}, C_2)$ Let $x = x_1$ and $y \in C_2$. Suppose that lower C_2 has c points and upper C_2 has $\lfloor k/2 \rfloor - 1 - c$ points. Note that there is a line through x_1 that separates C_1 and C_2 . Then the only segments crossing xy are those connecting C_1 to $A \cup \{x_2\}$ or (upper C_2) to $B_1 \cup C_1 \cup (\text{lower } C_2)$. The total is

$$\begin{aligned} \lfloor \frac{k}{2} \rfloor [(k+2) + 1] + (\lfloor \frac{k}{2} \rfloor - 1 - c) [(k+2 - \lfloor \frac{k}{2} \rfloor) + \lfloor \frac{k}{2} \rfloor + c] \\ = c(\lfloor \frac{k}{2} \rfloor - k - 3 - c) + (k+2)(2\lfloor \frac{k}{2} \rfloor - 1) + \lfloor \frac{k}{2} \rfloor. \end{aligned}$$

This is maximized when $c = 0$ and it gives $(k+2)(2\lfloor k/2 \rfloor - 1) + \lfloor k/2 \rfloor \leq k^2 + 3k/2 - 2$.

(B_1, B_1) Let xy be a segment with both points in B_1 . Suppose that there are b points of B_1 between x and y , and $k - \lfloor k/2 \rfloor - b$ outside. The only segments crossing xy are those connecting one of the b inside points in B_1 to points in $B_2 \cup C_2 \cup \{x_2\}$ or to points in B_1 outside xy . The total is

$$b [(k+2 - \lfloor \frac{k}{2} \rfloor) + \lfloor \frac{k}{2} \rfloor + 1 + (k - \lfloor \frac{k}{2} \rfloor - b)] = b[2k + 3 - \lfloor \frac{k}{2} \rfloor - b].$$

This is maximized when $b = k - \lfloor k/2 \rfloor$ and it gives $(k - \lfloor k/2 \rfloor)(k+3) \leq k^2 + 3k$ for $k \geq 4$.

(B_1, B_2) Let $x \in B_1$ and $y \in B_2$. Suppose that lower B_1 , upper B_1 , lower B_2 , and upper B_2 have b_1 , $k+1 - \lfloor k/2 \rfloor - b_1$, b_2 , and $k+1 - \lfloor k/2 \rfloor - b_2$ points, respectively. The only segments crossing xy are those connecting (lower B_1) to (upper B_2) \cup (upper B_1) $\cup C_2 \cup \{x_2\}$, or (lower B_2) to (upper B_1) \cup (upper B_2) $\cup C_1 \cup \{x_1\}$. The total is

$$\begin{aligned} & b_1[(k+1 - \lfloor \frac{k}{2} \rfloor - b_2) + (k+1 - \lfloor \frac{k}{2} \rfloor - b_1) + \lfloor \frac{k}{2} \rfloor + 1] \\ & + b_2[(k+1 - \lfloor \frac{k}{2} \rfloor - b_1) + (k+1 - \lfloor \frac{k}{2} \rfloor - b_2) + \lfloor \frac{k}{2} \rfloor + 1] \\ & = (b_1 + b_2)(2k+3 - \lfloor \frac{k}{2} \rfloor - (b_1 + b_2)) \\ & < (b_1 + b_2)(2k+3 - (b_1 + b_2)). \end{aligned}$$

As a function of $b_1 + b_2$, this is maximized when $b_1 + b_2 \in \{k+1, k+2\}$ and it gives $k^2 + 3k + 2$.

(B_1, C_1) Let $x \in B_1$ and $y \in C_1$. Suppose that lower B_1 , upper B_1 , lower C_1 , and upper C_1 have b , $k+1 - \lfloor k/2 \rfloor - b$, $\lfloor k/2 \rfloor - 1 - c$, and c points, respectively. The only segments crossing xy are those connecting (lower C_1) to (lower B_1) $\cup A \cup \{x_1\}$, or (upper B_1) to (upper C_1) $\cup A \cup \{x_1\}$, or x_1 to B_2 . The total is

$$\begin{aligned} & (\lfloor \frac{k}{2} \rfloor - 1 - c)[b + (k+2) + 1] + (k+1 - \lfloor \frac{k}{2} \rfloor - b)[c + (k+2) + 1] + (k+2 - \lfloor \frac{k}{2} \rfloor) \\ & = -b(2c+4+k - \lfloor \frac{k}{2} \rfloor) - c(2 + \lfloor \frac{k}{2} \rfloor) + k^2 + 4k + 2 - \lfloor \frac{k}{2} \rfloor. \end{aligned}$$

As a function of b and c , this is maximized when $b = c = 0$ and it gives $k^2 + 4k + 2 - \lfloor k/2 \rfloor$.

(B_1, C_2) Let $x \in B_1$ and $y \in C_2$. Suppose that lower B_1 , upper B_1 , lower C_2 , and upper C_2 have b , $k+1 - \lfloor k/2 \rfloor - b$, $\lfloor k/2 \rfloor - 1 - c$, and c points, respectively. The only segments crossing xy are those connecting (lower B_1) to (upper C_2) \cup (upper B_1), or (upper B_1) to $B_2 \cup$ (lower C_2) $\cup \{x_2\}$, or $B_2 \cup$ (lower C_2) to $C_1 \cup \{x_1\}$, or (upper C_2) to (lower C_2). The total is

$$\begin{aligned} & b[c + (k+1 - \lfloor \frac{k}{2} \rfloor - b)] + (k+1 - \lfloor \frac{k}{2} \rfloor - b)[(k+2 - \lfloor \frac{k}{2} \rfloor) + (\lfloor \frac{k}{2} \rfloor - 1 - c) + 1] \\ & + [(k+2 - \lfloor \frac{k}{2} \rfloor) + (\lfloor \frac{k}{2} \rfloor - 1 - c)](\lfloor \frac{k}{2} \rfloor + 1) + c(\lfloor \frac{k}{2} \rfloor - 1 - c) \\ & = -(b-c)^2 - b(\lfloor \frac{k}{2} \rfloor + 1) - c(k - \lfloor \frac{k}{2} \rfloor + 3) + k^2 + 4k + 3 - \lfloor \frac{k}{2} \rfloor. \end{aligned}$$

As a function of b and c , this is maximized when $b = c = 0$ and it gives $k^2 + 4k + 3 - \lfloor k/2 \rfloor$.

(C_1, C_1) Let xy be a segment with both points in C_1 . Suppose that there are c points of C_1 between x and y , and $\lfloor k/2 \rfloor - 2 - c$ outside. The only segments crossing xy are those connecting one of the c inside points in C_1 to points in $B_2 \cup C_2 \cup \{x_2\}$ or to points in C_1 outside xy . The total is

$$c[(k+2 - \lfloor \frac{k}{2} \rfloor) + \lfloor \frac{k}{2} \rfloor + 1 + (\lfloor \frac{k}{2} \rfloor - 2 - c)] = c[k + \lfloor \frac{k}{2} \rfloor + 1 - c].$$

This is maximized when $c = \lfloor k/2 \rfloor - 2$ and it gives $(k+3)(\lfloor k/2 \rfloor - 2) < k^2/2$.

(C_1, C_2) Let $x \in C_1$ and $y \in C_2$. Suppose that lower C_1 , upper C_1 , lower C_2 , and upper C_2 have $\lfloor k/2 \rfloor - 1 - c_1$, c_1 , $\lfloor k/2 \rfloor - 1 - c_2$, and c_2 points, respectively. The only segments crossing xy are those connecting (upper C_1) to $B_2 \cup (\text{lower } C_2) \cup (\text{lower } C_1)$, or (upper C_2) to $B_1 \cup (\text{lower } C_1) \cup (\text{lower } C_2)$, or x_1 to (lower C_2), or x_2 to (lower C_1). The total is

$$\begin{aligned}
& c_1[(k + 2 - \lfloor \frac{k}{2} \rfloor) + (\lfloor \frac{k}{2} \rfloor - 1 - c_2) + (\lfloor \frac{k}{2} \rfloor - 1 - c_1)] \\
& + c_2[(k + 2 - \lfloor \frac{k}{2} \rfloor) + (\lfloor \frac{k}{2} \rfloor - 1 - c_1) + (\lfloor \frac{k}{2} \rfloor - 1 - c_2)] \\
& + (\lfloor \frac{k}{2} \rfloor - 1 - c_1) + (\lfloor \frac{k}{2} \rfloor - 1 - c_2) \\
& = (c_1 + c_2)(k + \lfloor \frac{k}{2} \rfloor - 1 - c_1 - c_2) + 2 \lfloor \frac{k}{2} \rfloor - 2 \\
& < (c_1 + c_2)[2k - (c_1 + c_2)] + k - 2.
\end{aligned}$$

As a function of $c_1 + c_2$, this is maximized when $c_1 + c_2 = k$ and it gives $k^2 + k - 2$.

In all cases the maximum possible number of crossings is at most $k^2 + 4k + 3 - \lfloor k/2 \rfloor$. Equality occurs in the case (B_1, C_2) when $b = c = 0$, and for k even in the case (A, C_1) when $(a, c) \in \{(0, k/2 - 1), (k/2 - 1, 0), (k/2 - 2, 0)\}$; together with the symmetric cases (B_2, C_1) and (A, C_2) . \square

5 Proof of Theorem 1

If $n \leq 2$, then the result is trivially true because $\overline{\text{lcr}}(K_n) = 0$. Suppose $n \geq 3$. The required formula as lower bound follows directly by Theorem 3. If $n \equiv 0, 1 \pmod{3}$, then the matching upper bound follows from Theorem 4. If $n \equiv 2 \pmod{3}$ and $n \geq 20$, then the matching upper bound follows from Theorem 5. The upper bounds $\overline{\text{lcr}}(K_5) \leq 1$ and $\overline{\text{lcr}}(K_8) \leq 4$ follow from Theorem 4. The drawings shown in Figure 5 show that $\overline{\text{lcr}}(K_{11}) \leq 8$, $\overline{\text{lcr}}(K_{14}) \leq 15$, and $\overline{\text{lcr}}(K_{17}) \leq 23$. Finally, for $n = 8$, we used the data base by Aichholzer et al. [7, 6] to calculate the local crossing number of each of the 3315 order types of 8 points. Our calculations verified that $\overline{\text{lcr}}(K_8) \geq 4$ for all of them, with equality achieved by 39 order types. For $n = 14$, Aichholzer [5] extended his data base from $n = 11$ to $n = 14$ for this specific problem. He confirmed our results for $n < 14$ and verified our earlier conjecture [1] that there are no geometric sets of 14 points where every edge is crossed at most 14 times. Thus $\overline{\text{lcr}}(K_{14}) = 15$.

6 Final Remarks

The construction in Theorem 3 shows that Lemma 2 is best possible. That is, the point-sets P constructed in Theorem 3 verify that every path of length 2 whose endpoints are vertices of the convex hull of P separates the rest of the set into parts that differ by no less than $(n - 3)/3$.

Another consequence of Theorem 1 is that for $n \notin \{8, 14\}$, there is always an edge achieving $\overline{\text{lcr}}(K_n)$ with one endpoint that is a vertex of the convex hull. The same is true for $n = 8$ by inspecting all order types that achieve $\overline{\text{lcr}}(K_8) = 4$. However, it is not true that




Figure 5: The construction for $n \in \{11, 14, 17\}$. The 11-point set consists of the black points, the 14-point set consists of the black or white points, and the 17-point set consists of all of the points shown.

all of the maximal edges on the optimal examples have an endpoint that is a vertex of the convex hull. Another property shared by all the optimal constructions we know is that their convex hull is a triangle. Is it true that all the optimal point sets have triangular convex hulls? A related problem is to determine the maximum number of edges that can achieve $\overline{\text{lcr}}(K_n)$ crossings in an optimal point set.

Acknowledgments. We warmly thank the referees for their comments and suggestions, which certainly improved the presentation of this paper. We also thank Oswin Aichholzer for listening to our results, settling the case $n = 14$, and verifying Theorem 1 for small values of n . S. Fernández-Merchant’s research was supported by the NSF grant DMS-1400653.

References

- [1] B. M. Ábrego and S. Fernández-Merchant. The rectilinear local crossing number of K_n . [arXiv:1508.07926v1 \[math.CO\]](#), 2015.
- [2] B. M. Ábrego and S. Fernández-Merchant. Geometric drawings of K_n with few crossings. *J. Combin. Theory Ser. A* 114:373–379, 2007.
- [3] B. M. Ábrego, M. Cetina, S. Fernández-Merchant, J. Leaños, and G. Salazar. On $(\leq k)$ -edges, crossings, and halving lines of geometric drawings of K_n . *Discrete Comput. Geom.*, 48:192–215, 2012.
- [4] B. M. Ábrego, M. Cetina, S. Fernández-Merchant, J. Leaños, and G. Salazar. 3-symmetric and 3-decomposable drawings of K_n , *Discrete Appl. Math.*, 158:1240–1258, 2010.

- [5] O. Aichholzer, personal communication, 2016.
- [6] O. Aichholzer, Enumerating order types for small point sets with applications, <http://www.ist.tugraz.at/staff/aichholzer/research/rp/triangulations/order/types/>.
- [7] O. Aichholzer, F. Aurenhammer, and H. Krasser, Enumerating order types for small point sets with applications. *Order* 19:265–281, 2002.
- [8] E. de Klerk, D. V. Pasechnik, and A. Schrijver. Reduction of symmetric semidefinite programs using the regular *-representation. *Math. Program*, 109:613–624, 2007.
- [9] R. K. Guy. A combinatorial problem. *Bull. Malayan Math. Soc.*, 7:68–72, 1960.
- [10] R. K. Guy, T. Jenkyns, and J. Schaer. The toroidal crossing number of the complete graph. *J. Combinatorial Theory*, 4:376–390, 1968.
- [11] P. C. Kainen. Thickness and coarseness of graphs. *Abh. Math. Sem. Univ. Hamburg*, 39:88–95, 1973.
- [12] D. Lara. Problem posed at the 2nd Austrian-Japanese-Mexican-Spanish Workshop on Discrete Geometry, Barcelona, Spain, July 6–10, 2015.
- [13] D. Lara, C. Rubio-Montiel, F. Zaragoza. Grundy and pseudo-Grundy indices for complete graphs. In *Abstracts of the XVI Spanish Meeting on Computational Geometry*, Barcelona, July 1–3, 2015.
- [14] M. Schaefer. The Graph Crossing Number and its Variants: A Survey. *Electron. J. Combin.* DS21: May 15, 2014.
- [15] C. Thomassen. Rectilinear drawings of graphs. *J. Graph Theory*, 12(3):335–341, 1988.