

Two integral representations for the logarithm of the Glaisher-Kinkelin constant

Jean-Christophe PAIN^{1,2,*}

¹CEA, DAM, DIF, F-91297 Arpajon, France

²Université Paris-Saclay, CEA, Laboratoire Matière en Conditions Extrêmes,
91680 Bruyères-le-Châtel, France

May 10, 2024

Abstract

We present two integral representations of the logarithm of the Glaisher-Kinkelin constant. Both are based on a definite integral of $\ln [\Gamma(x+1)]$, Γ being the usual Gamma function. The first one relies on an integral representation of $\ln [\Gamma(x+1)]$ due to Binet, and the second one results from the so-called Malmstén formula. The numerical evaluation is easier with the latter expression than with the former.

1 Introduction

The Glaisher-Kinkelin constant A is related to the Barnes G -function:

$$G(n) = \prod_{k=1}^{n-2} k! \quad (1)$$

by

$$A = \lim_{n \rightarrow \infty} \frac{(2\pi)^{\frac{n}{2}} n^{\frac{n^2}{2} - \frac{1}{12}} e^{-\frac{3n^2}{4} + \frac{1}{12}}}{G(n+1)}. \quad (2)$$

The logarithm of the Glaisher-Kinkelin constant A can be expressed as the following integral [1, 2]:

$$\ln A = \frac{1}{12} - 2 \int_0^\infty \frac{x \ln x}{e^{2\pi x} - 1} dx. \quad (3)$$

Another integral is given by [1]:

$$\int_0^{1/2} \ln [\Gamma(x+1)] dx = -\frac{1}{2} - \frac{7}{24} \ln 2 + \frac{1}{4} \ln \pi + \frac{3}{2} \ln A, \quad (4)$$

*jean-christophe.pain@cea.fr

where Γ represents the usual Gamma function. Only a few integral representations are available in the literature (see for instance the work of Choi and Nash [3]). Using two different improper integral representations of $\ln [\Gamma(x+1)]$ (the first one due to Binet (see Sec. 2), and the second one to Malmstén (see Sec. 3)), we derive two integral representations of the logarithm of the Glaisher-Kinkelin constant $\ln A$, which, to our knowledge, were not published elsewhere.

2 Integral representation deduced from the Binet formula

The following important result is due to Binet [4, 5]:

$$\Gamma(x+1) = \left(\frac{x}{e}\right)^x \sqrt{2\pi x} e^{\theta(x)}, \quad (5)$$

where

$$\theta(x) = \int_0^\infty \left(\frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) \frac{e^{-xt}}{t} dt. \quad (6)$$

One has

$$\ln [\Gamma(x+1)] = x \ln x - x + \frac{1}{2} \ln(2\pi x) + \theta(x), \quad (7)$$

and

$$\int_0^{1/2} \ln [\Gamma(x+1)] dx = \int_0^{1/2} (x \ln x - x) dx + \frac{1}{2} \int_0^{1/2} \ln(2\pi x) dx + \int_0^{1/2} \theta(x) dx. \quad (8)$$

The first integral is equal to

$$\int_0^{1/2} (x \ln x - x) dx = -\frac{1}{8} \left(\ln 2 + \frac{3}{2} \right), \quad (9)$$

the second one is

$$\int_0^{1/2} \ln(2\pi x) dx = \frac{1}{2} (\ln \pi - 1), \quad (10)$$

and the third one, applying the Fubini theorem, reads

$$\int_0^{1/2} \theta(x) dx = \int_0^\infty \left(\frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) \frac{1}{t} \left(\int_0^{1/2} \frac{e^{-xt}}{t} dx \right) dt, \quad (11)$$

which is equal to

$$\int_0^{1/2} \theta(x) dx = \int_0^\infty \left(\frac{1}{e^t - 1} - \frac{1}{t} + \frac{1}{2} \right) \frac{(1 - e^{-t/2})}{t^2} dt, \quad (12)$$

or also

$$\int_0^{1/2} \theta(x) dx = \int_0^\infty \frac{(1 - e^{-t/2}) [t \coth(t/2) - 2]}{2t^3} dt. \quad (13)$$

Inserting Eqs. (9), (10) and (13) in Eq. (4) yields

$$-\frac{1}{2} - \frac{7}{24} \ln 2 + \frac{1}{4} \ln \pi + \frac{3}{2} \ln A = -\frac{1}{8} \left(\ln 2 + \frac{3}{2} \right) + \frac{1}{4} (\ln \pi - 1) + \int_0^\infty \frac{(1 - e^{-t/2}) [t \coth(t/2) - 2]}{2t^3} dt. \quad (14)$$

or equivalently

$$\ln A = \frac{1}{9} \ln 2 + \frac{1}{24} + \frac{2}{3} \int_0^\infty \frac{(1 - e^{-t/2}) [t \coth(t/2) - 2]}{2t^3} dt, \quad (15)$$

which is the first main result of the present work.

3 Integral representation resulting from the Malmstén formula

The Malmstén formula reads [6]:

$$\ln \Gamma(z + 1) = \int_0^\infty \left[z - \frac{(1 - e^{-zt})}{(1 - e^{-t})} \right] \frac{e^{-t}}{t} dt. \quad (16)$$

Using the Fubini theorem, one can write

$$\int_0^{1/2} \ln [\Gamma(x + 1)] dx = \int_0^\infty \frac{e^{-t}}{t} \left(\int_0^{1/2} \left[x - \frac{(1 - e^{-xt})}{(1 - e^{-t})} \right] dx \right) dt, \quad (17)$$

which is equal to

$$\int_0^{1/2} \ln [\Gamma(x + 1)] dx = \int_0^\infty \left[\frac{1}{8} - \frac{1}{2(1 - e^{-t})} + \frac{1}{t(1 + e^{-t/2})} \right] \frac{e^{-t}}{t} dt, \quad (18)$$

or equivalently

$$\int_0^{1/2} \ln [\Gamma(x + 1)] dx = \int_0^\infty \frac{e^{-t} [(8 - 3t) e^t - 8 e^{t/2} - t]}{8t^2(e^t - 1)} dt. \quad (19)$$

Inserting the latter expression in Eq. (4) yields

$$-\frac{1}{2} - \frac{7}{24} \ln 2 + \frac{1}{4} \ln \pi + \frac{3}{2} \ln A = \int_0^{1/2} \ln [\Gamma(x + 1)] dx = \int_0^\infty \frac{e^{-t} [(8 - 3t) e^t - 8 e^{t/2} - t]}{8t^2(e^t - 1)} dt, \quad (20)$$

leading to

$$\ln A = \frac{1}{3} + \frac{7}{36} \ln 2 - \frac{1}{6} \ln \pi + \frac{2}{3} \int_0^\infty \frac{e^{-t} [(8 - 3t) e^t - 8 e^{t/2} - t]}{8t^2(e^t - 1)} dt, \quad (21)$$

which is the second main result of the present work. The convergence of the latter integral (Eq. (21)), is much faster than for the representation (15).

4 Conclusion

In this letter, we presented two integral representations of the logarithm of the Glaisher-Kinkelin constant. Both are based on a definite integral representation involving $\ln [\Gamma(x + 1)]$, which can itself be expressed by an improper integral, either using a representation proposed a long time ago by Binet, or using the Malmstén formula. The two new expressions may help investigating new properties of the Glaisher-Kinkelin constant.

References

- [1] J. W. L. Glaisher, *On the product $1^1.2^2.3^3\dots n^n$* , Messenger Math. **7**, 43-47 (1878).
- [2] G. Almkvist, *Asymptotic formulas and generalized Dedekind sums*, Experim. Math. **7**, 343-359 (1998).
- [3] J. Choi and C. Nash, *Integral representations of the Kinkelin's constant A* , Math. Japon. **45**, 223-230 (1997).
- [4] Z. Sasvari, *An elementary proof of Binet's formula for the Gamma function*, Amer. Math. Mon. **106**, 156-158 (1999).
- [5] E. T. Whittaker and G. N. Watson, *A course in modern analysis*, 4th ed. Cambridge, England: Cambridge University Press, 1990.
- [6] A. Erdélyi, W. Magnus, F. Oberhettinger and F. G. Tricomi, *Higher transcendental functions*, Vol. 1. New York: Krieger, pp. 20-21, 1981.