Widom's conjecture: variance asymptotics and entropy bounds for counting statistics of free fermions

Alix Deleporte, Gaultier Lambert

Université Paris-Saclay, CNRS, Laboratoire de mathématiques d'Orsay, 91405, Orsay, France. KTH Royal Institute of Technology, Department of Mathematics, 11428, Stockholm, Sweden.

May 14, 2024

Abstract

We obtain a central limit theorem for bulk counting statistics of free fermions in smooth domains of \mathbb{R}^n with an explicit description of the covariance structure. This amounts to a study of the asymptotics of norms of commutators between spectral projectors of semiclassical Schrödinger operators and indicator functions supported in the bulk. In the spirit of the Widom conjecture, we show that the squared Hilbert-Schmidt norm of these commutators is of order $\hbar^{-n+1}\log(\hbar)$ as the semiclassical parameter \hbar tends to 0. We also give a new upper bound on the trace norm of these commutators and applications to estimations of the entanglement entropy for free fermions.

Contents

L	Intr	oduction	
	1.1	Statement of results	4
	1.2	Organisational remarks	6
	1.3	Relation to random matrices and determinants in the Fisher-Hartwig class	6
	1.4	Widom's conjecture & entanglement entropy	8
	1.5	Acknowledgements	9
2	Pse	udodifferential operators and Fourier Integral operators	9
	2.1	Notation	9
	2.2	Pseudodifferential calculus	10
	2.3	Preliminary commutator estimates	11
3	Hill	pert-Schmidt estimates for mild spectral functions	15
	3.1	Hilbert-Schmidt norm of products; proof of (3.1)	16
	3.2	Hilbert-Schmidt norm of commutators; proof of (3.2)–(3.3) at small spectral scales	18
	3.3	Hilbert-Schmidt norm of commutators; proof of (3.2) – (3.3) at large spectral scales	20

^{*}alix.deleporte@universite-paris-saclay.fr

[†]glambert@kth.se

4	Hill	bert-Schimdt norm of commutators: Proof of Theorem 1.	22
	4.1	Preliminary estimates	24
	4.2	Reduction to an oscillatory integral	28
		4.2.1 Step 1: Stationary phase	30
		4.2.2 Step 2: Study of the phase Ψ_1	31
		4.2.3 Step 3: the main symbol is in the class \mathcal{F}	33
		4.2.4 Step 4: Control of $I_{1,\hbar}$	35
	4.3	Case of a contractible open set	37
		4.3.1 Change of variables	38
		4.3.2 Asymptotics in case of the ball	40
		4.3.3 Proof of Proposition 4.12	44
	4.4		
	4.5	Central limit theorem; Proof of Theorem 2	47
5	Tra	ce norm of commutators & entropy estimates	49
	5.1	Proof of Theorem 4: Multi-scale argument	49
	5.2	Bounds for the entanglement entropy: Proof of Theorem $3 \dots \dots \dots \dots \dots$.	52
\mathbf{A}	Stat	tionary phase with mild amplitudes	54

1 Introduction

The goal of this article is to explore the relationships between the number variance and entanglement entropy of certain free fermionic systems, on one hand, and the spectral theory of semiclassical Schrödinger operators on the other hand. Consider a Schrödinger operator on $L^2(\mathbb{R}^n)$:

$$H_{\hbar} := -\hbar^2 \Delta + V,$$

where Δ is the standard (negative) Laplacian on \mathbb{R}^n , $\hbar > 0$ plays the role of the Planck constant, and $V : \mathbb{R}^n \to \mathbb{R}$ is bounded from below. The operator H_{\hbar} is essentially self-adjoint with domain

$$H^2(\mathbb{R}^n) \cap \{u \in L^2, Vu \in L^2\}.$$

We are interested in the fluctuations of the N particles free fermionic state associated with the operator H_{\hbar} at zero temperature. Under assumptions on V and N which are specified below, this (random) point process is described by the probability density \mathbb{P}_N on $\mathbb{R}^{n\times N}$ with density

$$\mathbb{P}_N[\mathrm{d}\mathbf{x}] = \frac{1}{N!} \left| \det_{N \times N} \left[v_k(x_j) \right] \right|^2, \qquad (x_1, \dots, x_N) \in \mathbb{R}^{n \times N}, \tag{1.1}$$

where $(\lambda_k, v_k)_{1 \leq k \leq N}$ are the N lowest eigenvalues, and associated orthonormal eigenfunctions, of H_{\hbar} . Even though it is not emphasized, this spectral data and the measure \mathbb{P}_N depend on the semiclassical parameter \hbar . We are interested in a joint limit where $\hbar \to 0$ and $N \to +\infty$, while keeping fixed the Fermi energy μ . To this end, we fix $\mu \in \mathbb{R}$ and choose

$$N = N(\hbar) := \max \{ k \in \mathbb{N} : \lambda_k(\hbar) \le \mu \}.$$

We work under the following assumptions on the external potential V:

Assumptions 1. Fix $\mu \in \mathbb{R}$ and assume that $V \in L^1_{loc}(\mathbb{R}^n)$ with $-\infty < \inf(V) < \mu$. We assume that $\mathcal{D} := \{V < \mu\}$ is relatively compact, V is C^{∞} on a neighborhood of \mathcal{D} , and $\partial_x V(x) \neq 0$ for all $x \in \partial \mathcal{D} = \{V = \mu\}$.

The probability measure (1.1) gives rise to a determinantal point process $\mathbf{X} := \sum_{j=1}^{N} \boldsymbol{\delta}_{x_j}$ on \mathbb{R}^n associated with the integral kernel (also denoted $\Pi_{\hbar,\mu}$) of the spectral projection

$$\Pi_{\hbar,\mu} = \mathbb{1}_{(-\infty,\mu]}(H_{\hbar}). \tag{1.2}$$

The set \mathcal{D} is usually called the *bulk* or *droplet*, and it is the region where the fermions concentrate. This framework and our assumptions are similar to that of our previous work [19] where we study the scaling limits of the kernel Π both in the bulk and at the boundary of the droplet (see also [14]) and give applications to the (real) random variables $\mathbf{X}(\mathbf{f}) = \sum_{j=1}^{N} \mathbf{f}(x_j)$ for certain smooth functions \mathbf{f} .

The determinantal structure means that the correlation functions or marginals of the measure (1.1) are of the form

$$\rho_{N,k}(x_1,\cdots,x_k) = \det_{k \times k} \left[\Pi_{\hbar,\mu}(x_i,x_j) \right], \qquad k \le N.$$

Moreover, for any $f \in L^{\infty}(\mathbb{R}^n)$, the random variable $\mathbf{X}(f) = \sum_{j=1}^{N} f(x_j)$ has the following Laplace transform:

$$\mathbb{E}[\exp \mathbf{X}(\mathbf{f})] = \det(1 + (e^{\mathbf{f}} - 1)\Pi_{\hbar,\mu}) \tag{1.3}$$

where the right-hand side is a Fredholm determinant on $L^2(\mathbb{R}^n)$, since $\Pi_{\hbar,\mu}$ is a finite-rank projection with $\operatorname{tr} \Pi_{\hbar,\mu} = N(\hbar)$.

In [19], we showed that with overwhelming probability, as $\hbar \to 0$, one has weak-* convergence of probability measures

$$N^{-1}\mathbf{X} \to \mathrm{d}\varrho := Z^{-1}(\mu - V)^{\frac{n}{2}}_{\perp} \mathrm{d}x,$$
 (1.4)

where dx denotes the Lebesgue measure on \mathbb{R}^n , and

$$Z = \int (\mu - V)_{+}^{\frac{n}{2}} dx$$

$$N(\hbar) \sim \frac{|B_n|Z}{(2\pi\hbar)^n} \quad \text{as } \hbar \to 0,$$
(1.5)

where $|B_n|$ is the volume of the Euclidean ball $B_n = \{x \in \mathbb{R}^n : |x| \leq 1\}$. The convergence (1.4) yields a law of large numbers for counting statistics, in the sense that $\mathbf{X}(\mathbf{f})$ concentrates around its mean $N \int \mathrm{fd}\rho$ for $\mathbf{f} \in C_c^{\infty}(\mathbb{R}^n)$. In [18, 19], we also studied the fluctuations of $\mathbf{X}(\mathbf{f})$ and we obtain a central limit theorem as $\hbar \to 0$ with the variance being at most of order $\hbar^{-n+1} \approx N\hbar$. Moreover, in dimension 1, [57, 18] the variance converges and its limit is described by a weighted $H^{1/2}$ Sobolev seminorm.

This article focuses instead on the case where $f = \mathbb{1}_{\Omega}$ is the indicator function of an open, relatively compact subset Ω of the droplet \mathcal{D} with a smooth boundary. The *counting statistics* $\mathbf{X}(\Omega)$ are directly relevant for physical applications; in particular the problem of measuring the *entanglement entropy* of subsystems that we shortly review.

Entanglement is a crucial property of quantum states which, for free fermions, results from the Pauli exclusion principle, and is quantified by suitable concepts of entropy. For any open, relatively compact set $\Omega \in \mathbb{R}^n$, one can define its von Neumann entanglement entropy \mathcal{S}_{Ω} which measures the correlations between the state restricted to Ω and its complement. This quantity plays a crucial role in condensed matter physics and in quantum information, however it is difficult to estimate, both experimentally and theoretically. Remarkably, at zero temperature, the entanglement entropy is not extensive and is often of order $\hbar^{n-1}|\partial\Omega|$ where \hbar is the typical inter-particle distance. This is known as the area law in the physics literature; see [64, 8] for an historic references. This property is expected for states with a finite correlation length (gapped systems), while for gapless systems, which exhibit long-range correlations, it is expected that the entropy is enhanced by an extra $\log(\hbar^{-1})$ factor. This has been demonstrated for critical quantum

systems [5, 38, 21]; for one-dimensional models, conformal field theory methods predict that the leading behavior of the entanglement entropy is universal and

$$S_{\Omega} \sim \frac{c}{3} \log(\hbar^{-1})$$
 where c is the central charge of the corresponding CFT. (1.6)

In dimension n > 1, physical arguments also predict that, for a smooth domain Ω , the entropy \mathcal{S}_{Ω} is of order $\hbar^{1-n}\log(\hbar^{-1})$. This problem has been investigated for gapless systems of free fermions, both numerically [67] and analytically [28, 7, 6], providing many evidence for this *enhanced area law*. Remarkably, these works predict a universal relationship between the fluctuations of the counting statistic $\mathbf{X}(\Omega)$ and the entanglement entropy, independent of the dimension:

$$S_{\Omega} \underset{N \to +\infty}{\sim} \frac{\pi^2}{3} \text{var} \mathbf{X}(\Omega).$$
 (1.7)

This relation provides an experimental mean to measure the entanglement entropy of a fermionic system by estimating its quantum fluctuations, which is easier to measure.

Apart from the case of constant coefficients differential operators, there is no rigorous bound in the literature on the entanglement entropy. The goal of this article is to rigorously explore these questions for the (Schrödinger) free Fermion model defined above by using techniques from semiclassical analysis. In particular, we obtain an equivalent for $\operatorname{var}\mathbf{X}(\Omega)$ which is consistent with Widom's conjecture (Theorem 1) and deduce a central limit theorem for $\mathbf{X}(\Omega)$ (Theorem 2). We also obtain lower and upper bounds for the entanglement entropy (Theorem 3) which match up to $\log \log(\hbar^{-1})$.

1.1 Statement of results

We first compute an equivalent for the variance $\operatorname{var}\mathbf{X}(\Omega)$ for an open set $\Omega \in \mathcal{D}$ with smooth boundary, also known as *number variance* in random matrix theory (Section 1.3). By the determinantal structure, this variance corresponds to the Hilbert-Schmidt norm (J²-norm – see [56] for some background on Schatten norms) of the commutator $[\Pi_{\hbar,\mu}, \mathbb{1}_{\Omega}]$.

Theorem 1. Suppose that Assumptions 1 hold. Let $\Omega \in \mathcal{D}$ be an open set with a smooth boundary and let $f \in C^{\infty}(\mathbb{R}^n)$. Then as $\hbar \to 0$,

$$\operatorname{var} \mathbf{X}(f|_{\Omega}) = \frac{1}{2} \| [f|_{\Omega}, \Pi_{\hbar, \mu}] \|_{\mathbf{J}^{2}}^{2} = (2\pi\hbar)^{1-n} (C_{\Omega, f} \log(\hbar^{-1}) + \mathcal{O}(1))$$

where

$$C_{\Omega,f} = \frac{c_{n-1}}{2\pi^2} \int_{\partial\Omega} (\mu - V(\hat{x}))_+^{\frac{n-1}{2}} f(\hat{x})^2 d\hat{x}, \tag{1.8}$$

 $d\hat{x}$ is the volume measure on the boundary $\partial\Omega$, and $c_n = \frac{\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2}+1)}$ for $n \geq 0$ (in particular $c_n = |B_n|$ for $n \geq 1$).

We expect that these asymptotics also hold for arbitrary sets $\Omega \in \mathbb{R}^n$ with a piecewise smooth boundary. An important consequence of Theorem 1 is a central limit theorem for counting statistics.

Theorem 2. Suppose that Assumptions 1 hold. Let $k \in \mathbb{N}$, $\Omega_1, \dots, \Omega_k \in \mathcal{D}$ be a collection of open sets with smooth boundaries. Let $C_{\Omega} = C_{\Omega,1}$ be as in (1.8). Assume that for all $1 \leq i, j \leq k$, if $i \neq j$, $|\partial \Omega_i \cap \partial \Omega_j| = 0$ for the (n-1)-Haussdorff measure. Then, one has in distribution as $\hbar \to 0$,

$$\left(\frac{\mathbf{X}(\Omega_1) - \mathbb{E}[\mathbf{X}(\Omega_1)]}{\sqrt{(2\pi\hbar)^{1-n}\log(\hbar^{-1})}}, \cdots, \frac{\mathbf{X}(\Omega_k) - \mathbb{E}[\mathbf{X}(\Omega_k)]}{\sqrt{(2\pi\hbar)^{1-n}\log(\hbar^{-1})}}\right) \Rightarrow \mathcal{N}_{0,\Sigma}$$

where the limit covariance matrix is

$$\Sigma = \operatorname{diag}(C_{\Omega_1}, \cdots, C_{\Omega_k}).$$

Physical systems whose number variance are lower-order compared to the volume (or expected number of points) are called hyperuniform. This concept has been introduced in the theoretical chemistry and statistical physics literature [66] where it provides a framework to classify quasicrystals and other disordered systems; we refer to the survey [65] for examples of hyperuniform point processes and their properties. in the framework of [66, 65], Theorem 1 shows that the free fermion ground states are Class II hyperuniform point processes.

An important physical quantity related to $\mathbf{X}(\Omega)$ is the entanglement entropy defined as follows. Let $s: \lambda \in [0,1] \mapsto -\lambda \log(\lambda) - (1-\lambda) \log(1-\lambda)$ – this function is continuous with s(0) = s(1) = 0, and $s(\lambda)$ is the entropy of a Bernoulli random variable \mathbf{B}_{λ} with parameter λ .

Given $\Omega \subseteq \mathbb{R}^n$, we now define

$$S_{\Omega}(\mathbf{X}) = \operatorname{tr} s(\Pi|_{\Omega}) = \sum_{n \in \mathbb{N}} s(\sigma_n), \tag{1.9}$$

where σ_n are the eigenvalues of operator $\Pi|_{\Omega} = \mathbb{1}_{\Omega}\Pi_{\hbar,\mu}\mathbb{1}_{\Omega}$. In particular, $0 \leq \Pi|_{\Omega} \leq 1$ as operators so that $\sigma_n \in [0,1]$. Formula (1.9) comes from the fact that $\mathbf{X}(\Omega) \stackrel{\text{law}}{=} \sum_{n \in \mathbb{N}} \mathbf{B}_{\sigma_n}$ where $(\mathbf{B}_{\sigma_n})_{n \in \mathbb{N}}$ are independent Bernoulli random variables, so that there entropies sum up. This key observation is a consequence of Wick's Theorem for free fermions, e.g. [54, 55] or [32] for a probabilistic interpretation. Moreover, (1.9) agrees with the physical definition of the entanglement entropy of the reduced state $\rho_{\Omega} = \operatorname{tr}_{\mathcal{H}(\Omega^c)}(\Pi_{\hbar,\mu})$ which comes from the decomposition of the fermionic Fock space $\mathcal{H}(\mathbb{R}^n) = \mathcal{H}(\Omega) \otimes \mathcal{H}(\Omega^c)$ where $\mathcal{H}(\Omega) = \bigwedge L^2(\Omega)$. Indeed, using the canonical decomposition of the state $\Pi_{\hbar,\mu}$, one verifies that $\operatorname{tr}(\rho_{\Omega} \log \rho_{\Omega}) = \sum_{n \in \mathbb{N}} s(\sigma_n)$: see the introduction of [27] or [9, Section 7] for a detailed computation.

A natural observation is that the entropy is bounded from below by the variance, since $s(x) \ge x(1-x)$. Hence, Theorem 1 yields a lower-bound for the entanglement entropy of Ω . We prove a complementary upper bound of the same order up to a factor $\log \log (\hbar^{-1})$.

Theorem 3. Suppose that Assumptions 1 hold. Let $\Omega \in \mathcal{D}$ be an open set with a smooth boundary, there exists constants C_{Ω} , $c_{\Omega} > 0$ so that

$$c_{\Omega}\hbar^{1-n}\log(\hbar^{-1}) \le \mathcal{S}_{\Omega} \le C_{\Omega}\hbar^{1-n}\log(\hbar^{-1})\log\log(\hbar^{-1}).$$

The upper estimate is proved by interpolation between Theorem 1 and the following estimate on the trace-norm $(J^1$ -norm) of a commutator.

Theorem 4. Suppose that Assumptions 1 hold. Let $\Omega \in \mathcal{D}$ be an open set with a smooth boundary. Then as $\hbar \to 0$,

$$\|[\Pi_{\hbar,\mu}, \mathbb{1}_{\Omega}]\|_{J^1} = \mathcal{O}(\hbar^{1-n} \log^2 \hbar).$$

From the Widom conjecture (Conjecture 1 below), we expect that as $\hbar \to 0$, $\mathcal{S}_{\Omega} \sim \frac{\pi^2}{3} \text{var} \mathbf{X}(\Omega)$ and $\|[\Pi_{\hbar,\mu}, \mathbb{1}_{\Omega}]\|_{\mathbf{J}^1} \sim \pi \text{var} \mathbf{X}(\Omega)$ so that the magnitude of the entropy is expected to be captured by our lower bound.

The more trivial estimate

$$\big\|[\Pi_{\hbar,\mu},\mathbb{1}_\Omega]\big\|_{J^1}=\mathcal{O}\big(\hbar^{-n}\big)$$

leads via the same interpolation to the upper-bound $S_{\Omega} = \mathcal{O}(\hbar^{1-n} \log^2(\hbar))$ appearing in the physics literature [28].

Note that the trace-norm of a commutator is generally more technical to estimate than its Hilbert-Schmidt norm (Theorem 1). This is the main reason why we presume that the estimate of Theorem 4 is not sharp. Theorem 4 should be compared, both in its result and its methods, to the main result of [25] that we now recall.

Proposition 1. Let $f \in C_c^{\infty}(\mathbb{R}^n)$ and suppose that $\int |t||\widehat{f}(t)|dt < \infty$. Under the Assumptions 1,

$$\|[\Pi_{\hbar,\mu}, f]\|_{\mathcal{J}^1} = \mathcal{O}\left(\hbar^{1-n} \int |t||\widehat{f}(t)|dt\right) \tag{1.10}$$

where the implied constant depends only on (V, μ) .

1.2 Organisational remarks

In Subsection 1.3, we concisely review the connections between our results and the random matrix literature with an emphasize on the Szegő asymptotics for determinants in the Fisher-Hartwig class. Then, in Subsection 1.4, we report on a conjecture of Widom [72] which relates number variance and entanglement entropy of (free) fermionic systems to spectral function of Schrödinger operators and we review the main progress on this conjecture.

As in our previous work [18, 19], our methods to study the commutator $[\Pi_{\hbar,\mu}, \mathbb{1}_{\Omega}]$ rely on the semiclassical machinery, in particular, on an approximation of the projection $\Pi_{\hbar,\mu}$ by a Fourier Integral operator. Under Assumptions 1, this approximation holds modulo an error of order \hbar^{1-n} in trace-norm, which is negligible in this context. In Section 2, we regroup the main notations and results from the literature on semiclassical Schrödinger operator H_{\hbar} that we need to prove our main results.

Section 3 gathers several additional Hilbert-Schmidt estimates for commutators between regular or mild spectral functions of H_{\hbar} and spatial functions at arbitrary scale.

Section 4 is dedicated to the proofs of Theorem 1 and its consequence, Theorem 2. The method relies on applying several stationary phase argument when the amplitude is discontinuous. The arguments are rather involved and therefore the proof is divided in several intermediate steps.

In Section 5, we rely on a multi-scale argument (we decompose both $\Pi_{\hbar,\mu}$ and $\mathbbm{1}_{\Omega}$ in dyadic pieces) and the estimates from Section 3 to prove Theorem 4. Then, Theorem 3 follows by a simple Schatten-norm interpolation which is explained in Subsection 5.2.

Finally, in the Appendix A, we gather different versions of the *stationary phase method* that we need when the amplitude varies on possibly arbitrary small scales.

1.3 Relation to random matrices and determinants in the Fisher-Hartwig class

In dimension 1, for a non-trivial interval $[a,b] \subset \mathcal{D}$, Theorem 1 states that, with N the particle number,

$$\operatorname{var} \mathbf{X}([a,b]) \sim \frac{1}{\pi^2} \log N.$$

In particular, this quantity is independent of the confining potential V and it matches with the Gaussian Unitary Ensemble (and sine process) number variance. The GUE variance computation goes back to Dyson and Mehta [47] and the study of the fluctuations to [13], see [58] for additional precisions and [46] for a study of large deviations. This universality is consistent with (1.6) and not surprising as it comes from the microscopic fluctuations (given by the sine process) of the particles around a, b. Moreover, there is an exact correspondence between the model (1.1) with $V: x \in \mathbb{R} \mapsto x^2$ and the Gaussian Unitary Ensemble (GUE) eigenvalues [62, 6].

These number variance asymptotics are explained by the log-correlated structure of the eigenvalues of random matrices, an observation which originates from [33, 29, 26] and has been revisited in [12] from the perspective of Gaussian multiplicative chaos. In short, one can view $b \in \mathbb{R} \mapsto \mathbf{X}((-\infty, b])$ as a Gaussian log-correlated field regularized on scale $\hbar \approx N^{-1}$ in the bulk. This has been a very active research topic lately and we cannot review all related results here. We simply mention that for the Gaussian β -ensemble¹ (normalised

¹The Gaussian β -ensemble corresponds to a fermionic model with Calogero-Sutherland type interaction for $\beta \neq 2$ and the $\mathcal{O}(1)$ -term has been conjectured in [58]. This term has been computed in the classical cases $\beta \in \{1, 2, 4\}$.

so that $\mathcal{D} = (-1,1)$, with $\mathbf{Z}(b) := \frac{\mathbf{X}((-\infty,b]) - \frac{2}{\pi} \int_{\lambda \leq b} (1-\lambda^2)_+^{1/2} d\lambda}{\sqrt{\log N}}$, one has (uniformly) for $b_1, \ldots, b_k \in \mathcal{D}$ such that Σ exists,

$$(\mathbf{Z}(b_1), \cdots, \mathbf{Z}(b_k)) \Rightarrow \mathcal{N}_{0,\Sigma} \quad \text{where} \quad \Sigma_{ij} = \frac{1}{\pi^2 \beta} \lim_{N \to \infty} \left(\frac{1}{\log N} \log_+ \min \left\{ N(1 - b_j^2)^{3/2}, \frac{1 - b_j^2}{|b_i - b_j|} \right\} \right). \quad (1.11)$$

(1.11) is due to [29] for GUE ($\beta = 2$) and also holds for general Wigner matrices [50, 48] and β -ensembles [3]. This CLT captures the behavior down to the microscopic scale as well as the edge effect, it should be compared to Theorem 2 for free fermions on \mathbb{R} . We refer to [42, 41] for more details on the log-correlated structure of the Gaussian β -ensemble, and [40, 51, 2] for the most recent results on the maximum fluctuations of the eigenvalue counting function of different random matrix model. A result analogous to Theorem 2 has also been obtained recently for the circular Riesz gas using methods from statistical mechanics [4].

For the eigenvalues of Hermitian (or unitary) matrix models, which are also determinantal processes, there are explicit formulae for the Laplace transform of any linear statistic in terms of large Hankel (or Toeplitz) determinants. The basic example, known as the circular unitary ensemble (CUE), corresponds to determinants of the type (1.3) involving spectral projector $\Pi_{\hbar} = \mathbb{1}(-\hbar^2 \Delta_{\mathbb{S}^1} \leq 1)$. In contrast to the Schrödinger model, Toeplitz determinants with a smooth symbol, as well as their continuous counterpart with $\mathbb{1}(-\hbar^2\Delta_{\mathbb{R}}\leq 1)$, can be analyzed by a miscellary of different techniques and the (Gaussian) behavior of such determinants are known as Szegő asymptotics. These asymptotics have constituted a central problem in mathematical physics because of a plethora of applications (Ising model, impenetrable bosons, Toda chain, random matrices, etc.), see [16] for a review and [18] for more on the relation to one-dimensional free fermions. A special class of singular symbols, called the Fisher-Hartwig class, plays a fundamental role in these applications, in particular to describe the fluctuations of the characteristic polynomial and eigenvalue counting functions of matrix models. These determinants also arise when computing the entanglement entropy of one-dimensional quantum spin chains [34, 36, 37]. Fisher-Hartwig [24] already conjectured that the asymptotics of these determinants involved extra $\log N$ terms. The first results on Fisher-Hartwig determinants go back to Widom's seminal work [69] (see also [70, 71]) and the full conjecture was resolved in [15] based on the connection with orthogonal polynomials with varying weights and the corresponding Riemann-Hilbert problem. The article [15] relies on the (non-linear) steepest descent method [17] for oscillatory Riemann-Hilbert problem to obtain precise asymptotics valid, for instance, for the Laplace transform of counting statistics. This requires to develop new local parametrices at the singular points. Our approach (to the free fermions case) is similar in spirit, since by using Fourier integral approximation for the Schrödinger propagator, we reduce the problem of computing the $\log N$ term in the variance to applying the stationary phase method to an oscillatory integral with a discontinuous amplitude. We cannot review the rich literature on Fisher-Hartwig determinants and we refer instead to the survey [16], [10, 22] for recent developments and further references, as well as [39, 68, 12, 35] for some probabilistic applications to random matrices.

In dimension $n \geq 2$, there are several results in the physics literature concerning disk counting statistics of free fermions confined by a rotation-invariant potential [57, 59] (by scaling, one can compare these predictions to the asymptotics of Theorem 1 in case $V(x) = |x|^q$ with q > 0 and $\mu = 1$). According to (1.8), writing V(x) = v(r), for a disk $\Omega = \{|x| \leq r\}$ within the bulk $(v(r) < \mu)$,

$$C_{\Omega} = \frac{\mathbf{c}_{n-1}|\partial B_n|}{2\pi^2} r^{n-1} (\mu - \mathbf{v}(r))^{\frac{n-1}{2}} = \frac{n\mathbf{c}_{n-1}\mathbf{c}_n}{2\pi^2} r^{n-1} (\mu - \mathbf{v}(r))^{\frac{n-1}{2}} = \frac{1}{\pi^2 \Gamma(n)} (2\pi r)^{n-1} (\mu - \mathbf{v}(r))^{\frac{n-1}{2}}$$

since $c_{n-1}c_n = 2\frac{(2\pi)^{n-1}}{\Gamma(n+1)}$ by Legendre duplication formula. Hence, we obtain as $\hbar \to 0$,

$$\operatorname{var} \mathbf{X}(rB_n) = \frac{\hbar^{1-n} (\log \hbar^{-1} + \mathcal{O}(1))}{\pi^2 \Gamma(n)} (r\sqrt{1 - \mathbf{v}(r)})^{n-1}$$

which is consistent with [57], formula (4) with $\mu \leftarrow 1/\hbar$. We also refer to formulae (S55) and (S56) in the supplementary material of [57] for an expression of the $\mathcal{O}(1)$ correction terms in case of a rotation-invariant potential.

1.4 Widom's conjecture & entanglement entropy

From the viewpoint of spectral theory and physical applications, it is of interest to describe spectral functions of the type $g(\Pi_{\hbar}|_{\Omega})$ where $g:[0,1] \to \mathbb{R}_{+}$ is continuous, Π_{\hbar} is a semiclassical (self-adjoint) projection on $L^{2}(\mathbb{R}^{n})$, $\Omega \in \mathbb{R}^{n}$ be open and let $\Pi_{\hbar}|_{\Omega} = \mathbb{1}_{\Omega}\Pi_{\hbar}\mathbb{1}_{\Omega}$. The case where g(0) = g(1) = 0 is of special interest. For instance, with $g: t \mapsto t(1-t)$,

$$g(\Pi_{\hbar}|_{\Omega}) = \mathbb{1}_{\Omega}(\Pi_{\hbar} - \Pi_{\hbar}\mathbb{1}_{\Omega}\Pi_{\hbar})\mathbb{1}_{\Omega}$$

so that

$$\operatorname{tr} g(\Pi_{\hbar}|_{\Omega}) = \operatorname{tr}((\Pi_{\hbar} - \Pi_{\hbar} \mathbb{1}_{\Omega} \Pi_{\hbar}) \mathbb{1}_{\Omega}) = -\frac{1}{2} \operatorname{tr}[\Pi_{\hbar}, \mathbb{1}_{\Omega}]^{2},$$

which corresponds to the case of Theorem 1; see more generally Lemma 5.4.

The model case where $\Pi_{\hbar} = \mathbb{1}(-\Delta < \hbar^{-2})$ in dimension 1 and Ω is an interval has been studied in [43] using the Mellin transform and the results were generalised to pseudodifferential operators with discontinuous symbols (replacing Π_{\hbar} by $\operatorname{Op}_{\hbar}(a)$ where the amplitude $a : \mathbb{R}^2 \to \mathbb{C}$ is not necessarily continuous) in [72]. Widom also conjectured in [72] an analogous result for pseudodifferential operators with discontinuous symbols in higher dimensions. A notable first step was the computation of the variance in [28] (that is, the case $g : \lambda \mapsto \lambda(1-\lambda)$), as well as an upper estimate on the entropy (using a cruder bound on the J¹-norm) of the form $S_{\Omega} \leq C\hbar^{1-n}\log(\hbar)^2$. The proof of the Widom conjecture when $g : \mathbb{R} \to \mathbb{R}$ is analytic, along with some history and further references, is gathered in [60].

Recent developments focus on the case where g is less regular, including the entropy [61], pseudodifferential operators with matrix-valued symbols [23, 1], and magnetic Laplacians with large magnetic fields [9, 45, 52]; in this latter case the analysis is slightly different due to a spectral gap in the ground state. Having in mind the rich applications to fermionic many-body physics and determinantal point processes, it is relevant to try and extend this program to more general spectral projectors. Inspired by Widom's conjecture [72], in the context of this article, we propose the following conjecture.

Conjecture 1. Under the Assumptions 1, let $\Pi_{\hbar,\mu}$ be as in (1.2) and let $\Pi_{\hbar}|_{\Omega} = \mathbb{1}_{\Omega}\Pi_{\hbar,\mu}\mathbb{1}_{\Omega}$. Let $g:[0,1] \to \mathbb{R}$ be continuous with g(0) = g(1) = 0 such that either $g \ge 0$ or $t \mapsto \frac{g(t)}{t(1-t)}$ is in L^1 . Then, as $\hbar \to 0$,

$$\frac{\operatorname{tr} g(\Pi_{\hbar}|_{\Omega})}{(2\pi\hbar)^{1-n}\log\hbar^{-1}} \to 2C_{\Omega} \int_{[0,1]} \frac{g(\lambda)}{\lambda(1-\lambda)} d\lambda$$

where C_{Ω} is given by (1.8) with f = 1.

Lemma 5.4 gives a relationship between Schatten-like norms of $[\Pi_{\hbar,\mu}, \mathbb{1}_{\Omega}]$ and traces of spectral functions of $\Pi_{\hbar}|_{\Omega}$. In particular, Conjecture 1 contains as a particular case the asymptotics of $\|[\Pi_{\hbar,\mu}, \mathbb{1}_{\Omega}\|_{J^1}$, of the von Neumann entropy of Theorem 3, and also the Rényi entropies. In these cases, as we already mentioned, this conjecture is also supported by numerical simulations [67] and theoretical physics computations [7, 6, 21]. Moreover, formula (1.8) agrees with the geometric constants appearing in previously studied forms of the conjecture [60, 27, 44].

In future work, we intend to return to proving Widom-type asymptotics for general spectral functions, and to bridge the gap with the existing results concerning pseudodifferential operators with discontinuous symbols.

1.5 Acknowledgements.

A.D. acknowledges the support of the CNRS PEPS 2021 grant. G.L. acknowledges the support of the starting grant 2022-04882 from the Swedish Research Council and of the starting grant from the Ragnar Söderbergs Foundation.

2 Pseudodifferential operators and Fourier Integral operators

2.1 Notation

Throughout the article, we will not always emphasize that the operators $H = H_{\hbar}, \Pi = \Pi_{\hbar,\mu}$, and that functions, depend on the semiclassical parameter \hbar . In particular, we will consider a class S^{δ} of functions (symbols) which depend on a small scale $\delta(\hbar)$.

Definition 2.1. Let $d \in \mathbb{N}$ and $\delta = \delta(\hbar) \in [\hbar, 1]$. Given $\Omega \subseteq \mathbb{R}^d$ open and bounded, we define the following class of functions:

$$S^{\delta}(\Omega) = \{ a \in \mathscr{C}_c^{\infty}(\Omega) : ||a||_{\mathscr{C}^k} \le C_k \delta^{-k} \text{ for every } k \in \mathbb{N}_0 \}.$$

We also use the notations $a \in S^{\delta}(\mathbb{R}^d)$, or $a \in S^{\delta}$, if $a \in S^{\delta}(\Omega)$ for some $\Omega \in \mathbb{R}^d$ bounded. For $\Theta = \Theta(\hbar) \in (0,1]$, we also write $a = \mathcal{O}_{S^{\delta}}(\Theta)$ if $a/\Theta \in S^{\delta}$.

In particular, $a \in S^1(\mathbb{R}^d)$ if it is $C_c^{\infty}(\mathbb{R}^d)$, $\operatorname{supp}(a) \subseteq \Omega$ with Ω bounded, and Ω , $||a||_{\mathscr{C}^k}$ are independent of \hbar for all $k \in \mathbb{N}$.

If a symbol depends on several variables, e.g. $(x,z) \in \mathbb{R}^{n \times m}$, with different regularity exponents, say $\epsilon, \delta \in [\hbar, 1]$, we will denote $a \in S_x^{\epsilon} \times S_z^{\delta}$ if for every indices $\alpha \in \mathbb{N}_0^n, \beta \in \mathbb{N}_0^m$ with $|\alpha| + |\beta| \le k$,

$$\sup_{x,z} \left| \partial_x^{\alpha} \partial_z^{\beta} a(x,\xi) \right| \le C_k \epsilon^{-|\alpha|} \delta^{-|\beta|}.$$

In the sequel, by *constants*, we mean positive numbers, e.g. C, c, independent of \hbar . Such *constants* are allowed to depend on the potential V and other auxiliary parameters. Sometimes, we denote c_{α} to emphasize that the constant c depends on a parameter α .

We declare right now the following objects and conventions, which we fix for the rest of the article.

- 1. Replacing V with $V \mu$, we assume that $\mu = 0$. The droplet is $\mathcal{D} = \{V < 0\}$.
- 2. $\Omega \in \mathcal{D}$ is a fixed relatively compact open set, with a smooth boundary. $w : \mathbb{R}^n \to \mathbb{R}$ is a smooth, tends to $+\infty$ at ∞ , such that $\Omega := \{w < 0\}$ and $\partial_x w \neq 0$ on $\partial\Omega = \{w = 0\}$. $\Omega' \in \{V < 0\}$ is an open neighbourhood of Ω inside the droplet.
- 3. $\underline{\tau}, \underline{\ell}, \underline{c}$ are small positive constants. The constant $\underline{\ell}$ is chosen much smaller than \underline{c} , and the constant $\underline{\tau}$ is chosen much smaller than $\underline{\ell}$, and all constants are small, in a way which depends on V and Ω . In particular, we assume that V is C^{∞} on $\{V < 2\underline{c}\}$, that $\partial_x V(x) \neq 0$ for all $x \in \{|V| \leq \underline{c}\}$, and that $\Omega' \subset \{V \leq -2\underline{c}\}$.
- 4. $\rho \in \mathcal{S}(\mathbb{R}, \mathbb{R}_+)$ is even with $\int_{\mathbb{R}} \rho(\lambda) d\lambda = 1$. Its Fourier transform $\hat{\rho}$ is supported inside $[-\underline{\tau}, \underline{\tau}]$.
- 5. Given $f \in L^{\infty}(\mathbb{R}^d)$, for $\delta \in (0,1]$, we denote $f_{\delta} : x \mapsto f(\delta^{-1}x)$. This notation admits one exception: we will denote $\rho_{\hbar} : u \mapsto \hbar^{-1}\rho(\hbar^{-1}u)$, that is, ρ is scaled as a density rather than a function.
- 6. Given $\vartheta \in \mathscr{C}_c^{\infty}(\mathbb{R})$ and given $\varkappa \in L^{\infty}(\mathbb{R})$ with compact support, we define $\widetilde{\varkappa} := \vartheta \cdot (\varkappa * \rho_{\hbar})$. Note that if $\varkappa \in S^{\eta}$ with $\eta \in [\hbar, 1]$ (see Definition 2.1), then $\widetilde{\varkappa} \in S^{\eta}$.

2.2 Pseudodifferential calculus

The goal of this section is to gather classical results on spectral functions of H. The main ingredient is that spectral functions involving the propagator, of the form $\vartheta(H)e^{-i\frac{tH}{h}}$ where $\vartheta \in \mathscr{C}_c^{\infty}(\mathbb{R})$ and $t \in \mathbb{R}$ is small, are given by Fourier Integral operators, up to negligible errors. Such strong approximations go back to [11, 30], but for consistency, we extract these results from our previous work [19, Proposition 2.11] as they fit the exact hypotheses of our main claims (see also [53, 20] for further reference). Recall that the potential V satisfies Assumptions 1 with $\mu = 0$ and $\underline{c} > 0$ is a small constant such that $V \in \mathscr{C}^{\infty}$ on $\{V < \underline{c}\}$.

Proposition 2.2. Let $\vartheta \in \mathscr{C}_c^{\infty}((-\infty,\underline{c}])$. There exists a smooth function $\varphi : \mathbb{R}^{2n+1} \to \mathbb{R}$ (independent of \hbar) and $s \in S^1(\mathbb{R}^{3n+1})$, such that, uniformly for $t \in [-\underline{\tau},\underline{\tau}]$,

$$\vartheta(H)e^{i\frac{tH}{\hbar}}:(x,y)\mapsto \frac{1}{(2\pi\hbar)^n}\int e^{i\frac{\varphi(t,x,\xi)-\xi\cdot y}{\hbar}}s(t,x,y,\xi)\mathrm{d}\xi+\mathcal{O}_{\mathrm{J}^1}(\hbar^\infty).$$

For every $t \in [-\underline{\tau},\underline{\tau}]$, the amplitude s is supported in a neighbourhood of

$$\{|x-y| \le \underline{\ell}, \ \vartheta(V(x) + |\xi|^2) \ne 0\}.$$

Moreover, its principal part is given on the diagonal by, at t = 0,

$$s(0, x, x, \xi)|_{\hbar=0} = \vartheta(|\xi|^2 + V(x)), \qquad (x, \xi) \in \mathbb{R}^{2n}$$

The phase φ is the solution of the following initial value problem for the Hamilton-Jacobi equation on a neighbourhood of the support of s:

$$\partial_t \varphi(t, x, \xi) = V(x) + |\partial_x \varphi(t, x, \xi)|^2$$
 $\varphi|_{t=0} : (x, \xi) \mapsto x \cdot \xi.$

Remark 2.3. Note that s silently depends on \hbar ; in fact it can be taken to be a smooth function of \hbar on $[0, \hbar_0]$. Usually, s is represented by its Taylor expansion at $\hbar = 0$, and the first term of this expansion (the value at 0) is called the *principal symbol*. We will use this notion throughout this article, without writing down explicitly the dependence of s (and other amplitudes) on \hbar .

Using the Fourier inversion formula, Proposition 2.2 provides an approximation of spectral functions of H which are smooth on scales larger than \hbar . More precisely we obtain the following corollary.

Corollary 2.4. Let φ be the phase from Proposition 2.2. Let $\vartheta \in \mathscr{C}_c^{\infty}((-\infty,\underline{c}])$, let $\varkappa \in L^{\infty}(\mathbb{R})$ with compact support and let $\widetilde{\varkappa} = \vartheta \cdot (\varkappa * \rho_{\hbar})$ as in the notations of Section 2.1. Then, there exists $a \in S^1(\mathbb{R}^{3n+1})$ so that

$$\widetilde{\varkappa}(H): (x,y) \mapsto \frac{1}{(2\pi\hbar)^{n+1}} \int e^{i\frac{\varphi(t,x,\xi)-y\cdot\xi-t\lambda}{\hbar}} b(t,x,y,\xi) \varkappa(\lambda) d\xi dt d\lambda + \mathcal{O}_{J^1}(\hbar^{\infty}).$$

The amplitude b is supported in $\{t \in [\underline{\tau},\underline{\tau}], |x-y| \leq \underline{\ell}\}$ and on a small neighbourhood of

$$\{(x,\xi): \vartheta(V(x) + |\xi|^2) \neq 0\}.$$

Up to a negligible error, controlled in the trace-norm, the kernel of the operator $\widetilde{\varkappa}(H)$ is given by an oscillatory integral. In fact, this kernel can be further simplified if $\varkappa \in S^{\eta}$ with $\eta \geq \hbar^{\frac{1}{2}}$; see Proposition 3.2. Let us also record two important observations regarding the support of the amplitude and the approximation of the phase.

Since $\underline{\tau}$ is much smaller than $\underline{\ell}$, which itself is much smaller than \underline{c} , we obtain that, if ϑ is supported in $[-\underline{c},\underline{c}]$, one has

$$x \in \Omega', y \in \Omega', (t, x, y, \xi) \in \text{supp}(a) \implies |\xi| \ge \underline{c}.$$
 (2.1)

Moreover, using the Hamilton-Jacobi equation, the phase has the following expansion as $t \to 0$,

$$\varphi(t, x, \xi) = x \cdot \xi + t(V(x) + |\xi|^2) + \mathcal{O}(t^2)$$
(2.2)

uniformly over the support of a, with a smooth error term, so that $\partial_t \varphi(t, x, \xi) = V(x) + |\xi|^2 + \mathcal{O}(t)$ as $t \to 0$.

2.3 Preliminary commutator estimates

We are interested in estimates for the Hilbert-Schmidt and trace-norm of the commutator $[\Pi, f|_{\Omega}]$ where $\Pi = \mathbb{1}_{\mathbb{R}_{-}}(H)$ is a spectral projector and $\Omega \in \mathcal{D}$ a subset of the droplet. A preliminary step to prove Theorem 1 consists in replacing Π by a Fourier integral operator $\widetilde{\Pi}$, up to a negligible error when computing $[\Pi, f|_{\Omega}]$. This operator is obtained by regularizing Π on scale \hbar using Proposition 2.2.

Proposition 2.5. Let φ be as in Proposition 2.2 and let $f \in C^{\infty}(\mathbb{R}^n)$. There exists $a \in S^1(\mathbb{R}^{3n+2})$ such that

$$[\Pi, f|_{\Omega}] = [\widetilde{\Pi}, f|_{\Omega}] + \mathcal{O}_{\mathcal{I}^1}(\hbar^{1-n})$$

where $\widetilde{\Pi}$ is a trace-class operator with a kernel

$$\widetilde{\Pi}: (x,y) \mapsto \frac{1}{(2\pi\hbar)^{n+1}} \int e^{i\frac{\varphi(t,x,\xi) - y \cdot \xi - t\lambda}{\hbar}} a(t,x,y,\xi,\lambda) \mathbb{1}\{\lambda \le 0\} d\xi dt d\lambda.$$
(2.3)

The amplitude a is supported in $\{t \in [-\underline{\tau},\underline{\tau}], |x-y| \leq \underline{\ell}, |V(x)+|\xi|^2| \leq \underline{c}, |\lambda| \leq \underline{\ell}\}$, satisfies (2.1) and its principal part satisfies at t=0,

$$a(0, x, x, \xi, \lambda)|_{\hbar=0} = \vartheta(|\xi|^2 + V(x))\chi(\lambda).$$

We expect the error $\mathcal{O}_{J^1}(\hbar^{1-n})$ to be sharp; it is negligible compared to the asymptotics of Theorem 1. Note also that the kernel (2.3) has a discontinuous amplitude, which makes its analysis non-trivial.

The proof of Proposition 2.5 occupies the rest of this section, the operator Π being constructed using Corollary 2.4 with $\varkappa = \chi \mathbb{1}_{\mathbb{R}_-}$ and two cutoffs:

- $\vartheta : \mathbb{R} \to [0,1]$ supported in $[-\underline{c},\underline{c}]$ and equal to 1 on $[-\underline{c}/2,\underline{c}/2]$.
- $\chi: \mathbb{R} \to [0,1]$, compactly support in $[-\underline{\ell},\underline{\ell}]$ and equal to 1 on a neighborhood of 0.

Then, we have

$$\widetilde{\varkappa}(H) = \vartheta(H)(\chi \mathbb{1}_{\mathbb{R}_{-}} * \rho_{\hbar})(H) = \widetilde{\Pi} + \mathcal{O}_{J^{1}}(\hbar^{\infty})$$
 (2.4)

where $\widetilde{\Pi}$ is the Fourier integral operator (2.3) and the amplitude $a:(t,x,y,\xi,\lambda)\mapsto\sqrt{2\pi}\widehat{\rho}(t)s(t,x,y,\xi)\chi(\lambda)$ with s as in Proposition 2.2. This yields both the support condition for a and its principal part.

It remains to relate $\widetilde{\varkappa}(H)$ to the true projection $\Pi = \mathbb{1}_{\mathbb{R}_-}(H)$. The first (simple) step is to replace Π by $\vartheta(H)\mathbb{1}_{\mathbb{R}_-}(H)$ where $\vartheta \in \mathscr{C}_c^{\infty}(\mathbb{R})$ is supported in an arbitrary small neighbourhood of $\mu = 0$.

Proposition 2.6. Let $\vartheta : \mathbb{R} \to \mathbb{R}$ be smooth and compactly supported in $(-\infty,\underline{\mathbf{c}}]$. Let $\Omega \subseteq \mathbb{R}^n$ be a set with a smooth boundary and $f \in \mathscr{C}^{\infty}(\mathbb{R})$, then as $\hbar \to 0$,

$$\|[\vartheta(H), f|_{\Omega}]\|_{\mathcal{J}^1} = \mathcal{O}(\hbar^{1-n}).$$

Since it requires some auxiliary estimates, we postpone the proof of Proposition 2.6 to the end of this Section.

For the second step, going from $\vartheta(H)\mathbb{1}_{\mathbb{R}_-}(H)$ to Π , we need the following estimates for the eigenvalue counting function on microscopic scale. The proof follows the reasoning from [30, Theorem 4]. We emphasize that the argument is similar in spirit, albeit technically much more straightforward, to the application of the stationary phase method we will use to prove Theorems 1 and 4.

Lemma 2.7. Uniformly for $\lambda \in [-\underline{c}, \underline{c}]$,

$$\operatorname{tr} \mathbb{1}\{|H - \lambda| \le \hbar\} = \mathcal{O}(\hbar^{1-n}).$$

Proof. First observe that it follows directly from Proposition 2.2 with t = 0 that for any smooth function $\vartheta : \mathbb{R} \to \mathbb{R}_+$ supported on $(-\infty, \underline{c}]$,

$$\|\vartheta(H)\|_{\mathsf{J}^1} = \int_{\mathbb{R}^n} \vartheta(H)(x,x) \mathrm{d}x = \frac{1}{(2\pi\hbar)^n} \int b(0,x,x,\xi) \mathrm{d}x \mathrm{d}\xi + \mathcal{O}(\hbar^{\infty}) = \mathcal{O}(\hbar^{-n}). \tag{2.5}$$

Here, we used that the operator $\vartheta(H) \geq 0$ is trace-class so that $\|\vartheta(H)\|_{J^1} = \operatorname{tr} \vartheta(H)$.

Let $\mathcal{H}(x,\xi) := V(x) + |\xi|^2$ for $(x,\xi) \in \mathbb{R}^{2n}$. Using the Notation of Section 2.1, with $\widetilde{\delta_{\lambda}} := \vartheta \cdot (\delta_{\lambda} * \rho_{\hbar})$ for $\rho \in \mathcal{S}(\mathbb{R}, \mathbb{R}_+)$ and ϑ a cutoff equal to 1 on $[-\underline{c}/2, \underline{c}/2]$, one has $\rho(\underline{-\lambda}) = \hbar \widetilde{\delta_{\lambda}} + \mathcal{O}(\hbar^{\infty}\vartheta)$ if $\lambda \in [-\underline{c}/2, \underline{c}/2]$. Then, by Proposition 2.2 and (2.5),

$$\rho(\frac{H-\lambda}{\hbar}): (x,y) \mapsto \frac{1/2\pi}{(2\pi\hbar)^n} \int e^{i\frac{\varphi(t,x,\xi)-y\cdot\xi-t\lambda}{\hbar}} b(t,x,y,\xi) d\xi dt + \mathcal{O}_{J^1}(\hbar^{\infty})$$

where $(t, x, \xi) \mapsto b(t, x, x, \xi)$ is supported on $t \in [-\underline{\tau}, \underline{\tau}]$ and a small neighbourhood of $\{\mathcal{H}(x, \xi) = \lambda\}$, and the error term is independent of $\lambda \in [-\underline{c}/2, \underline{c}/2]$. In addition, this holds for any $\rho \in \mathcal{S}(\mathbb{R}, \mathbb{R}_+)$, not necessarily a probability density. Thus,

$$\operatorname{tr} \rho(\frac{H-\lambda}{\hbar}) \leq \frac{1}{(2\pi\hbar)^n} \left| \int e^{i\frac{\varphi(t,x,\xi) - x \cdot \xi - t\lambda}{\hbar}} b(t,x,x,\xi) \mathrm{d}x \mathrm{d}\xi \mathrm{d}t \right| + (\hbar^{\infty}).$$

We now estimate the previous integral using the stationary phase method. Observe that according to (2.2), the phase is $t(\mathcal{H}(x,\xi) - \lambda + \mathcal{O}(t))$ and, by Assumptions 1, the sets $\{\mathcal{H}(x,\xi) = \lambda\}$ for $\lambda \in [-\underline{c},\underline{c}]$ are diffeomorphic to spheres. In particular, letting $\eta = \mathcal{H}(x,\xi)$, we can make a (non-degenrate) change of coordinates $(x,\xi) \in \mathbb{R}^{2n} \to (\eta,\omega) \in \mathbb{R} \times \mathbb{S}^{2n-1}$ such that

$$\int e^{i\frac{\varphi(t,x,\xi)-x\cdot\xi-t\lambda}{\hbar}}b(t,x,x,\xi)d\xi dt = \int e^{i\frac{t(\varpi(t,\eta,\omega)-\lambda)}{\hbar}}a(t,\eta,\omega)d\omega d\eta dt$$

where $\varpi(t,\eta,\omega)=\eta+\mathcal{O}(t)$ with a smooth error and the amplitude $a\in S^1(\mathbb{R}^{2n+1})$ is supported in $\{t\in [-\underline{\tau},\underline{\tau}]\}$. Now, we apply the stationary phase method in the variable (t,η) , keeping $\omega\in\mathbb{S}^{n-1}$ fixed. There is a unique critical point $(t,\eta)=(0,\lambda)$ and the Hessian is the identity at the critical point. By Proposition A.1, we obtain

$$\left| \int e^{i\frac{\varphi(t,x,\xi) - x \cdot \xi - t\lambda}{\hbar}} b(t,x,x,\xi) dx d\xi dt \right| = \mathcal{O}(\hbar)$$

where the error term is uniform for $\{\lambda \in [-\underline{c},\underline{c}]\}$. By positivity, we conclude that there is a constant C so that

$$\operatorname{tr} \mathbb{1}\{|H - \lambda| \le \hbar\} \le \operatorname{tr} \rho(\frac{H - \lambda}{\hbar}) \le C\hbar^{1 - n}.$$

We are now ready to finish the proof of Proposition 2.5.

End of the proof of Proposition 2.5. By Proposition 2.6, since the operator H is bounded from below

$$[\Pi,\mathbb{1}_{\Omega}] = [\vartheta(H)\Pi,\mathbb{1}_{\Omega}] + [(1-\vartheta(H))\Pi,\mathbb{1}_{\Omega}] = [\vartheta(H)\Pi,\mathbb{1}_{\Omega}] + \mathcal{O}_{\mathsf{J}^1}(\hbar^{1-n}).$$

Recall that that $\widetilde{\varkappa} := \vartheta \cdot (\chi \mathbb{1}_{\mathbb{R}_-} * \rho_{\hbar})$ where the mollifier ρ is a Schwartz function and ϑ, χ are appropriate cutoffs. By construction, for every $k \in \mathbb{N}$, there is a function $\chi_k : \mathbb{R} \to [-1, 1]$, smooth with compact support so that

$$\widetilde{\varkappa} = \vartheta \mathbb{1}_{\mathbb{R}_-} + \chi_k(\hbar^{-1}\cdot) + \mathcal{O}(\hbar^k \vartheta).$$

Consequently, by Lemma 2.7 (see also (2.5)),

$$\widetilde{\varkappa}(H) = \vartheta(H)\Pi + \mathcal{O}_{J^1}(\hbar^{1-n})$$

By (2.4), this concludes the proof.

It remains to prove Proposition 2.6. We start with the following estimate from [20], formula (9.2): Let $K \in L^1(\mathbb{R}^{2n})$ be an integral kernel, then the corresponding operator satisfies

$$||K||_{\mathcal{J}^1} \le \sum_{\alpha \in \mathbb{N}_0^{2n}: |\alpha| \le 2n+1} ||\partial_{x,y}^{\alpha} K||_{L^1(\mathbb{R}^{2n})}. \tag{2.6}$$

Lemma 2.8. Let $\vartheta : \mathbb{R} \to \mathbb{R}$ be smooth and supported in $(-\infty,\underline{c}]$ and let $f : \mathbb{R}^n \to \mathbb{R}$ with compact support and $||f||_{\mathscr{C}^1} \leq 1$. Then, as $\hbar \to 0$,

$$[\vartheta(H), f] = O_{\mathcal{J}^1}(\hbar^{1-n}).$$

Proof. By Proposition 2.2 with t = 0, there is $a \in S^1(\mathbb{R}^{3n})$ so that

$$\vartheta(H): (x,y) \mapsto \frac{1}{(2\pi\hbar)^n} \int e^{i\frac{\xi \cdot (x-y)}{\hbar}} a(x,y,\xi) d\xi + \mathcal{O}_{J^1}(\hbar^{\infty}). \tag{2.7}$$

Hence,

$$[\vartheta(H), f] = K + \mathcal{O}_{J^1}(\hbar^{\infty}), \qquad K : (x, y) \mapsto \frac{1}{(2\pi\hbar)^n} \int e^{i\frac{\xi \cdot (x - y)}{\hbar}} a(x, y, \xi) (f(x) - f(y)) d\xi.$$

Up to a unitary scaling (changing $x \leftarrow x\hbar$), which fixes the J¹-norm, the kernel K is equivalent to

$$\widetilde{K}: (x,y) \mapsto \frac{1}{(2\pi)^n} \int e^{i(x-y)\cdot\xi} a(\hbar x, \hbar y, \xi) (f(\hbar x) - f(\hbar y)) d\xi = \widehat{a}(\hbar x, \hbar y, x - y) (f(\hbar x) - f(\hbar y)).$$

where $\widehat{a}(x,y,\cdot)$ denotes the Fourier transform (appropriately normalized) of $\xi \mapsto a(x,y,\xi)$ for $(x,y) \in \mathbb{R}^{2n}$ – in particular, $\widehat{a} \in \mathcal{S}(\mathbb{R}^{3n})$ with norms independent of \hbar .

Hence, if $||f||_{\mathscr{C}^1} \leq 1$, we can bound for any $k \in \mathbb{N}_0$,

$$|\widetilde{K}(x,y)| \le \hbar \frac{\varphi_k(\hbar x)||x-y|}{1+|x-y|^k}$$

where $\varphi_k \in \mathcal{S}(\mathbb{R}^n)$. This estimate implies that $\|\widetilde{K}\|_{L^1(\mathbb{R}^{2n})} = \mathcal{O}(\hbar^{1-n})$.

In addition, observe that by a similar computation for any index $\alpha \in \mathbb{N}_0^{2n}$ and $k \in \mathbb{N}_0$,

$$\partial_{x,y}^{\alpha}\widetilde{K}(x,y) = \mathcal{O}_{\alpha}\left(\hbar \frac{\varphi_k(\hbar x)||x-y|}{1+|x-z|^k}\right)$$

Thus, using (2.6), this shows that $||K||_{J^1} = \mathcal{O}(\hbar^{1-n})$, which completes the proof.

In order to prove Proposition 2.6, we need a similar estimates for (smooth) function which are constant except on a \hbar -neighborhood of the boundary $\partial\Omega$. Let w be as in Notation 5.1 below; this includes functions of the type $f = \chi(\hbar^{-1}w)$ with $\chi : \mathbb{R} \to \mathbb{R}_+$ is smooth and compactly supported.

Lemma 2.9. Let $\vartheta : \mathbb{R} \to \mathbb{R}$ be smooth and supported in $(-\infty,\underline{\mathbf{c}}]$. Let $\mathbf{f} \in S^{\hbar}(\mathbb{R}^n)$ and suppose that \mathbf{f} is constant on $\{x \in \mathbb{R}^n : \mathbf{w}(x) > c\hbar\}$ and constant on $\{x \in \mathbb{R}^n : \mathbf{w}(x) < -c\hbar\}$. Then, as $\hbar \to 0$,

$$\|[\vartheta(H), f]\|_{J^1} = O(\hbar^{1-n}).$$

Proof. Let $\triangle := \{x \in \mathbb{R}^n : |\mathbf{w}(\hbar x)| \le c\hbar\}$ be a rescaled \hbar -neighborhood of $\partial\Omega$. In particular, since Ω is a smooth compact set, $|\triangle| = \mathcal{O}(\hbar^{1-n})$. Let $f(x) := f(\hbar x)$ so that all the norms $||f||_{\mathscr{C}^k}$ are all controlled independently of \hbar ; since $\mathbf{f} \in S^{\hbar}(\mathbb{R}^n)$. Like in the proof of Lemma 2.8, one has $||[\vartheta(H), \mathbf{f}]||_{J^1} = ||\widetilde{K}||_{J^1} + O(\hbar^{1-n})$ where the operator \widetilde{K} has kernel

$$\widetilde{K}:(x,y)\mapsto \widehat{a}(\hbar x,\hbar y,x-y)\{f(x)-f(y)\}$$

where $\hat{a} \in \mathcal{S}(\mathbb{R}^{3n})$ with norms independent of \hbar . In particular, by a simple volume estimate

$$\|\widetilde{K}\|_{L^1} = \int_{\triangle^c \times \triangle^c} |\widetilde{K}(x,y)| dy dx + \mathcal{O}(|\triangle|), \qquad |\triangle| = \mathcal{O}(\hbar^{1-n}).$$

Then, by assumptions, for $x, y \in \triangle^c = \mathbb{R}^n / \triangle$,

$$|f(x) - f(y)| \le \mathbb{1}\{w(x) \ge c\hbar, -w(y) \ge c\hbar\} + \mathbb{1}\{w(y) \ge c\hbar, -w(x) \ge c\hbar\}.$$

If $|\varkappa(z)| \leq \frac{C_k}{(1+|z|)^{2k}}$ for $z \in \mathbb{R}^n$, $k \in \mathbb{N}$, using that $w(x) \asymp \operatorname{dist}(x, \partial\Omega)$, we also have

$$\int \mathbb{1}\{\mathbf{w}(\hbar x) \ge c\hbar, -\mathbf{w}(\hbar y) \ge c\hbar\} \varkappa(x - y) dx dy \le C|\triangle| = \mathcal{O}(\hbar^{1-n}).$$

Since $|\widehat{a}(x,y,z)| \leq \varkappa(z)$ for $(x,y,z) \in \mathbb{R}^{3n}$ with \varkappa as above, this shows that

$$\int_{\Lambda^c \times \Lambda^c} |\widetilde{K}(x,y)| \mathrm{d}y \mathrm{d}x = \mathcal{O}(\hbar^{1-n}).$$

This establishes that $\|\widetilde{K}\|_{L^1} = \mathcal{O}(\hbar^{1-n})$. In addition, since all derivatives of \widetilde{K} are controlled independently of \hbar , we can similarly bound for any $\alpha \in \mathbb{N}_0^n$, $\|\partial_{x,y}^{\alpha}\widetilde{K}\|_{L^1(\mathbb{R}^{2n})} = \mathcal{O}_{\alpha}(\hbar^{1-n})$ using the same argument. By (2.6), we conclude that $\|\widetilde{K}\|_{J^1} = \mathcal{O}(\hbar^{1-n})$.

Remark 2.10. The same argument shows that if f is constant on $\{x \in \mathbb{R}^n : |\mathbf{w}(x)| > c\hbar\}$, without any smoothness condition (such as $\mathbf{f} \in S^{\hbar}$),

$$\|[\vartheta(H), \mathbf{f}]\|_{\mathbf{J}^2}^2 = \|\widetilde{K}\|_{L^2(\mathbb{R}^{2n})}^2 = O(\hbar^{1-n}).$$

Indeed, there is no need to differentiate the kernel K in order to obtain this estimate. In particular, this shows that

$$\|[\vartheta(H), \mathbb{1}_{\Omega}]\|_{\mathcal{J}^2}^2 = O(\hbar^{1-n}).$$

Equipped with these estimates, we are ready to proceed to prove Proposition 2.6.

Proof of Proposition 2.6. Recall that $\Omega \in \mathbb{R}^n$ be a smooth compact set and let $\chi \in C_c^{\infty}(\mathbb{R})$. Let $f = \chi(\hbar^{-1}w)$ where $\chi : \mathbb{R} \to [0,1]$ smooth, compactly supported, so that f satisfies the assumptions Lemma 2.9. The first step is to expand

$$\begin{split} [\vartheta(H),\mathbb{1}_{\Omega}] &= \vartheta(H)\mathbb{1}_{\Omega}(1-\vartheta(H)) - (1-\vartheta(H))\mathbb{1}_{\Omega}\vartheta(H) \\ &= f\vartheta(H)\mathbb{1}_{\Omega}(1-\vartheta(H)) - (1-\vartheta(H))\mathbb{1}_{\Omega}\vartheta(H)f + (1-f)\vartheta(H)\mathbb{1}_{\Omega}(1-\vartheta(H)) - (1-\vartheta(H))\mathbb{1}_{\Omega}\vartheta(H)(1-f). \end{split}$$

Let $h = (1 - f)\mathbb{1}_{\Omega}$. The function h also satisfies the assumptions of Lemma 2.9 so that

$$\|[\vartheta(H),f]\|_{J^1},\|[\vartheta(H),h]\|_{J^1}=\mathcal{O}(\hbar^{1-n}).$$

Hence, we have

$$\begin{split} (1-f)\vartheta(H)\mathbb{1}_{\Omega}(1-\vartheta(H)) - (1-\vartheta(H))\mathbb{1}_{\Omega}\vartheta(H)(1-f) \\ &= \vartheta(H)h(1-\vartheta(H)) - (1-\vartheta(H))h\vartheta(H) + [\vartheta(H),f]\mathbb{1}_{\Omega}(1-\vartheta(H)) + (1-\vartheta(H))\mathbb{1}_{\Omega}[\vartheta(H),f] \\ &= [\vartheta(H),h] + \mathcal{O}_{J^1}\big([\vartheta(H),f]\big) \\ &= \mathcal{O}_{J^1}(\hbar^{1-n}). \end{split}$$

Moreover, we can decompose $\mathbb{1}_{\Omega} = \mathbf{k} + \theta$ where k satisfies the assumptions of Lemma 2.9 with $\mathbf{k} = 0$ on supp(f) and θ is discontinuous, supported in a \hbar -neighborhood of $\partial\Omega$. In particular, $\|\mathbf{f}\|_{L^2}^2$, $\|\theta\|_{L^2}^2 = \mathcal{O}(\hbar)$ and $\|[\vartheta(H),\mathbf{k}]\|_{\mathrm{J}^1} = \mathcal{O}(\hbar^{1-n})$ so that

$$f\vartheta(H)\mathbb{1}_{\Omega}(1-\vartheta(H)) = f\vartheta(H)k(1-\vartheta(H)) + f\vartheta(H)\theta(1-\vartheta(H))$$
$$= f[\vartheta(H),k](1-\vartheta(H)) + f\vartheta(H)[\vartheta(H),\theta] + f\psi(H)^2\theta$$

where $\psi = \sqrt{\vartheta(1-\vartheta)} - \psi \in \mathscr{C}_c^{\infty}(\mathbb{R})$. By Proposition 3.1 and Remark 2.10,

$$\|f\vartheta(H)[\vartheta(H),\theta]\|_{J^1} \le \|f\vartheta(H)\|_{J^2}\|[\vartheta(H),\theta]\|_{J^2} = \mathcal{O}(\hbar^{1/2-n}\|f\|_{L^2}) = \mathcal{O}(\hbar^{1-n})$$

and

$$\|f\psi(H)^{2}\theta|_{J^{1}} \leq \|f\psi(H)\|_{J^{2}} \|\theta\psi(H)\|_{J^{2}} = \mathcal{O}(\hbar^{-n}\|f\|_{L^{2}}\|\theta\|_{L^{2}}) = \mathcal{O}(\hbar^{1-n}).$$

This shows that

$$\|f\vartheta(H)\mathbb{1}_{\Omega}(1-\vartheta(H))\|_{J^1}\leq \|[\vartheta(H),k]\|_{J^1}+\mathcal{O}(\hbar^{1-n})=\mathcal{O}(\hbar^{1-n}).$$

Similarly,
$$\|(1-\vartheta(H))\mathbb{1}_{\Omega}\vartheta(H)f\|_{J^1} = \mathcal{O}(\hbar^{1-n})$$
, so we conclude that $\|[\vartheta(H),\Omega]\|_{J^1} = \mathcal{O}(\hbar^{1-n})$.

3 Hilbert-Schmidt estimates for mild spectral functions

This section is devoted to preliminary estimates that we will use in the proof of Theorem 4: these estimates are Hilbert-Schmidt norms of products or commutators between spectral functions of H_{\hbar} and multiplication operators, with relatively small supports: the spectral functions will be supported on small neighbourhoods of 0, and the multiplication operators will be supported on small neighbourhoods of $\partial\Omega$.

Even though they are relatively independent, these estimates also serve as a warm-up for Section 4 where we will compute the limit of the Hilbert-Schmidt norm of the "full" commutator $[\Pi_{\hbar}, \mathbb{1}_{\Omega}]$ for Theorem 2. The same kind of stationary phase arguments are used in both cases and they are simpler to apply in the context of this section.

Recall the notations from Section 2.1 and assume that $\vartheta \in \mathscr{C}_c^{\infty}(\mathbb{R})$ is supported in $[-\underline{c},\underline{c}]$.

Proposition 3.1. Let $f, g : L^{\infty}(\mathbb{R}, \mathbb{R})$, be two functions supported in $[-\underline{c}, \underline{c}]$. For every $\hbar \in (0, 1]$, for every $\eta, \varepsilon \in [\hbar, 1]$,

$$\|\widetilde{f}_{\eta}(H)g_{\varepsilon}(\mathbf{w})\|_{\mathbf{J}^{2}}^{2} + \|f_{\eta}(H)g_{\varepsilon}(\mathbf{w})\|_{\mathbf{J}^{2}}^{2} \le C\hbar^{-n}\eta\varepsilon.$$
(3.1)

Let $f, g, k \in C_c^{\infty}(\mathbb{R})$ be supported in $[-\underline{c}, \underline{c}]$. For every $\hbar \in (0, 1]$, for every $\eta, \varepsilon \in [\hbar, 1]$ with $\eta \varepsilon \geq \hbar$,

$$\|[\widetilde{f}_{\eta}(H), g_{\varepsilon}(\mathbf{w})]\|_{\mathbf{J}^{2}}^{2} \le C \frac{\hbar^{n-2}}{\eta \varepsilon}$$
 (3.2)

$$\|[[\widetilde{f}_{\eta}(H), g_{\varepsilon}(\mathbf{w})], k_{\varepsilon}(\mathbf{w})]\|_{\mathbf{J}^{2}}^{2} \le C \frac{\hbar^{n-4}}{\eta^{3} \varepsilon^{3}}.$$
(3.3)

Observe that \widetilde{f}_{η} is a smooth function in the class $S^{\eta}(\mathbb{R})$, according to Definition 2.1, then using Corollary 2.4, one can replace (up to a negligible error) $\widetilde{f}_{\eta}(H)$ by a Fourier integral operator whose amplitude is in $S^1_{t,x,y,\xi} \times S^{\eta}_{\lambda}$. This allows us to apply the stationary phase method (Appendix A) in order to prove the above Hilbert-Schmidt estimates. The proof of Proposition 3.1 occupies the rest of this section and it is organized as follows.

- In Section 3.1, we proceed with the estimate for $\|\widetilde{f}_{\eta}(H)g_{\varepsilon}(\mathbf{w})\|_{\mathbf{J}^2}^2$. This quantity is given (up to a negligible error) by the oscillatory integral (3.4). After an appropriate change of coordinates, we can simplify (3.4) by a stationary phase argument. This reduces the problem to the computation of another oscillatory integral (3.10), with a mild amplitude, but with the classical phase $(x,\xi) \mapsto x \cdot \xi$. This method is important and we will use variants of it throughout the rest of this article. Then, we can deduce a similar estimate for $\|f_{\eta}(H)g_{\varepsilon}(\mathbf{w})\|_{\mathbf{J}^2}^2$ by monotonicity.
- In Section 3.2, we adapt the method from Section 3.1 to estimate the Hilbert-Schmidt norms of the commutators (3.2) and (3.3), if the spectral scale η is smaller than $\hbar^{\frac{1}{2}}$. Some difficulty comes from controlling the *support of the amplitude* after a first application of stationary phase. Note that the proofs of (3.2) and (3.3) are analogous.
- In Section 3.3, we prove (3.2) and (3.3) in the complementary regime where the spectral scale η is larger than $\hbar^{\frac{1}{2}}$. The proof relies on the fact that in this regime, $\tilde{f}_{\eta}(H)$ is (up to a negligible error) a pseudo-differential operator (Proposition 3.2). Then, using again the stationary phase method in a more direct way, we complete the proofs of (3.2)–(3.3).

3.1 Hilbert-Schmidt norm of products; proof of (3.1)

This subsection focuses on the proof of (3.1). Starting from Corollary 2.4, we have

$$\|\widetilde{f}_{\eta}(H_{\hbar})g_{\varepsilon}(\mathbf{w})\|_{\mathbf{J}^{2}}^{2} = \frac{1}{(2\pi\hbar)^{2n+2}} \int e^{i\frac{\Psi_{0}(x,y,\xi_{1},\xi_{2},t_{1},t_{2},\lambda_{1},\lambda_{2})}{\hbar}} B_{0}(x,y,\xi_{1},\xi_{2},t_{1},t_{2}) \times f_{n}(\lambda_{1})f_{n}(\lambda_{2})g_{\varepsilon}(\mathbf{w}(x))^{2} dt_{1}dt_{2}d\lambda_{1}d\lambda_{2}d\xi_{1}d\xi_{2}dxdy + \mathcal{O}(\hbar^{\infty})$$
(3.4)

where the error is independent of (η, ε) , the amplitude B_0 belongs to $S^1(\mathbb{R}^{3n+2})$, and we have introduced the phase

$$\Psi_0: (x, y, \xi_1, \xi_2, t_1, t_2, \lambda_1, \lambda_2) \mapsto \varphi(t_1, x, \xi_1) - \varphi(t_2, x, \xi_2) - y \cdot (\xi_1 - \xi_2) - t_1\lambda_1 + t_2\lambda_2. \tag{3.5}$$

The amplitude

$$B_0: (x, y, \xi_1, \xi_2, t_1, t_2) \mapsto b(t_1, x, y, \xi_1) \overline{b(t_2, x, y, \xi_2)} \chi(x, y)$$
(3.6)

where χ is a smooth cutoff supported in Ω'^2 . We can introduce this cutoff since $g_{\varepsilon}(\mathbf{w})$ is supported in $\{|\mathbf{w}| < \underline{\mathbf{c}}\}$ and b is supported in $|x-y| \le \underline{\ell}$ with $\underline{\ell} \ll \underline{\mathbf{c}}$. In particular, according to (2.1), the amplitude B_0 is supported in

$$\{t_1, t_2 \in [\underline{\tau}, \underline{\tau}], x, y \in \Omega', |x - y| \le \underline{\ell}, |\xi_1|, |\xi_2| \ge \underline{c}\}.$$

This allows to make the following change of coordinates in the integral (3.4):

$$\begin{cases} t = \frac{t_1 + t_2}{2}, & \lambda = \frac{\lambda_1 + \lambda_2}{2}, & \xi = \frac{\xi_1 + \xi_2}{2} = r\omega & \text{with } r > 0, \ \omega \in \mathbb{S}^{n-1} \\ s = t_1 - t_2, & \sigma = \lambda_2 - \lambda_1, & \zeta = \xi_1 - \xi_2 \end{cases}$$
(3.7)

We write

$$\Psi_0(x, y, \xi_1, \xi_2, t_1, t_2, \lambda_1, \lambda_2) = \Psi_1(x, y, r, t, \lambda, \omega, \zeta, s) + \sigma t$$

so that Ψ_1 is independent of σ and

$$\|\widetilde{f}_{\eta}(H_{\hbar})g_{\varepsilon}(w)\|_{J^{2}}^{2} = \frac{1}{(2\pi\hbar)^{2n+2}} \int e^{i\frac{\Psi_{1}(x,y,r,t,\lambda,\omega,\zeta,s)+\sigma t}{\hbar}} B_{1}(x,y,\zeta,\omega,r,s,t)$$

$$\times f_{\eta}(\lambda + \frac{\sigma}{2})f_{\eta}(\lambda - \frac{\sigma}{2})g_{\varepsilon}(w(x))^{2} \mathrm{d}y \mathrm{d}\zeta \mathrm{d}r \mathrm{d}s \mathrm{d}\omega \mathrm{d}t \mathrm{d}\sigma \mathrm{d}\lambda \mathrm{d}x + \mathcal{O}(\hbar^{\infty}).$$
(3.8)

At this stage, we apply the stationary phase method to the previous integral in the variable (y, ζ, r, s) keeping $(t, \omega, \sigma, \lambda, x)$ fixed. The equations for the critical point(s) are (using the notation (3.7))

$$\begin{cases} \partial_y \Psi_1 = 0 \\ \partial_\zeta \Psi_1 = 0 \\ \partial_r \Psi_1 = 0 \\ \partial_s \Psi_1 = 0 \end{cases} \iff \begin{cases} \zeta = 0 \\ y = \frac{1}{2} (\partial_\xi \varphi(t_1, x, \xi_1) + \partial_\xi \varphi(t_2, x, \xi_2)) \\ 0 = \frac{1}{2} \omega \cdot (\partial_\xi \varphi(t_1, x, \xi_1) - \partial_\xi \varphi(t_2, x, \xi_2)) \\ \lambda = \frac{1}{2} (\partial_t \varphi(t_1, x, \xi_1) + \partial_t \varphi(t_2, x, \xi_2)). \end{cases}$$

The first equation is equivalent to $\xi_1 = \xi_2 = r\omega$ with $r \geq c_2$. Then, using that $\partial_{\xi}\varphi(t_1, x, \xi) = x + t\xi + \mathcal{O}(t^2)$ and $\partial_t\varphi(t, x, \xi) = V(x) + |\xi|^2 + \mathcal{O}(t)$, these equations reduce to (uniformly in $\omega, \sigma, \lambda, x$)

$$\begin{cases} \zeta = 0 \\ y = x + \mathcal{O}(rt) \\ 0 = sr + \mathcal{O}(st) \\ \lambda = \partial_t \varphi(t, x, r\omega) \end{cases} \iff \begin{cases} \zeta = 0 \\ y = x + \mathcal{O}(rt) \\ s = 0 \\ r = \sqrt{\lambda - V(x) + \mathcal{O}(t)}. \end{cases}$$

Since we restrict our attention to $|t|, |s| < \underline{\tau}$, there is a unique critical point $(y, \zeta, r, s) = (y_{\star}, 0, r_{\star}, 0)$. At the critical point, $\Psi_1 = 0$ and

$$\operatorname{Hess}\Psi_{1}(x,t,\lambda,\omega) = \begin{pmatrix} 0 & \mathrm{I} & 0 & 0 \\ \mathrm{I} & * & * & 0 \\ 0 & * & 0 & 2r_{\star} + \mathcal{O}(t) \\ 0 & 0 & 2r_{\star} + \mathcal{O}(t) & 0 \end{pmatrix}; \qquad \begin{cases} r_{\star}(x,t,\lambda,\omega) = \sqrt{\lambda - V(x) + \mathcal{O}(t)} \\ y_{\star}(x,t,\lambda,\omega) = x + \mathcal{O}(t) \end{cases} . (3.9)$$

In particular det $\text{Hess}\Psi_1 = 4(\lambda - V(x)) + \mathcal{O}(t)$ is non-degenerate. As far as the phases are concerned, we are in position to apply stationary phase, and y_{\star}, r_{\star} are smooth functions of $\omega, \sigma, \lambda, x$.

Since B_1 belongs to S^1 , by Proposition A.1, there exists a symbol $B_2 \in S^1(\mathbb{R}^{2n+2})$ such that

$$\frac{1}{(2\pi\hbar)^{n+1}} \int e^{i\frac{\Psi_1(x,y,r,t,\lambda,\omega,\zeta,s)}{\hbar}} B_1(x,y,\zeta,\omega,r,s,t) dy d\zeta dr ds = B_2(x,\omega,\lambda,t).$$

Let $B_3:(x,\lambda,t)\mapsto \int_{\mathbb{S}^{n-1}}B_2(x,\omega,\lambda,t)\mathrm{d}\omega$, so that $B_3\in S^1(\mathbb{R}^{n+2})$. We have simplified (3.4) into

$$\|\widetilde{f}_{\eta}(H_{\hbar})g_{\varepsilon}(\mathbf{w})\|_{\mathbf{J}^{2}}^{2} = \frac{1}{(2\pi\hbar)^{n+1}} \int e^{i\frac{\sigma t}{\hbar}} B_{3}(x,\lambda,t) f_{\eta}(\lambda + \frac{\sigma}{2}) f_{\eta}(\lambda - \frac{\sigma}{2}) g_{\varepsilon}(\mathbf{w}(x))^{2} dt d\sigma d\lambda dx + \mathcal{O}(\hbar^{\infty}).$$
 (3.10)

Now, we compute the integral (3.10) by applying the stationary phase method in the variables (t, σ) keeping (x, λ) as parameters. Note that the amplitude is in $S_t^1 \times S_{\sigma}^{\eta}$ with $\eta \geq \hbar$ and it satisfies

$$\int B_3(x,\lambda,t) f_{\eta}(\lambda + \frac{\sigma}{2}) f_{\eta}(\lambda - \frac{\sigma}{2}) g_{\varepsilon}(\mathbf{w}(x))^2 d\lambda dx = \mathcal{O}_{S_t^1 \times S_{\sigma}^{\eta}}(\eta \varepsilon)$$

since $||g_{\varepsilon}(\mathbf{w})||_{L^{2}(\mathbb{R}^{n})}^{2} = \mathcal{O}(\varepsilon)$ and $||f_{\eta}||_{L^{2}(\mathbb{R})}^{2} = \mathcal{O}(\eta)$. Hence, by Corollary A.7, we conclude that

$$\|\widetilde{f}_{\eta}(H_{\hbar})g_{\varepsilon}(\mathbf{w})\|_{\mathbf{J}^{2}}^{2} = \mathcal{O}(\hbar^{-n}\eta\varepsilon).$$

This proves the first estimate in (3.1). To deduce the second estimate, we can assume that $f, g \geq 0$ and we choose the function $\vartheta \in C_c^{\infty}(\mathbb{R} \to \mathbb{R}_+)$ so that $\tilde{f}_{\delta} \geq f_{\delta}$. Then, as operators, for any $\varkappa : \mathbb{R}^n \to \mathbb{R}_+$ bounded with compact support,

$$\varkappa \widetilde{f}_{\eta}(H_{\hbar})^2 \varkappa \geq \varkappa f_{\eta}(H_{\hbar})^2 \varkappa.$$

With $\varkappa = \sqrt{g_{\varepsilon}(\mathbf{w})}$, taking traces on both sides, we conclude that $\|\widetilde{f}_{\eta}(H_{\hbar})g_{\varepsilon}(\mathbf{w})\|_{\mathbf{J}^{2}}^{2} \geq \|f_{\eta}(H_{\hbar})g_{\varepsilon}(\mathbf{w})\|_{\mathbf{J}^{2}}^{2}$.

3.2 Hilbert-Schmidt norm of commutators; proof of (3.2)–(3.3) at small spectral scales

This section is devoted to the proof of (3.2) and (3.3) under the assumption that the spectral scale η is smaller than $\hbar^{\frac{1}{2}}$. The arguments follow the strategy from the proof of (3.1) in the previous section, but they rely on the stationary phase with mild symbols (Proposition A.4); the hypotheses will be satisfied because $\varepsilon \eta \geq \hbar$ and in particular $\varepsilon \geq \hbar^{\frac{1}{2}}$, since we have assumed that $\eta \leq \hbar^{\frac{1}{2}}$.

We begin with (3.2). Introducing a dyadic sequence of open sets

$$\mho_k := \{ |\mathbf{w}| < \varepsilon_k \} \text{ with } \varepsilon_k = 2^k \varepsilon \text{ for } k \ge 0,$$

since without loss of generality supp $(g) \subset [-1,1]$, one has supp $(g_{\varepsilon}(\mathbf{w})) \subset \mho_0$.

As in (3.4)–(3.8), our starting point is the change of variables (3.7) leading to the expression

$$\|[\widetilde{f}_{\eta}(H_{\hbar}), g_{\varepsilon}(\mathbf{w})]\|_{\mathbf{J}^{2}}^{2} = \frac{1}{(2\pi\hbar)^{2n+2}} \int e^{i\frac{\Psi_{1}(x, y, r, t, \lambda, \omega, \zeta, s) + \sigma t}{\hbar}} B_{1,\varepsilon}(x, y, \zeta, \omega, r, s, t) \times f_{n}(\lambda + \frac{\sigma}{2}) f_{n}(\lambda - \frac{\sigma}{2}) \mathrm{d}y \mathrm{d}\zeta \mathrm{d}r \mathrm{d}s \mathrm{d}\omega \mathrm{d}t \mathrm{d}\sigma \mathrm{d}\lambda \mathrm{d}x + \mathcal{O}(\hbar^{\infty}).$$
(3.11)

Here

$$B_{1,\varepsilon}(x,y,\zeta,\omega,r,s,t) = B_0(x,y,\zeta,\omega,r,s,t) (g_{\varepsilon}(\mathbf{w}(x)) - g_{\varepsilon}(\mathbf{w}(y)))^2$$

The change of coordinates (3.7) is still well-defined, and as in the previous subsection, with respect to the variables (y, ζ, r, s) , there is a unique critical point $(y_{\star}, 0, r_{\star}, 0)$, satisfying (3.9). The amplitude $B_{1,\varepsilon}$ belongs to $S^{\varepsilon}(\mathbb{R}^{4n+2})$ with $\varepsilon \geq \hbar^{\frac{1}{2}}$, so we can apply the stationary phase method to the integral (3.11), with respect to the variables (y, ζ, r, s) , the other variables being treated as parameters. All in all, by Proposition A.4, there exists a symbol $B_{3,\varepsilon} \in S^{\varepsilon}(\mathbb{R}^{n+2})$ such that (integrating also over $\omega \in \mathbb{S}^{n-1}$)

$$\frac{1}{(2\pi\hbar)^{n+1}} \int e^{i\frac{\Psi_1(x,y,r,t,\lambda,\omega,\zeta,s)}{\hbar}} B_{1,\varepsilon}(x,y,\zeta,\omega,r,s,t) dy d\zeta dr ds d\omega = B_{3,\varepsilon}(x,\lambda,t).$$

Moreover, this symbol has the following properties: $B_{3,\varepsilon}$ is supported in $\{x \in \Omega\}$,

$$B_{3,\varepsilon}(x,\lambda,0) = \mathcal{O}(\hbar\varepsilon^{-2}) \tag{3.12}$$

and, for any fixed $\ell \in \mathbb{N}$, there is a small constant c > 0 so that for every $k \geq 2$,

$$B_{3,\varepsilon}(x,\lambda,t) = \mathcal{O}_{S_t^{\varepsilon}}\left(\left(\frac{\hbar}{\varepsilon\varepsilon_k}\right)^{\ell}\right), \quad \text{for } x \in (\mathcal{O}_k \setminus \mathcal{O}_{k-1}), |t| \le c\varepsilon_k, |\lambda| \le c.$$
 (3.13)

Here, (3.12) follows directly from (A.3) with $\ell = 1$, using that L^0 is a multiplication operator and $y_{\star}|_{t=0} = x$. Then, (3.13) follows from (A.4). Indeed, since $y_{\star}(x,t,\lambda,\omega) = x + \mathcal{O}(t)$ and we assume that $\partial_x w \neq 0$ on $\{w = 0\}$, we have for every $k \geq 2$,

$$\operatorname{dist}(y_{\star}(x,t,\lambda,\omega),\mho_0) \geq \varepsilon_k, \qquad \text{for } x \in (\mho_k \setminus \mho_{k-1}), \ |t| \leq c\varepsilon_k, \ |\lambda| \leq c, \ \omega \in \mathbb{S}^{n-1}.$$

By (3.11), this computation implies that

$$\|[\widetilde{f}_{\eta}(H_{\hbar}), g_{\varepsilon}(\mathbf{w})]\|_{\mathbf{J}^{2}}^{2} = \frac{1}{(2\pi\hbar)^{n+1}} \int e^{i\frac{\sigma t}{\hbar}} B_{3,\varepsilon}(x,\lambda,t) f_{\eta}(\lambda + \frac{\sigma}{2}) f_{\eta}(\lambda - \frac{\sigma}{2}) dt d\sigma d\lambda dx + \mathcal{O}(\hbar^{\infty}).$$
(3.14)

Since $|\mho_1| \leq C\varepsilon$ and $||f_\eta||^2_{L^2(\mathbb{R})} = \mathcal{O}(\eta)$, we have

$$\int_{\{x \in \mathcal{O}_1\}} B_{3,\varepsilon}(x,\lambda,t) f_{\eta}(\lambda + \frac{\sigma}{2}) f_{\eta}(\lambda - \frac{\sigma}{2}) d\lambda dx = \mathcal{O}_{S_t^{\varepsilon} \times S_{\sigma}^{\eta}}(\eta \varepsilon),$$

thus by Corollary A.7 (with $\ell = 2$),

$$\frac{1}{2\pi\hbar} \int_{\{x \in \mathcal{V}_1\}} e^{i\frac{\sigma t}{\hbar}} B_{3,\varepsilon}(x,\lambda,t) f_{\eta}(\lambda + \frac{\sigma}{2}) f_{\eta}(\lambda - \frac{\sigma}{2}) dt d\sigma d\lambda dx$$

$$= \int_{\{x \in \mathcal{V}_1\}} B_{3,\varepsilon}(x,\lambda,0) f_{\eta}(\lambda)^2 dx d\lambda + \mathcal{O}(\frac{\hbar^2}{\eta \varepsilon}). \quad (3.15)$$

Here, we used that the amplitude of the previous integral is in $S_t^{\varepsilon} \times S_{\sigma}^{\eta}$ with $\varepsilon \eta \geq \hbar$, that $B_{3,\varepsilon}$ is independent of σ , and that $\sigma \mapsto f_{\eta}(\lambda + \frac{\sigma}{2})f_{\eta}(\lambda - \frac{\sigma}{2})$ is a odd function. Then, by (3.12), the previous integral is $\mathcal{O}(\hbar \varepsilon^{-1} \eta)$. Since by assumption $\eta^2 \leq \hbar$, we conclude that

$$\int \mathbb{1}\{x \in \mathcal{O}_1\} e^{i\frac{\sigma t}{\hbar}} B_{3,\varepsilon}(x,\lambda,t) f_{\eta}(\lambda + \frac{\sigma}{2}) f_{\eta}(\lambda - \frac{\sigma}{2}) dt d\sigma d\lambda dx = \mathcal{O}(\frac{\hbar^3}{\eta \varepsilon}).$$

In addition, according to (3.13), by Corollary A.8, for every $\ell \geq 2$,

$$\int \mathbb{1}\{x \in \mathcal{O}_k \setminus \mathcal{O}_{k-1}\}e^{i\frac{t\sigma}{\hbar}}B_{3,\varepsilon}(x,\lambda,t)f_{\eta}(\lambda + \frac{\sigma}{2})f_{\eta}(\lambda - \frac{\sigma}{2})\mathrm{d}t\mathrm{d}\sigma\mathrm{d}\lambda\mathrm{d}x = \mathcal{O}(\frac{\hbar^{\ell+2}}{\eta^{\ell}\varepsilon_k^{\ell}})$$

using that $|\mho_k| \leq C\varepsilon_k$ and $(\lambda, \sigma) \mapsto f_{\eta}(\lambda + \frac{\sigma}{2})f_{\eta}(\lambda - \frac{\sigma}{2})$ is in S^{η} and supported in $\{|\lambda|, |\sigma| \leq C\eta\}$. Since $B_{3,\varepsilon}$ is supported in $\{x \in \Omega\}$ and $\sum_{k \geq 2} \varepsilon_k^{-\ell} = \mathcal{O}(\varepsilon^{-\ell})$, we deduce

$$\int \mathbb{1}\{x \notin \mathcal{O}_1\} e^{i\frac{t\sigma}{\hbar}} B_{3,\varepsilon}(x,\lambda,t) f_{\eta}(\lambda + \frac{\sigma}{2}) f_{\eta}(\lambda - \frac{\sigma}{2}) dt d\sigma d\lambda dx = \mathcal{O}(\frac{\hbar^{\ell+2}}{\eta^{\ell}\varepsilon^{\ell}}).$$

Going back to (3.14), by combing these estimates, we obtain

$$\|[\widetilde{f}_{\eta}(H_{\hbar}), g_{\varepsilon}(\mathbf{w})]\|_{\mathbf{J}^{2}}^{2} = \mathcal{O}(\frac{\hbar^{2-n}}{n\varepsilon}).$$

This completes the proof of (3.2) in the case where the spectral scale η is smaller than $\hbar^{\frac{1}{2}}$.

Now, we turn to the proof of (3.3) in the same regime. The Hilbert-Schmidt norm of the double commutator $\|[[\tilde{f}_{\eta}(H_{\hbar}), g_{\varepsilon}(\mathbf{w})], k_{\varepsilon}(\mathbf{w})]\|_{\mathbf{J}^{2}}^{2}$ is expressed as in (3.11) with, instead of $B_{1,\varepsilon}$, the symbol

$$\widetilde{B}_{1,\varepsilon}(x,y,\zeta,\omega,r,s,t) = B_0(x,y,\zeta,\omega,r,s,t) \left(g_{\varepsilon}(\mathbf{w}(x)) - g_{\varepsilon}(\mathbf{w}(y))\right)^2 \left(k_{\varepsilon}(\mathbf{w}(x)) - k_{\varepsilon}(\mathbf{w}(y))\right)^2.$$

We can again apply the same stationary phase method; the amplitude $\widetilde{B}_{1,\varepsilon}$ belongs to $S^{\varepsilon}(\mathbb{R}^{4n+2})$ with $\varepsilon \geq \hbar^{\frac{1}{2}}$, so by Proposition A.4, there exists a symbol $\widetilde{B}_{3,\varepsilon} \in S^{\varepsilon}(\mathbb{R}^{n+2})$ such that

$$\|[[\widetilde{f}_{\eta}(H_{\hbar}), g_{\varepsilon}(\mathbf{w})], k_{\varepsilon}(\mathbf{w})]\|_{\mathbf{J}^{2}}^{2} = \frac{1}{(2\pi\hbar)^{n+1}} \int e^{i\frac{\sigma t}{\hbar}} \widetilde{B}_{3,\varepsilon}(x,\lambda,t) f_{\eta}(\lambda + \frac{\sigma}{2}) f_{\eta}(\lambda - \frac{\sigma}{2}) dt d\sigma d\lambda dx + \mathcal{O}(\hbar^{\infty}),$$

and $\widetilde{B}_{3,\varepsilon}$ also satisfies (3.12)–(3.13). In particular, using (3.13) and Corollary A.8 as above, for any fixed $\ell \in \mathbb{N}$,

$$\int \mathbb{1}\{x \notin \mathcal{O}_1\} e^{i\frac{t\sigma}{\hbar}} \widetilde{B}_{3,\varepsilon}(x,\lambda,t) f_{\eta}(\lambda + \frac{\sigma}{2}) f_{\eta}(\lambda - \frac{\sigma}{2}) dt d\sigma d\lambda dx = \mathcal{O}(\frac{\hbar^{\ell+2}}{\eta^{\ell}\varepsilon^{\ell}}).$$

In fact, $\widetilde{B}_{3,\varepsilon}$ satisfies stronger estimates than (3.12). Since $y_{\star}|_{t=0} = x$, we have for any smooth differential operator $L = L_{(y,\zeta,r,s)}$ of degree 2,

$$L\widetilde{B}_{1,\varepsilon}(x,y,\zeta,\omega,r,s,t)\big|_{(y,\zeta,r,s)=(y_{\star},0,r_{\star},0),\,t=0}=0$$

so that, according to (A.3) with $\ell = 2$,

$$B_{3,\varepsilon}(x,\lambda,0) = \mathcal{O}(\hbar^2 \varepsilon^{-4}).$$

Similarly, by (A.3) with $\ell = 1$,

$$\partial_t^2 B_{3,\varepsilon}(x,\lambda,t)\big|_{t=0} = \mathcal{O}(\hbar\varepsilon^{-4}).$$

Hence, instead of (3.15), we apply Corollary A.7 (with $\ell = 4$), and obtain

$$\frac{1}{2\pi\hbar} \int \mathbb{1}\{x \in \mathcal{O}_1\} e^{i\frac{\sigma t}{\hbar}} \widetilde{B}_{3,\varepsilon}(x,\lambda,t) f_{\eta}(\lambda + \frac{\sigma}{2}) f_{\eta}(\lambda - \frac{\sigma}{2}) dt d\sigma d\lambda dx
= \int \mathbb{1}\{x \in \mathcal{O}_1\} \Big(B_{3,\varepsilon}(x,\lambda,0) f_{\eta}(\lambda)^2 + \left(\frac{\hbar}{2\eta}\right)^2 \partial_t^2 B_{3,\varepsilon}(x,\lambda,0) \big(f'_{\eta}(\lambda)^2 - f''_{\eta}(\lambda) f_{\eta}(\lambda) \big) \Big) dx d\lambda + \mathcal{O}(\frac{\hbar^4}{\eta^3 \varepsilon^3}).$$

Since $\eta^2 \leq \hbar$ in this regime and $\|f_{\eta}^{(j)}\|_{L^2(\mathbb{R})}^2 = \mathcal{O}_j(\eta)$ for $j \in \mathbb{N}$, we conclude that

$$\int \mathbb{1}\{x \in \mathcal{O}_1\} e^{i\frac{\sigma t}{\hbar}} \widetilde{B}_{3,\varepsilon}(x,\lambda,t) f_{\eta}(\lambda + \frac{\sigma}{2}) f_{\eta}(\lambda - \frac{\sigma}{2}) dt d\sigma d\lambda dx = \mathcal{O}(\frac{\hbar^5}{\eta^3 \varepsilon^3}).$$

By combining the previous estimates, we conclude that

$$\|[[\widetilde{f}_{\eta}(H_{\hbar}), g_{\varepsilon}(\mathbf{w})], k_{\varepsilon}(\mathbf{w})]\|_{\mathbf{J}^{2}}^{2} = \mathcal{O}(\frac{\hbar^{4}}{\eta^{3}\varepsilon^{3}}).$$

This completes the proof of (3.3) in the case where the spectral scale η is smaller than $\hbar^{\frac{1}{2}}$.

3.3 Hilbert-Schmidt norm of commutators; proof of (3.2)–(3.3) at large spectral scales

This section is concerned with the proof of (3.2) and (3.3) under the assumption that $\eta \geq \hbar^{\frac{1}{2}}$. In this regime, $f_{\eta}(H_{\hbar})$ is a pseudodifferential operator with symbol in S^{η} and small support.

Proposition 3.2. Let $\mathcal{H}(x,\xi) := V(x) + |\xi|^2$ for $(x,\xi) \in \mathbb{R}^{2n}$ and $\chi : \mathbb{R}^n \to [0,1]$ be a smooth function with compact support, and equal to 1 on the ball of radius $\underline{\ell}$. Let \widetilde{f}_{η} be as in Proposition 3.1 for $\eta \in [\sqrt{\overline{h}}, 1]$. Then there exists a symbol $p_{\eta} \in S^{\eta}(\mathbb{R}^{2n})$ such that

$$\widetilde{f}_{\eta}(H_{\hbar}): (x,y) \mapsto \frac{\chi(x-y)}{(2\pi\hbar)^n} \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} p_{\eta}(x,\xi) d\xi + \mathcal{O}_{J^1}(\hbar^{\infty})$$

and for any $\delta \in [\eta, 1]$,

$$p_{\eta} = \mathcal{O}_{S^{\eta}}\left(\left(\frac{\hbar}{n\delta}\right)^{\infty}\right) \quad \text{uniformly on } \{|\mathcal{H}(x,\xi)| > C\delta\}.$$
 (3.16)

In particular, $||p_{\eta}||_{L^{1}_{\xi} \times L^{\infty}_{x}} = \mathcal{O}(\eta)$.

Proof. Recall that according to Proposition 2.2,

$$\widetilde{f}_{\eta}(H_{\hbar}):(x,y)\mapsto \frac{1}{(2\pi\hbar)^{n+1}}\int e^{i\frac{\varphi(t,x,\xi)-y\cdot\xi-t\lambda}{\hbar}}a(x,y,\xi,t)f_{\eta}(\lambda)\mathrm{d}t\mathrm{d}\lambda\mathrm{d}\xi+O_{\mathrm{J}^{1}}(\hbar^{\infty}).$$

We define

$$b: (x, y, \xi) \mapsto \frac{1}{2\pi\hbar} \int e^{i\frac{\Phi(t, \lambda, x, \xi)}{\hbar}} a(x, y, \xi, t) f_{\eta}(\lambda) dt d\lambda, \qquad \Phi(t, \lambda, x, \xi) := \varphi(t, x, \xi) - x \cdot \xi - t\lambda. \tag{3.17}$$

This oscillatory integral satisfies the hypotheses of Proposition A.4: the amplitude belongs to $S_{t,\lambda}^{\eta}(\mathbb{R}^2)$, the phase Φ has a unique critical point $(t,\lambda)=(0,V(x)+|\xi|^2)$, and the Hessian at this point is non-degenerate. Moreover, the amplitude is in S_y^1 , the phase is independent of y. Consequently, there is a symbol $b\in S_{x,\xi}^{\eta}\times S_y^1$ such that

b satisfies (A.8) with
$$\mathcal{A} = \{(x,\xi) \in \mathbb{R}^{2n}, f_{\eta}(V(x) + |\xi|^2) \neq 0\},$$

and we have

$$\widetilde{f}_{\eta}(H_{\hbar})(x,y) = \frac{1}{(2\pi\hbar)^n} \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} b(x,y,\xi) d\xi + O_{J^1}(\hbar^{\infty}).$$

Observe also that since a is supported in a neighborhood of size $\underline{\ell}$ of the diagonal $\{x=y\}$, we can freely include the cutoff $\chi(x-y)$ in the previous kernel. Now, it remains to eliminate the fact that b depends on y. To this end, let $\varkappa : \mathbb{R}^n \to [0,1]$ be a smooth cutoff, with a fixed compact support, such that $\varkappa(\xi) = 1$ on a neighborhood of $\mathcal{B} := \bigcup_{(x,y) \in \mathbb{R}^{2n}} \operatorname{supp} (\xi \mapsto b(x,y,\xi))$, and introduce

$$p_{\eta}: (x,\xi) \mapsto \frac{\varkappa(\xi)}{(2\pi\hbar)^n} \int e^{-i\frac{u\cdot\zeta}{\hbar}} b(x,x+u,\xi+\zeta) \,\mathrm{d}u \mathrm{d}\zeta.$$

Note that we introduce the cutoff \varkappa simply so that p_{η} has a fixed compact support. Then, applying again Proposition A.4, $p_{\eta} \in S^{\eta}(\mathbb{R}^{2n})$ and, according to Remark A.5, p_{η} satisfies (3.16). Finally, by Proposition A.6 (the saddle point is $(z,\xi) = (y,\zeta)$), since $b(x,z,\zeta)\varkappa(\xi)$ is in $S_{z,\xi}^0 \times S_{x,\xi}^{\eta}$, we obtain for any $y \in \mathbb{R}^n$,

$$\int e^{i\frac{(x-y)\cdot\xi}{\hbar}} p_{\eta}(x,\xi) d\xi = \frac{1}{(2\pi\hbar)^n} \int e^{i\frac{(x-y)\cdot\xi + (x-z)\cdot(\zeta - \xi)}{\hbar}} b(x,z,\zeta) \varkappa(\xi) dz d\xi d\zeta$$
$$= \int e^{i\frac{(x-y)\cdot\zeta}{\hbar}} \{b(x,y,\zeta) + \mathcal{O}_{S_{x,\zeta}^{\eta}}(\hbar^{\infty})\} d\zeta$$

where we used that $\varkappa(\xi) = 1$ on \mathcal{B} , so that $\partial_{\xi}^k \varkappa(\xi) = 0$ for every $k \geq 1$ on the support of b. In particular, the error (times the cutoff $\chi(x-y)$) is a kernel in $\mathcal{O}_{S_{x,y}^{\eta}}(\hbar^{\infty})$. Using the estimate (2.6), this corresponds to an operator $\mathcal{O}_{J^1}(\hbar^{\infty})$ as claimed.

Finally, to estimate $||p_{\eta}||_{L^{1}_{\xi}\times L^{\infty}_{x}} = \mathcal{O}(\eta)$, we observe that by assumption, $\partial_{x,\xi}\mathcal{H} \neq 0$ for $x \in \{\mathcal{H} = 0\}$, so that $|\xi \in \mathbb{R}^{n} : |\mathcal{H}(x,\xi)| \leq C\delta| = \mathcal{O}(\delta)$ for any $0 < \delta \leq \underline{c}$ for some small constant \underline{c} . Then, by considering dyadic scales $\delta_{k} = \eta 2^{k}$ for $k \geq 0$, we deduce from the estimate (3.16) that uniformly for $x \in \mathbb{R}^{n}$,

$$\int |p_{\eta}(x,\xi)| d\xi = \mathcal{O}\left(\eta + \sum_{k>1} \frac{\hbar^2}{\eta^2 \delta_k}\right) = \mathcal{O}(\eta)$$

since $\hbar^2 \eta^{-4} \leq 1$. This concludes the proof.

From Proposition 3.2, by applying the stationary phase for mild symbols (in the form of Proposition A.6), one can prove the relevant commutator estimates when the spectral scale η is larger than $\hbar^{\frac{1}{2}}$, in which case we only need to assume that $\varepsilon \eta \geq \hbar$. We focus on (3.2).

Since $f_{\eta}(H_{\hbar})$ and $g_{\epsilon}(\mathbf{w})$ are self-adjoint, from the expression of the kernel of $f_{\eta}(H_{\hbar})$ in Proposition 3.2, we obtain

$$\begin{aligned} \|[f_{\eta}(H_{\hbar}), g_{\varepsilon}(\mathbf{w})]\|_{\mathbf{J}^{2}}^{2} \\ &= \frac{1}{(2\pi\hbar)^{2n}} \int e^{i\frac{(x-y)\cdot(\xi-\zeta)}{\hbar}} p_{\eta}(x,\xi) p_{\eta}(y,\zeta) \chi(x-y)^{2} (g_{\varepsilon}(\mathbf{w}(x)) - g_{\varepsilon}(\mathbf{w}(y)))^{2} d\zeta dy dx d\xi + \mathcal{O}(\hbar^{\infty}). \end{aligned} (3.18)$$

We split this integral in two parts by introducing a cutoff. Let $\chi_1 : \mathbb{R} \to [0,1]$ be a smooth compactly supported function, let $\chi_{\varepsilon} := \chi(\varepsilon^{-1}\mathbf{w})$ and $\chi_{\varepsilon}^{\dagger} := 1 - \chi_{\varepsilon}$ be such that $\chi_{\varepsilon}^{\dagger} \cdot g_{\varepsilon}(\mathbf{w}) = 0$. We have

$$||[f_{\eta}(H_{\hbar}), g_{\varepsilon}(\mathbf{w})]||_{\mathbf{J}^{2}}^{2} = \frac{1}{(2\pi\hbar)^{2n}} \int p_{\eta}(x,\xi) \left(\int e^{i\frac{(x-y)\cdot(\xi-\zeta)}{\hbar}} p_{\eta}(y,\zeta)Q(x,y)\mathrm{d}\zeta\mathrm{d}y \right) \mathrm{d}x\mathrm{d}\xi + \frac{1}{(2\pi\hbar)^{n}} \int p_{\eta}(y,\zeta)g_{\varepsilon}(\mathbf{w}(y))^{2}R(y,\zeta)\mathrm{d}\zeta\mathrm{d}y + \mathcal{O}(\hbar^{\infty})$$
(3.19)

where

$$Q: (x,y) \mapsto \chi(x-y)^2 \chi_{\varepsilon}(x) (g_{\varepsilon}(\mathbf{w}(x)) - g_{\varepsilon}(\mathbf{w}(y)))^2$$
$$R: (y,\zeta) \mapsto \frac{1}{(2\pi\hbar)^n} \int e^{i\frac{(x-y)\cdot(\xi-\zeta)}{\hbar}} \chi(x-y)^2 p_{\eta}(x,\xi) \chi_{\varepsilon}^{\dagger}(x) dx d\xi.$$

The relevant property is that, with $\ell = 2$,

$$(y,\zeta) \mapsto \varepsilon^{-1} p_{\eta}(y,\zeta) \int Q(x,y) dx$$
 is in $S_y^{\varepsilon} \times S_{\zeta}^{\eta}$ with $\varepsilon \eta \ge \hbar$ and $\partial_y^k Q(x,y)|_{y=x} = 0$ for $k < \ell$. (3.20)

Then, applying Corollary A.7 with $\ell = 2$, we obtain (uniformly for $\xi \in \mathbb{R}^n$)

$$\frac{1}{(2\pi\hbar)^n}\int e^{i\frac{(x-y)\cdot(\xi-\zeta)}{\hbar}}p_{\eta}(y,\zeta)Q(x,y)\mathrm{d}\zeta\mathrm{d}y\mathrm{d}x=\mathcal{O}\big(\tfrac{\hbar^2}{\eta^2\varepsilon}\big).$$

Since

$$\sup_{x \in \mathbb{R}^n} \int |p_{\eta}(x,\xi)| d\xi = \mathcal{O}(\eta),$$

this implies that $(3.19) = \mathcal{O}(\frac{\hbar^{2-n}}{\eta \varepsilon})$.

Then, the amplitude of the integral R is in $S_x^{\delta} \times S_{\xi}^{\eta}$ with $\delta = \min\{\varepsilon, \eta\}$, so applying either Proposition A.6 in case $\delta = \varepsilon$ or Lemma A.2 in case $\delta = \eta$ with $\eta \geq \hbar^{\frac{1}{2}}$ (here, $\chi_{\varepsilon}^{\dagger}(x) = 0$ for $x \in B(y, \varepsilon)$ if $y \in \{g_{\varepsilon}(w) \neq 0\}$), we obtain in both cases

$$R(y,\zeta) = \mathcal{O}\left(\left(\frac{\hbar}{\eta\varepsilon}\right)^{\infty}\right)$$

with the required uniformity. Thus, this implies that the second line of (3.19) is $\mathcal{O}((\frac{\hbar}{\eta\varepsilon})^{\infty}\hbar^{-n}\eta\varepsilon)$. Combining these estimates, this concludes the proof of (3.2).

Now, we turn to the estimate (3.3); the argument being the same. Using Proposition 3.2, we also have

$$\|[[f_{\eta}(H_{\hbar}), g_{\varepsilon}(\mathbf{w})], k_{\varepsilon}(\mathbf{w})]\|_{\mathbf{J}^{2}}^{2} = \frac{1}{(2\pi\hbar)^{2n}} \int p_{\eta}(x, \xi) \left(\int e^{i\frac{(x-y)\cdot(\xi-\zeta)}{\hbar}} p_{\eta}(y, \zeta) \widetilde{Q}(x, y) d\zeta dy \right) dx d\xi + \frac{1}{(2\pi\hbar)^{n}} \int p_{\eta}(y, \zeta) g_{\varepsilon}(\mathbf{w}(y))^{2} k_{\varepsilon}(\mathbf{w}(y))^{2} \widetilde{R}(y, \zeta) d\zeta dy + \mathcal{O}(\hbar^{\infty})$$
(3.21)

where

$$\widetilde{Q}: (x,y) \mapsto \chi(x-y)^2 \chi_{\varepsilon}(x) (g_{\varepsilon}(\mathbf{w}(x)) - g_{\varepsilon}(\mathbf{w}(y)))^2 (k_{\varepsilon}(\mathbf{w}(x)) - k_{\varepsilon}(\mathbf{w}(y)))^2$$

$$\widetilde{R}: (y,\zeta) \mapsto \frac{1}{(2\pi\hbar)^n} \int e^{i\frac{(x-y)\cdot(\xi-\zeta)}{\hbar}} \chi(x-y)^2 p_{\eta}(x,\xi) \chi_{\varepsilon}^{\dagger}(x) dx d\xi.$$

Now \widetilde{Q} satisfies the condition (3.20) with $\ell=4$. Thus, the first term of the right-hand-side of (3.21) is $\mathcal{O}(\frac{\hbar^{4-n}}{\eta^3\varepsilon^3})$ and the integral involving \widetilde{R} is again $\mathcal{O}((\frac{\hbar}{\eta\varepsilon})^\infty\hbar^{-n}\eta\varepsilon)$. This yields the estimate (3.3).

4 Hilbert-Schimdt norm of commutators: Proof of Theorem 1.

Throughout this section, we use the notation from Section 2.1. In particular, $\Omega \in \mathcal{D}$ is a fixed relatively compact open subset of the bulk with smooth boundary, $f \in C^{\infty}(\mathbb{R}^n)$, and we denote $f|_{\Omega} = f\mathbb{1}_{\Omega}$ (this function is also viewed as a bounded operator on $L^2(\mathbb{R}^n)$). Using Proposition 2.5, we focus on analysing the

commutator $[\widetilde{\Pi}, f|_{\Omega}]$ where the regularized kernel $\widetilde{\Pi}$ is given by the oscillatory integral (2.3). Then, with the phase (3.5), one has

$$\|[\widetilde{\Pi}, f|_{\Omega}]\|_{J^{2}}^{2} = \operatorname{tr}\left([\widetilde{\Pi}, f|_{\Omega}][f|_{\Omega}, \widetilde{\Pi}^{*}]\right) = \frac{1}{(2\pi\hbar)^{2n+2}} \int e^{i\frac{\Psi_{0}(x,y,\xi_{1},\xi_{2},t_{1},t_{2},\lambda_{1},\lambda_{2})}{\hbar}} A_{0}(x,y,\xi_{1},\xi_{2},t_{1},t_{2},\lambda_{1},\lambda_{2})$$

$$(f|_{\Omega}(x) - f|_{\Omega}(y))^{2} \mathbb{1}_{\lambda_{1} \leq \mu} \mathbb{1}_{\lambda_{2} \leq \mu} dt_{1} dt_{2} d\lambda_{1} d\lambda_{2} d\xi_{1} d\xi_{2} dx dy,$$

$$(4.1)$$

where the amplitude A_0 belongs to $S^1(\mathbb{R}^{6n+4})$ and its principal part at $(t_1, t_2, y) = (0, 0, x)$ is given by

$$A_0(x, x, \xi_1, \xi_2, 0, 0, \lambda_1, \lambda_2)|_{\hbar=0} = \vartheta(V(x) + |\xi_1|^2)\vartheta(V(x) + |\xi_2|^2)\chi(\lambda_1)\chi(\lambda_2). \tag{4.2}$$

Moreover, since $\Omega \in \{V < 0\}$, the function $(x, y) \mapsto (f|_{\Omega}(x) - f|_{\Omega}(y))^2 \mathbb{1}\{|x - y| \le \underline{\ell}\}$ is supported in $\Omega' \times \Omega'$ where $\Omega' \subset \{V < -2\underline{c}\}$ is a neighbourhood of Ω in the bulk. Consequently, by (2.1), A_0 is supported in

$$\{t \in [-\underline{\tau}, \underline{\tau}], (x, y) \in \Omega' \times \Omega', |x - y| \le \underline{\ell}, |\xi_1|, |\xi_2| \ge \underline{c}, |\lambda_1|, |\lambda_2| \le \underline{\ell}\}. \tag{4.3}$$

for small constants $\underline{\tau} \ll \underline{\ell} \ll \underline{c}$.

The proof consists of the following steps.

- In Section 4.1, we gather some preliminary estimates for singular integrals with discontinuous amplitudes.
- In Section 4.2, using the stationary phase method, we perform a series of reductions to write the oscillatory integral (4.1) as

$$\|[\widetilde{\Pi}, f|_{\Omega}]\|_{\mathcal{J}^{2}}^{2} = \frac{-1}{(2\pi\hbar)^{2n}} \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} F(x, y, \xi) (f|_{\Omega}(x) - f|_{\Omega}(y))^{2} d\xi dx dy + \mathcal{O}(\hbar^{1-n}). \tag{4.4}$$

where the function F has compact support, is smooth on $\mathbb{R}^3 \setminus \{\xi = 0\}$ and $|F(x, y, \xi)| \leq C|\xi|$.

• In Section 4.3, we study integrals of the form (4.4) in the particular case where the set Ω is C^{∞} diffeomorphic to the ball $B = \{x \in \mathbb{R}^n : |x| < 1\}$. After a change of variables and further reduction steps,
we show that

$$(4.4) = \frac{1}{(2\pi\hbar)^{2n}} \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} g(\frac{x+y}{2}, \xi) \mathbb{1}_B(x) \mathbb{1}_B(y) d\xi dx dy + \mathcal{O}(\hbar^{1-n}),$$

where g is smooth on $\mathbb{R}^{2n} \setminus \{\xi = 0\}$ and $g(v, \xi) = |\xi| R(v) + \mathcal{O}(|\xi|^2)$ as $\xi \to 0$. Using the asymptotics of the Fourier transform $\hat{\mathbb{1}}_B$, which is related to Bessel functions, we obtain that the leading term of (4.4) is given by $(2\pi\hbar)^{1-n} \frac{\log \hbar^{-1}}{\pi^2} \int_{\partial B} R(\hat{x}) f(\hat{x})^2 d\hat{x}$.

- Finally, in Section 4.4, we use a partition of unity type argument to conclude the proof of Theorem 1 without the topological assumption.
- In Section 4.5, we prove Theorem 2. The Gaussian asymptotic fluctuations for counting statistics is a direct consequence of Theorem 1 (the variance of $\mathbf{X}(\Omega)$ diverges for any smooth set Ω) and we only need to estimate covariances. Using the off-diagonal decay of the regularized kernel (Proposition 4.17), we argue that for two smooth sets $\Omega_1, \Omega_2 \in \mathcal{D}$ such that $|\partial \Omega_1 \cap \partial \Omega_2| = 0$, the covariance satisfies $\operatorname{tr}([\Pi, \mathbb{1}_{\Omega_1}][\mathbb{1}_{\Omega_2}, \Pi]) = o(\log(\hbar^{-1})\hbar^{1-n})$ as $\hbar \to 0$.

4.1 Preliminary estimates

In this section we gather several estimates that will be important for the proof of Theorem 2.

Lemma 4.1. There exists a constant C_n such that the following is true: let $\Omega \in \mathbb{R}^n$ be a bounded open set with a smooth boundary. Denote by $|\partial\Omega|$ the (n-1)-Hausdorff measure of the smooth, compact hypersurface $\partial\Omega$. Then for every $\hbar > 0$ sufficiently small,

$$\int_{|x-y| \le \hbar} \frac{|\mathbb{1}_{\Omega}(x) - \mathbb{1}_{\Omega}(y)|^2}{2} \mathrm{d}x \mathrm{d}y = \int_{|x-y| \le \hbar} \mathbb{1}_{\Omega}(x) \mathbb{1}_{\Omega^c}(y) \mathrm{d}x \mathrm{d}y \sim C_n |\partial\Omega| \hbar^{n+1}.$$

For any $\varkappa : \mathbb{R}^{2n} \to [0,1]$ with compact support,

$$\int \frac{|\mathbb{1}_{\Omega}(x) - \mathbb{1}_{\Omega}(y)|^2}{2|x - y|^n} \varkappa(x, y) \mathrm{d}x \mathrm{d}y = \int \frac{\mathbb{1}_{\Omega}(x) \mathbb{1}_{\Omega^c}(y)}{|x - y|^n} \varkappa(x, y) \mathrm{d}x \mathrm{d}y < \infty. \tag{4.5}$$

Moreover, given $\overline{\chi}: \mathbb{R}^n \to [0,1]$ smooth such that $\overline{\chi}(x) = 0$ for $|x| \leq \underline{c}$, there exists a constant C so that for any $\hbar > 0$,

$$\int \overline{\chi}(\frac{x-y}{\hbar}) \frac{|\mathbb{1}_{\Omega}(x) - \mathbb{1}_{\Omega}(y)|^2}{2|x-y|^{n+2}} dxdy = \int \overline{\chi}(\frac{x-y}{\hbar}) \frac{\mathbb{1}_{\Omega}(x)\mathbb{1}_{\Omega^c}(y)}{|x-y|^{n+2}} dxdy \le C/\hbar.$$

$$(4.6)$$

Proof. Let

$$\Omega_{\hbar} = \{ x \in \Omega : \operatorname{dist}(x, \partial \Omega) < \hbar \}$$

be an \hbar -neighborhood of $\partial\Omega$ inside Ω . Since $\partial\Omega$ is smooth and compact, if \hbar is small enough, then the following change of variables is well defined

$$\Omega_{\hbar} \mapsto \partial\Omega \times (0,1)$$

$$x \mapsto (\hat{x}, t) = (\underset{q \in \partial\Omega}{\operatorname{arg\,min}} |q - x|^2, \hbar^{-1} \operatorname{dist}(x, \partial\Omega)).$$

This corresponds to the (orthogonal) projection of $x \in \Omega_{\hbar}$ onto $\partial\Omega$ and the volume form $\mathrm{d}x \sim \hbar \mathrm{d}t \mathrm{d}\hat{x}$ as $\hbar \to 0$ where $\mathrm{d}\hat{x}$ denotes the volume measure on $\partial\Omega$. Hence, as $\hbar \to 0$

$$\int \mathbb{1}_{\Omega}(x) \mathbb{1}_{|x-y| \le \hbar} \mathbb{1}_{\Omega^c}(y) dx dy \sim \hbar \int_{\partial \Omega \times (0,1)} \left(\int_{\Omega^c} \mathbb{1}_{|\hat{x}-\hbar t \nu_{\Omega}(\hat{x}) - y| \le \hbar} dy \right) d\hat{x} dt$$

where $\nu_{\Omega}(\hat{x})$ denotes the (exterior) unit normal vector at $\hat{x} \in \Omega$.

For a fixed $\hat{x} \in \partial \Omega$ and t > 0,

$$\int_{\Omega^c} \mathbb{1}_{|\hat{x} - \hbar t \nu_{\Omega} - y| \le \hbar} \mathrm{d}y \sim \hbar^n \int_{\{y \cdot \mathbf{e}_1 \ge 0\}} \mathbb{1}_{|y + t \mathbf{e}_1| \le 1} \mathrm{d}y$$

which follows by rescaling and using normal coordinates around \hat{x} . This shows that there exists a constant C_n so that as $\hbar \to 0$,

$$\int_{|x-y| \le \hbar} \mathbb{1}_{\Omega}(x) \mathbb{1}_{\Omega^c}(y) dx dy \sim C_n \hbar^{n+1} \int_{\partial \Omega} d\hat{x}.$$

To obtain (4.5), observe that for $s \in \mathbb{R}_+$

$$\int \frac{\mathbb{1}_{\Omega}(x)\mathbb{1}_{\Omega^{c}}(y)}{|x-y|^{s}} \varkappa(x,y) \mathrm{d}x \mathrm{d}y \leq \sum_{k \in \mathbb{Z}: 2^{k} \leq C/\hbar} (\hbar 2^{k})^{-s} \int_{|x-y| \leq \hbar 2^{k+1}} \mathbb{1}_{\Omega}(x)\mathbb{1}_{\Omega^{c}}(y) \mathrm{d}x \mathrm{d}y$$

where the constant C depends only on \varkappa . The pervious argument shows that the integrals on the RHS are $\mathcal{O}((\hbar 2^k)^{n+1})$ so that

$$\int \frac{\mathbb{1}_{\Omega}(x)\mathbb{1}_{\Omega^c}(y)}{|x-y|^s} \varkappa(x,y) \mathrm{d}x \mathrm{d}y \le C\hbar^{n+1-s} \sum_{k \in \mathbb{Z}: 2^k \le C/\hbar} 2^{k(n+1-s)} = O(1)$$

provided that s < n + 1; in which case the sum is geometrically growing. On the other hand, for s > n + 1, upon adding a different cutoff to exclude that diagonal, we obtain

$$\int \overline{\chi}(\frac{x-y}{\hbar}) \frac{\mathbb{1}_{\Omega}(x)\mathbb{1}_{\Omega^c}(y)}{|x-y|^s} \mathrm{d}x \mathrm{d}y \le \sum_{k \in \mathbb{N}_0} (\hbar 2^k)^{-s} \int_{|x-y| \le \hbar 2^{k+1}} \mathbb{1}_{\Omega}(x)\mathbb{1}_{\Omega^c}(y) \mathrm{d}x \mathrm{d}y \le C \hbar^{n+1-s}$$

since the series is convergent. This completes the proof of (4.6).

Lemma 4.1 has the two following consequences.

Proposition 4.2. Let $\Omega \in \mathbb{R}^n$ be a relatively compact open set with a smooth boundary, $f \in C^{\infty}(\mathbb{R}^n)$ and recall that $f|_{\Omega} = f\mathbb{1}_{\Omega}$. For any $\varkappa : \mathbb{R}^{2n} \to [0,1]$ with compact support,

$$\int \frac{|f|_{\Omega}(x) - f|_{\Omega}(y)|^2}{2|x - y|^n} \varkappa(x, y) dx dy < \infty.$$

Moreover, for any cutoff $\chi \in C_c^{\infty}$,

$$\int \chi(\frac{x-y}{\hbar})(f|_{\Omega}(x) - f|_{\Omega}(y))^2 dxdy = \mathcal{O}(\hbar^{n+1}).$$
(4.7)

Proof. We have

$$|f|_{\Omega}(x) - f|_{\Omega}(y)|^2 = (f(x) - f(y))(f|_{\Omega}(x) - f|_{\Omega}(y)) + f(x)f(y)(\mathbb{1}_{\Omega}(x)\mathbb{1}_{\Omega^c}(y) + \mathbb{1}_{\Omega^c}(x)\mathbb{1}_{\Omega}(y)).$$

Then according to Lemma 4.1 (4.5),

$$\int \frac{|f|_{\Omega}(x) - f|_{\Omega}(y)|^2}{|x - y|^n} \varkappa(x, y) dxdy = \int \frac{(f(x) - f(y))(f|_{\Omega}(x) - f|_{\Omega}(y))}{|x - y|^n} \varkappa(x, y) dxdy + \mathcal{O}(1).$$

Since f is smooth, the first term is controlled by $\int \frac{\varkappa(x,y)}{|x-y|^{n-1}} \mathrm{d}x \mathrm{d}y < \infty$.

Similarly, by Cauchy-Schwarz's inequality,

$$\int \chi(\frac{x-y}{\hbar})(f|_{\Omega}(x) - f|_{\Omega}(y))^{2} dxdy$$

$$\leq \sqrt{\int \chi(\frac{x-y}{\hbar})(f(x) - f(y))^{2} dxdy} \sqrt{\int \chi(\frac{x-y}{\hbar})(f|_{\Omega}(x) - f|_{\Omega}(y))^{2} dxdy} + \mathcal{O}(\hbar^{n+1})$$

$$= \sqrt{\mathcal{O}(\hbar^{n+2}) \int \chi(\frac{x-y}{\hbar})(f|_{\Omega}(x) - f|_{\Omega}(y))^{2} dxdy} + \mathcal{O}(\hbar^{n+1})$$

where we used that $\int \chi(\frac{x-y}{\hbar})(f(x)-f(y))^2 dx d = \mathcal{O}(\hbar^{n+2})$ for f Lipchitz-continuous. This inequality implies (4.7), otherwise upon dividing by $\sqrt{\int \chi(\frac{x-y}{\hbar})(f|\Omega(x)-f|\Omega(y))^2 dx dy}$ we obtain a contradiction.

Let us also record that by (4.6) and an analogous argument, given a cutoff $\overline{\chi}$, we have

$$\int \overline{\chi}(\frac{x-y}{\hbar}) \frac{|f|_{\Omega}(x) - f|_{\Omega}(y)|^2}{|x-y|^{n+2}} \mathrm{d}x \mathrm{d}y \leq \sqrt{\int \frac{|f(x) - f(y)|^2}{|x-y|^{n+2}} \mathrm{d}x \mathrm{d}y} \sqrt{\int \overline{\chi}(\frac{x-y}{\hbar}) \frac{|f|_{\Omega}(x) - f|_{\Omega}(y)|^2}{|x-y|^{n+2}} \mathrm{d}x \mathrm{d}y} + \mathcal{O}(\hbar^{-1}).$$

Since $\int \frac{|f(x) - f(y)|^2}{|x - y|^{n+2}} dxdy < \infty$, we conclude that

$$\int \overline{\chi}(\frac{x-y}{\hbar}) \frac{|f|_{\Omega}(x) - f|_{\Omega}(y)|^2}{|x-y|^{n+2}} dxdy = \mathcal{O}(\hbar^{-1}).$$
(4.8)

Lemma 4.3. Let $\Omega_1, \Omega_2 \in \mathbb{R}^n$ be two relatively compact open sets with smooth boundary such that the (n-1)-Hausdorff measure of the intersection $\partial \Omega_1 \cap \partial \Omega_2$ is zero. Then as $\hbar \to 0$,

$$\int \frac{|(\mathbb{1}_{\Omega_1}(x) - \mathbb{1}_{\Omega_1}(y))(\mathbb{1}_{\Omega_2}(x) - \mathbb{1}_{\Omega_2}(y))|}{(\hbar + |x - y|)^{n+1}} dxdy = o(\log(\hbar^{-1})).$$
(4.9)

If in fact there exists $\beta < n-1$ such that $H^{\beta}(\partial \Omega_1 \cap \partial \Omega_2) < +\infty$, then

$$\int \frac{|(\mathbb{1}_{\Omega_1}(x) - \mathbb{1}_{\Omega_1}(y))(\mathbb{1}_{\Omega_2}(x) - \mathbb{1}_{\Omega_2}(y))|}{(\hbar + |x - y|)^{n+1}} dxdy = \mathcal{O}(1). \tag{4.10}$$

Proof. We claim that as $\eta \to 0$, the integral

$$\int_{|x-y| \le \eta} (\mathbb{1}_{\Omega_1}(x) - \mathbb{1}_{\Omega_1}(y)) (\mathbb{1}_{\Omega_2}(x) - \mathbb{1}_{\Omega_2}(y)) dx dy = o(\eta^{1+n}). \tag{4.11}$$

Indeed, since the integrand is supported on $\{x \in \mathbb{R}^n : , \operatorname{dist}(x, \partial \Omega_1 \cap \partial \Omega_2) < \eta\}$ and $H^{n-1}(\partial \Omega_1 \cap \partial \Omega_2) = 0$, by a volume estimate we obtain

$$|\{(x,y) \in \mathbb{R}^{2n} : \operatorname{dist}(x,\partial\Omega_1 \cap \partial\Omega_2), \operatorname{dist}(x,y) < \eta\}| = o(\eta^{1+n}).$$

Moreover, for every $\hbar_0 > 0$,

$$\int_{|x-y|>\hbar_0} \frac{(\mathbb{1}_{\Omega_1}(x) - \mathbb{1}_{\Omega_1}(y))(\mathbb{1}_{\Omega_2}(x) - \mathbb{1}_{\Omega_2}(y))}{|x-y|^{n+1}} \mathrm{d}x \mathrm{d}y \le C/\hbar_0$$

The result follows directly form these estimates by a dyadic decomposition; one has for $\hbar \leq \hbar_0$,

$$(4.9) \leq \sum_{k < \log(1/\hbar)} \int_{\hbar 2^{k-1} \leq |x-y| \leq \hbar 2^k} \frac{|(\mathbb{1}_{\Omega_1}(x) - \mathbb{1}_{\Omega_1}(y))(\mathbb{1}_{\Omega_2}(x) - \mathbb{1}_{\Omega_2}(y))|}{|x-y|^{n+1}} dx dy + \mathcal{O}(\hbar_0^{-1})$$

Using (4.11), with $\eta = \hbar 2^k$, these integrals are all o(1) uniformly for $k \leq \log(\hbar^{-1})$, $\hbar \leq \hbar_0$, as $\hbar_0 \to 0$. This shows that (4.9) = $o(\log(\hbar^{-1}))$ by choosing \hbar_0 sufficiently small.

Under any positive improvement on the Hausdorff dimension of $\partial\Omega_1 \cap \partial\Omega_2$, one has instead, for some $\epsilon > 0$,

$$|\{(x,y) \in \mathbb{R}^{2n} : \operatorname{dist}(x,\partial\Omega_1 \cap \partial\Omega_2), \operatorname{dist}(x,y) < \eta\}| = o(\eta^{1+n+\epsilon}),$$

and now the estimates of the dyadic decomposition form a convergent series. This concludes the proof. \Box

As we already alluded to, the following class of functions plays a key role when bounding the integral (4.1).

Definition 4.4 (Class \mathcal{F}). Let $m, n \in \mathbb{N}_0$, define the class

$$\mathcal{F}(\mathbb{R}^m, \mathbb{R}^n) = \{ f : \mathbb{R}^m \times \mathbb{R}^n \to \mathbb{R} \text{ continuous with compact support}; x \mapsto f(x, \xi) \text{ is smooth for } \xi \in \mathbb{R}^n, \text{ and for every } j, k \in \mathbb{N}_0, \ \|\partial_{\xi}^k \partial_x^j f(x, \xi)\| \le C_{j,k} |\xi|^{1-k} \text{ for all } (x, \xi) \in \mathbb{R}^m \times \mathbb{R}^n \}.$$

We record the following two lemmas without proofs as they are direct consequences of this definition.

Lemma 4.5. Functions in \mathcal{F} are Lipschitz continuous with respect to ξ and there are smooth on $\mathbb{R}^n \setminus \{0\}$. If $g \in C_c^{\infty}(\mathbb{R}^m \times \mathbb{R}^n \times \mathbb{R})$ with $g(x, \xi, 0) = 0$, then $f(x, \xi) = g(x, \xi, |\xi|)$ is in the class \mathcal{F} . We also emphasize that the class \mathcal{F} is stable under smooth change of variables on $\mathbb{R}^m \times \mathbb{R}^n$ which coincide with the identity outside of a compact set.

Lemma 4.6. Let $a \in \mathcal{F}(\mathbb{R}^m, \mathbb{R}^n)$. For any cutoff $\chi \in C_c^{\infty}(\mathbb{R}^n)$, as $\hbar \to 0$,

$$\int \chi(\frac{\xi}{\hbar})|a(x,\xi)|d\xi dx = \mathcal{O}(\hbar^{n+1}).$$

The last lemma is complemented by the following claim on the zone $\{|\xi| \ge \hbar\}$.

Lemma 4.7. Let $n, k \in \mathbb{N}$ with $k \leq n+2$. Let $b : \mathbb{R}^{3n} \to \mathbb{R}$ have compact support, smooth with respect to ξ on $\mathbb{R}^n \setminus \{0\}$ and such that for every $j \in \mathbb{N}_0$, and $\xi \in \mathbb{R}^n$,

$$\int \frac{\|\partial_{\xi}^{j}b(x,y,\xi)\|}{|x-y|^{n+2-k}} \mathrm{d}x \mathrm{d}y \le |\xi|^{1-j-k}, \qquad \text{for } \xi \in \mathbb{R}^{n}.$$

$$(4.12)$$

Let $\chi \in C^{\infty}(\mathbb{R}^{3n})$ be such that, for every $x, y, \xi \in \mathbb{R}^n$,

$$\mathbb{1}_{|\xi|<1} \le \chi(x,y,\xi) \le \mathbb{1}_{|\xi|<2}.$$

Then as $\hbar \to 0$,

$$\int e^{i\frac{(x-y)\cdot\xi}{\hbar}} (1 - \chi(x,y,\frac{\xi}{\hbar}))b(x,y,\xi)d\xi dxdy = \mathcal{O}(\hbar^{n+1-k}).$$

Proof. We will prove this claim by decreasing order of the value of k: at k = n + 2, the claim follows from the simple fact that, for any fixed R > 0, as $\hbar \to 0$, one has

$$\int_{\hbar \le |\xi| \le R} |\xi|^{-n-1} = \mathcal{O}(\hbar^{-1}).$$

Suppose now that the claim holds for some value of k and let us prove it for k-1; to this end we integrate by parts

$$\int e^{i\frac{(x-y)\cdot\xi}{\hbar}} (1-\chi(x,y,\frac{\xi}{\hbar}))b(x,y,\xi)\mathrm{d}\xi\mathrm{d}x\mathrm{d}y$$

$$= i\hbar \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} (1-\chi(x,y,\frac{\xi}{\hbar}))\frac{(x-y)\cdot\partial_{\xi}b(x,y,\xi)}{\|x-y\|^2}\mathrm{d}\xi\mathrm{d}x\mathrm{d}y$$

$$-i\int e^{i\frac{(x-y)\cdot\xi}{\hbar}} (\partial_{\xi}\chi)(x,y,\frac{\xi}{\hbar})\cdot\frac{(x-y)b(x,y,\xi)}{\|x-y\|^2}\mathrm{d}\xi\mathrm{d}x\mathrm{d}y$$

Note that the first term in the equation is exactly of the form given before, where now $B = \frac{(x-y)\cdot\partial_{\xi}b(x,y,\xi)}{\|x-y\|^2}$ satisfies (4.12); thus

$$i\hbar \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} (1-\chi(x,y,\frac{\xi}{\hbar})) \frac{(x-y)\cdot\partial_{\xi}b(x,y,\xi)}{|x-y|^2} d\xi dx dy = \mathcal{O}(\hbar^{n+2-k}).$$

It remains to bound

$$i \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} (\partial_{\xi}\chi)(x,y,\frac{\xi}{\hbar}) \cdot \frac{(x-y)b(x,y,\xi)}{|x-y|^2} d\xi dx dy = \mathcal{O}\left(\int_{\hbar \le |\xi| \le 2\hbar} |\xi|^{2-k}\right) = \mathcal{O}(\hbar^{n+1-k}).$$

This concludes the proof.

Finally, we also need an estimate for the commutator between a pseudodifferential operators with smooth, compactly supported symbols and a function with jump discontinuities. This is to be compared with Proposition 2.6, albeit simpler.

Lemma 4.8. Let $f \in C^{\infty}(\mathbb{R}^n)$, $a \in S^1(\mathbb{R}^{3n})$ and $K : (x,y) \mapsto \frac{1}{(2\pi\hbar)^n} \int e^{i\frac{\xi \cdot (x-y)}{\hbar}} a(x,y,\xi) d\xi$. Then, for any bounded open set $\Omega \in \mathbb{R}^n$ with smooth boundary,

$$||[f|_{\Omega}, K]||_{\mathbf{J}^2}^2 = O(\hbar^{1-n}).$$

Proof. By definition, the Hilbert-Schmidt norm,

$$\|[f|_{\Omega}, K]\|_{J^{2}}^{2} = \frac{1}{(2\pi\hbar)^{2n}} \int e^{i\frac{(x-y)\cdot(\xi_{1}-\xi_{2})}{\hbar}} (f|_{\Omega}(x) - f|_{\Omega}(y))^{2} a(x, y, \xi_{1}) \,\overline{a}(y, x, \xi_{2}) \,\mathrm{d}\xi_{1} \,\mathrm{d}\xi_{2} \,\mathrm{d}x \,\mathrm{d}y.$$

We make a change of variable $\xi = \xi_1 - \xi_2$ and $\zeta = \frac{\xi_1 + \xi_2}{2}$ so that

$$\|[f|_{\Omega}, K]\|_{\mathcal{J}^2}^2 = \frac{1}{(2\pi\hbar)^{2n}} \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} (f|_{\Omega}(x) - f|_{\Omega}(y))^2 b(x, y, \xi, \zeta) \,\mathrm{d}\xi \,\mathrm{d}\zeta \,\mathrm{d}x \,\mathrm{d}y$$

where $b \in S^1(\mathbb{R}^{4n})$. Using the bound (4.7), up to an error $\mathcal{O}(\hbar^{1-n})$, we can introduce a cutoff excluding the diagonal in the previous integral. Then, we perform integrations by parts with respect to ξ ; we obtain for any $k \in \mathbb{N}_0$

$$\|[f|_{\Omega},K]\|_{J^2}^2 = \frac{(i\hbar)^k}{(2\pi\hbar)^{2n}} \int \overline{\chi}(\frac{x-y}{\hbar}) e^{i\frac{(x-y)\cdot\xi}{\hbar}} (f|_{\Omega}(x) - f|_{\Omega}(y))^2 \frac{(x-y)^{\otimes k} \cdot \partial_{\xi}^k b(x,y,\xi,\zeta)}{|x-y|^{2k}} d\xi d\zeta dx dy + \mathcal{O}(\hbar^{1-n}).$$

We can bound for $(x, y, \xi, \zeta) \in \mathbb{R}^{4n}$ with k = n + 2,

$$\left| \frac{(x-y)^{\otimes k} \cdot \partial_{\xi}^{k} b(x, y, \xi, \zeta, \hbar)}{|x-y|^{2k}} \right| \leq \frac{\varkappa(\xi, \zeta)}{|x-y|^{n+2}}$$

where $\varkappa \in C_c^{\infty}(\mathbb{R}^{2n})$. Thus, by (4.8), we conclude that $\|[f|_{\Omega}, K]\|_{J^2}^2 = O(\hbar^{1-n})$.

4.2 Reduction to an oscillatory integral

Starting from the expression (4.1), using the stationary phase method and certain well-chosen changes of variables, we can prove the following result.

Proposition 4.9. Under the assumptions from Section 2.1, there is a symbol $F \in \mathcal{F}(\mathbb{R}^{2n}, \mathbb{R}^n)$ (independent of Ω) such that as $\hbar \to 0$,

$$\|[\widetilde{\Pi}, f|_{\Omega}]\|_{J^{2}}^{2} = \frac{-1}{(2\pi\hbar)^{2n}} \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} F(x, y, \xi) (f|_{\Omega}(x) - f|_{\Omega}(y))^{2} d\xi dx dy + \mathcal{O}(\hbar^{1-n}).$$

Moreover, the principal part of F satisfies uniformly as $\xi \to 0$,

$$F(x, x, \xi)|_{\hbar=0} = c_{n-1}|\xi||V(x)|^{\frac{n-1}{2}} + \mathcal{O}(|\xi|^2)$$
(4.13)

with the constant c_{n-1} as in Theorem 1 for $n \in \mathbb{N}$.

Starting from formula (4.1), let us explain the main steps of the proof of Proposition 4.9.

• Step 1. As in Section 3.1, we perform a change of coordinates similar to (3.7)

$$\begin{cases} \lambda = \frac{\lambda_1 + \lambda_2}{2}, & \xi_1 = r_1 \omega_1 \\ \sigma = \lambda_2 - \lambda_1, & \xi_2 = r_2 \omega_2 \end{cases} \quad \text{with } r_1, r_2 > \underline{c}, \, \omega_1, \omega_2 \in \mathbb{S}^{n-1}.$$
 (4.14)

This is justified because of the support condition (4.3), which ensures that ξ_1 and ξ_2 are bounded away from zero. Observe that $\mathbb{1}_{\lambda_1 \leq 0} \mathbb{1}_{\lambda_2 \leq 0} = \mathbb{1}_{|\sigma| \leq -2\lambda}$ under this change of variables. Then, we can perform a stationary phase with respect to the variables (t_1, r_1, t_2, r_2) , keeping $(x, y, \omega_1, \omega_2, \lambda, \sigma)$ fixed, and we obtain an integral of the type

$$\frac{1}{(2\pi\hbar)^{2n}} \int e^{i\frac{\Psi_1(x,y,\omega_1,\omega_2,\lambda,\sigma)}{\hbar}} (f|_{\Omega}(x) - f|_{\Omega}(y))^2 A_1(x,y,\omega_1,\omega_2,\lambda,\sigma) \mathbb{1}_{\{|\sigma| \le -2\lambda\}} dx dy d\omega_1 d\omega_2 d\lambda d\sigma \qquad (4.15)$$

where the amplitude $A_1 \in S^1(\mathbb{R}^{4n+2})$ is also supported in $\{t \in [-\underline{\tau},\underline{\tau}], (x,y) \in \Omega'^2, |x-y| \leq \underline{\ell}, |\lambda| \leq \underline{\ell}\}.$

• Step 2. We study the phase Ψ_1 . This phase vanishes along the diagonal $\{x=y\}$ and it can be factorised into

$$\Psi_1(x, y, \omega_1, \omega_2, \lambda, \sigma) = (x - y) \cdot \zeta(x, y, \omega_1, \omega_1, \lambda, \sigma)$$

where $\zeta \in \mathbb{R}^{4n+2} \to \mathbb{R}^n$ is smooth and approximated by $\zeta \simeq \sqrt{\lambda - V(x)}(\omega_1 - \omega_2)$ for x - y small. This allows us to show that for $\omega_1 \in \mathbb{S}^{n-1}$, $x \in \Omega'$, $|x - y| \leq \underline{\ell}$ fixed, the map $(\omega_2, \sigma) \mapsto \zeta$ is a (smooth) diffeomorphism from a neighborhood of $(\omega_1, 0)$ in $S^{n-1} \times \mathbb{R}$ onto $\{\zeta \in \mathbb{R}^n : |\zeta| \leq \delta\}$, where δ can be chosen much larger than $\underline{\ell}$ and much smaller than \underline{c} .

We will justify in step 4 that the main contribution to the integral (4.15) comes from this region. Hence, by a change of coordinates, the main term in $\|[\widetilde{\Pi}, f|_{\Omega}]\|_{\mathbb{I}^2}^2$ is

$$\frac{-1}{(2\pi\hbar)^{2n}} \int e^{i\frac{(x-y)\cdot\zeta}{\hbar}} (f|_{\Omega}(x) - f|_{\Omega}(y))^2 A_2(x,y,\omega,\zeta,\lambda) \mathbb{1}_{\{|\sigma(x,y,\zeta,\omega,\lambda)| > -2\lambda > 0\}} dx dy d\zeta d\omega d\lambda$$

where $A_2 \in S^1(\mathbb{R}^{4n+1})$, $\sigma : \mathbb{R}^{4n+1} \to \mathbb{R}$ is smooth and $\sigma \simeq 2\sqrt{\lambda - V(x)} \zeta \cdot \omega$.

• Step 3. At this stage, we argue that symbols of the type

$$(x, y, \xi) \mapsto \int_{\mathbb{S}^{n-1} \times \mathbb{R}} A_2(x, y, \omega, \xi, \lambda) \mathbb{1}_{\{|\sigma(x, y, \xi, \omega, \lambda)| > -2\lambda > 0\}} d\omega d\lambda$$

are in the class \mathcal{F} and we prove (4.13).

• Step 4. It remains to show that the remaining part of the integral (4.15),

$$\frac{1}{(2\pi\hbar)^{2n}} \int e^{i\frac{(x-y)\cdot\zeta(x,y,\omega_1,\omega_1,\lambda,\sigma)}{\hbar}} (1-\chi_{\delta}) (\zeta(x,y,\omega_1,\omega_1,\lambda,\sigma)) (f|_{\Omega}(x)-f|_{\Omega}(y))^2$$
(4.16)

 $A_2(x, y, \omega_1, \omega_2, \lambda, \sigma) \mathbb{1}_{\{|\sigma| \le -2\lambda\}} dx dy d\omega_1 d\omega_2 d\lambda d\sigma,$

is $\mathcal{O}(\hbar^{1-n})$. Here, as usual $\chi_{\delta}(\cdot) = \chi(\delta \cdot)$, and the function $\chi : \mathbb{R}^n \to [0,1]$ is smooth, equal to 1 on $B(0,\frac{\delta}{2})$ and to 0 outside $B(0,\delta)$.

If $|x-y| \leq \underline{\ell}$ with $\underline{\ell}$ small with respect to δ , then the oscillating phase (4.16) has no stationary point. Thus we expect (4.16) to be rather small. The issue is that the amplitude is not smooth with respect to (x,y). Nevertheless, we can integrate by parts twice, once with respect to x and once with respect to y, and this reduces (4.16) (up to an error $\mathcal{O}(\hbar^{1-n})$) to an integral over the (smooth) boundary of Ω . It remains to bound an integral of the type

$$\frac{1}{(2\pi\hbar)^{2(n-1)}} \int_{\{\hat{x},\hat{y}\in\partial\Omega\}\times\{\omega_1,\omega_2\in\partial B\}} e^{i\frac{(\hat{x}-\hat{y})\cdot\zeta(\hat{x},\hat{y},\omega_1,\omega_1,\lambda,\sigma)}{\hbar}} \frac{1}{\chi_{\delta}} (\omega-\omega_2)) A_3(\hat{x},\hat{y},\omega_1,\omega_2,\lambda,\sigma) d\hat{x} d\hat{y} d\omega_1 d\omega_2$$

where $A_3 \in S^1(\mathbb{R}^{4n+2})$. To complete the proof, we argue that for a $\hat{x} \in \partial\Omega$, the phase has non-singular Hessian in one of the pair (\hat{y}, ω_1) or (\hat{y}, ω_2) . Hence, by a stationary phase argument, this integral is at most of order $\mathcal{O}(\hbar^{1-n})$. This completes the proof.

4.2.1 Step 1: Stationary phase

We make the change of coordinates (4.14) (the Jacobian of this change of variable is $(r_1r_2)^{n-1}$ for $n \ge 1$) and study the critical point of the phase Ψ_0 in the variables (t_1, r_1, t_2, r_2) . According to (3.5) and (2.2), we verify that

$$\partial_{t_1} \Psi_1(x, y, \xi_1, \xi_2, t_1, t_2, \lambda_1, \lambda_2) = \partial_{\xi} \varphi(t_1, x, \xi_1) - \lambda_1 = V(x) + r_1^2 - \lambda_1 + \mathcal{O}(t_1)
\partial_{r_1} \Psi_1(x, y, \xi_1, \xi_2, t_1, t_2, \lambda_1, \lambda_2) = \omega_1 \cdot (\partial_{\xi} \varphi(t_1, x, \xi_1) - y) = \omega_1 \cdot ((x - y) + \mathcal{O}(t_1^2)) + 2t_1 r_1$$
(4.17)

and similarly for $\partial_{t_2}\Psi_1, \partial_{r_2}\Psi_1$ switching $1 \leftarrow 2$ and the sign.

Let us first study the first line of (4.17). Since $|t_1| \leq \underline{\tau} \ll \underline{c} \leq \lambda - \sigma/2 - V(x)$, for every t_1 there exists a unique r_1^c solving the first equation, it is a smooth function of all other parameters, and $r_1^c|_{t_1=0} = \sqrt{\lambda + \sigma/2 - V(x)}$. In particular, $r_1^c \geq \underline{c}$. Now we turn to the second line of (4.17): with $r_1 = r_1^c$ bounded away from below, for every $|x-y| \leq \underline{\ell} \ll \underline{c}$ there exists at most one solution for t_1 such that $|t_1| \leq \underline{\tau}$; moreover at y = x this solution exists and is equal to 0.

All in all, for x - y small there exists a unique stationary point given by

$$r_{1}^{c} = \sqrt{\lambda + \sigma/2 - V(x)} + O(|x - y|)$$

$$t_{1}^{c} = -\frac{\omega_{1} \cdot (x - y)}{2\sqrt{\lambda + \sigma/2 - V(x)}} + O(|x - y|^{2})$$
(4.18)

and similarly for (r_2^c, t_2^c) replacing $(\lambda + \sigma) \leftarrow (\lambda - \sigma)$ and $\omega_1 \leftarrow \omega_2$. Differentiating (4.17) again, we obtain that the Hessian of Ψ_1 with respect to (t_1, r_1, t_2, r_2) is of the form

$$2 \begin{pmatrix} \star & r_1 + \mathcal{O}(t_1) & 0 & 0 \\ r_1 + \mathcal{O}(t_1) & t_1 + \mathcal{O}(t_1^2) & 0 & 0 \\ 0 & 0 & \star & -r_2 + \mathcal{O}(t_2) \\ 0 & 0 & r_2 + \mathcal{O}(t_2) & t_2 + \mathcal{O}(t_2^2) \end{pmatrix}.$$

For $r_1, r_2 \ge \underline{c}/4$ and $|t_1|, |t_2| \le \underline{\tau}$, this Hessian is non-degenerate and its determinant, evaluated at the critical point (4.18), is given by

$$J(x, y, \omega_1, \omega_2, \sigma, \lambda) = 16((\lambda - V(x))^2 - \sigma^2/4) + \mathcal{O}(|x - y|).$$
(4.19)

We introduce a new phase

$$\Psi_1: (x,y,\omega_1,\omega_2,\lambda,\sigma) \mapsto \Psi_0(x,y,t_1^c,t_2^c,r_1^c\omega_1,r_2^c\omega_2,\lambda+\frac{\sigma}{2},\lambda-\frac{\sigma}{2})$$

and we denote

$$R = R(x,\lambda) := \sqrt{\lambda - V(x)}.$$
(4.20)

By (4.18), if x = y and $\sigma = 0$ one has $R = r_i^c$ for $j \in \{1, 2\}$ as well as $J = (2R)^4$.

We are in position to apply Lemma A.1 to the integral (4.1), with the variables $(x, y, \omega_1, \omega_2, \lambda, \sigma)$ as parameters. There is an amplitude $A_1 \in S^1(\mathbb{R}^{4n+2})$ so that

$$\|[\widetilde{\Pi}, f|_{\Omega}]\|_{J^{2}}^{2} = \frac{1}{(2\pi\hbar)^{2n}} \int e^{i\frac{\Psi_{1}(x, y, \omega_{1}, \omega_{2}, \lambda, \sigma)}{\hbar}} (f|_{\Omega}(x) - f|_{\Omega}(y))^{2} A_{1}(x, y, \omega_{1}, \omega_{2}, \lambda, \sigma) \mathbb{1}_{\{|\sigma| \leq -2\lambda\}} dx dy d\omega_{1} d\omega_{2} d\sigma d\lambda.$$
(4.21)

Moreover, the principal symbol of A_1 is given as follows, in terms of (4.18) and (4.19),

$$A_1(x, y, \omega_1, \omega_2, \lambda, \sigma)|_{\hbar=0} = \frac{(r_1^c r_2^c)^{n-1}}{\sqrt{J}} A_0(x, y, r_1^c \omega_1, r_2^c \omega_2, t_1^c, t_2^c, \lambda + \frac{\sigma}{2}, \lambda - \frac{\sigma}{2}).$$

The factor $(r_1^c r_2^c)^{n-1}$ comes from the Jacobian of the change of coordinates. In particular, according to (4.2), at $(y, \sigma) = (x, 0)$,

$$A_{1}(x, x, \omega_{1}, \omega_{2}, \lambda, 0) = \frac{R^{2n-2}A_{0}(x, x, R\omega_{1}, R\omega_{2}, 0, 0, \lambda, \lambda)}{\sqrt{R^{4}}} = R^{2n-4}\vartheta(V(x) + R^{2})^{2}\chi(\lambda)^{2}/4$$
$$= R^{2(n-2)}g(\lambda)/4, \qquad x \in \Omega', \lambda \in \mathbb{R}, \omega_{1}, \omega_{2} \in \mathbb{S}^{n-1}, \tag{4.22}$$

where R is given by (4.20) and $g := \vartheta^2 \chi^2$ satisfies g(0) = 0. Moreover, A_1 is supported in

$$\{(x,y) \in \Omega' \times \Omega', |x-y| \le \underline{\ell}, |\lambda| \le \underline{\ell}\}.$$

4.2.2 Step 2: Study of the phase Ψ_1

By (4.18), $t_1^c = t_2^c = 0$ if x = y, then with $\xi_j = r_j^c \omega_j$, we have on the diagonal

$$\Psi_1(x, x, \omega_1, \omega_2, \lambda, \sigma) = \varphi(0, x, \xi_1) - \varphi(0, x, \xi_2) - (\xi_1 - \xi_2) \cdot x = 0.$$

Thus there exists a smooth function $\zeta: \mathbb{R}^{2n+2} \to \mathbb{R}^n$ such that we can write

$$\Psi_1(x, y, \omega_1, \omega_2, \lambda, \sigma) = (x - y) \cdot \zeta(x, y, \omega_1, \omega_2, \lambda, \sigma). \tag{4.23}$$

Moreover, the original phase Ψ_1 is anti-symmetric with respect to the exchange $\xi_1 \leftrightarrow \xi_2, t_1 \leftrightarrow t_2, \lambda_1 \leftrightarrow \lambda_2$. From the critical point equation, this implies that $(r_1^c, t_1^c) = (r_2^c, t_2^c) = (r^c, t^c)$ if $(\omega_1, \sigma) = (\omega_2, 0)$ so that

$$\Psi_1(x, y, \omega, \omega, \lambda, 0) = \Psi_0(x, y, t^c, t^c, r^c\omega, r^c\omega, \lambda, \lambda) = 0$$

Hence

$$\zeta(x, y, \omega, \omega, \lambda, 0) = 0,$$
 on $\{(x, y) \in \Omega' \times \Omega', |x - y| \le \underline{\ell}, |\lambda| \le \underline{\ell}\}.$

In particular, with $\mathcal{Z}(x,\omega_1,\omega_2,\sigma,\lambda) = \zeta(x,x,\omega_1,\omega_2,\sigma,\lambda)$,

$$\zeta(x, y, \omega_1, \omega_2, \sigma, \lambda) = \mathcal{Z}(x, \omega_1, \omega_2, \sigma, \lambda) + \mathcal{O}\left(\sqrt{|\omega_1 - \omega_2|^2 + |\sigma|^2}|x - y|\right). \tag{4.24}$$

Recall that $|x-y| < \underline{\ell}$ in the integral in questions, so we first study \mathcal{Z} instead of ζ . Note that

$$\mathcal{Z}(x,\omega_1,\omega_2,\sigma,\lambda) = -\partial_y \Psi_1\big|_{y=x} = -\partial_y \Psi_0\big|_{y=x,t_1=t_2=0,r_1=r_1^c,r_2=r_2^c}.$$
(4.25)

Here we used the property of the critical point for which $\partial_{t_j}\Psi_1|_{t_1^c,r_1^c,t_2^c,r_2^c}=\partial_{r_j}\Psi_1|_{t_1^c,r_1^c,t_2^c,r_2^c}=0$ for $j\in\{1,2\}$ and $t_1^c=t_2^c=0$ if x=y. Since $\partial_y\Psi_1=\xi_1-\xi_2$, by (4.18), this yields

$$\mathcal{Z}(x,\omega_1,\omega_2,\sigma,\lambda) = r_1^c \omega_1 - r_2^c \omega_2 = \omega_1 \sqrt{\lambda - V(x) + \frac{\sigma}{2}} - \omega_2 \sqrt{\lambda - V(x) - \frac{\sigma}{2}}.$$
 (4.26)

Linearizing this function for small σ (here, $|\sigma| \leq 2\underline{\ell}$), we obtain

$$\mathcal{Z}(x,\omega_1,\omega_2,\sigma,\lambda) = (\omega_1 - \omega_2)(R + \mathcal{O}(\sigma^2)) + (\omega_1 + \omega_2)\frac{\sigma}{4R}(1 + \mathcal{O}(\sigma^2))$$

with $R \ge \underline{\mathbf{c}}$ as in (4.20).

We now consider the equation $\mathcal{Z}(x,\omega_1,\omega_2,\sigma,\lambda)=\xi$ for fixed $(x,\omega_1,\lambda)\in\Omega'\times\mathbb{S}^{n-1}\times[-\underline{\ell},\underline{\ell}]$ and $\xi\in\mathbb{R}^n$. We have

$$\omega_2(R - \frac{\sigma}{4R}) = \omega_1(R + \frac{\sigma}{4R}) - \xi + \mathcal{O}(\sigma^2).$$

To solve this equation, we decompose $\omega_2 = \alpha \omega_1 + \nu$ where $\nu \in \omega_1^{\perp}$ and $\alpha = \sqrt{1 - |\nu|^2} \ge 0$, we obtain

$$\begin{cases} \alpha = 1 + \frac{\sigma}{2R^2} - \frac{\xi_1}{R - \frac{\sigma}{4R}} + \mathcal{O}(\sigma^2), & \text{where } \xi_1 = \xi \cdot \omega_1 \\ \nu = -\frac{\xi^{\perp}}{R - \frac{\sigma}{4R}} + \mathcal{O}(\sigma^2), & \text{where } \xi = \xi_1 + \xi^{\perp}. \end{cases}$$

In particular, we have

$$1 = \alpha^2 + |\nu|^2 = 1 + \frac{|\xi|^2}{(R - \sigma/4R)^2} + \frac{\sigma}{R^2} - \frac{2\xi_1}{R - \sigma/4R} + \mathcal{O}(\sigma^2).$$

Since $\delta \ll \underline{c} \leq R$, the last equation determines σ for $|\xi| < \delta$, and

$$\sigma = 2R\xi_1 + \mathcal{O}(|\xi|^2)$$
 and then $\omega_2 = \omega_1 - \xi^{\perp}/R + \mathcal{O}(|\xi|^2)$.

To summarize, choosing a small parameter δ with $\underline{\ell} \ll \delta \ll \underline{c}$, given $(x, \omega_1, \lambda) \in \Omega' \times \mathbb{S}^{n-1} \times \mathbb{R}$ and $\xi \in \delta B_n$, the equation $\mathcal{Z}(x, \omega_1, \omega_2, \sigma, \lambda) = \xi$ has a unique smooth solution $(\sigma, \omega_2) \in \mathbb{R} \times \mathbb{S}^{n-1}$. Moreover, by (4.26), we also compute

$$\partial_{\sigma} \mathcal{Z} = \frac{\omega_1 + \omega_2}{4R} + \mathcal{O}(\sigma(\omega_1 - \omega_2)), \qquad \partial_{\omega_2} \mathcal{Z} = RI_n + \mathcal{O}(\sigma),$$

where I_n denotes the identity matrix. These derivatives are non-degenerate in a neighbourhood of the previous solution, so according to (4.24), the equation $\zeta(x,y,\omega_1,\omega_2,\sigma,\lambda)=\xi$, given $(x,\omega_1,\lambda)\in\Omega'\times\mathbb{S}^{n-1}\times\mathbb{R}$ and $(y,\xi)\in\mathbb{R}^{2n}$ with $|x-y|<\underline{\ell}$, $|\xi|<\delta$, also has a unique smooth solution $(\sigma,\omega_2)\in\mathbb{R}\times\mathbb{S}^{n-1}$ with the expansions:

$$\begin{cases} \sigma = 2R\xi_1 + \mathcal{O}(|\xi|(|\xi| + |x - y|)) \\ \omega_2 = \omega_1 - \xi^{\perp}/R + \mathcal{O}(|\xi|(|\xi| + |x - y|)) \end{cases}, \qquad \xi = (\xi_1, \xi^{\perp}), \ \xi_1 = \xi \cdot \omega_1, \ R = \sqrt{\lambda - V(x)}. \tag{4.27}$$

Hence, we can make a change of variable $\xi \in \delta B_n \mapsto (\sigma, \omega_2)$, after which the phase becomes

$$\Psi_1(x, y, \omega_1, \omega_2, \sigma, \lambda) = (x - y) \cdot \xi.$$

To use this, we must split the integral (4.21) in two parts by introducing a cutoff in \mathcal{Z} (equivalently in ξ after the change of coordinates):

$$\|[\widetilde{\Pi}, f|_{\Omega}]\|_{\mathbf{J}^2}^2 = I_{1,\hbar} + I_{2,\hbar}$$

where

$$I_{1,\hbar} = \frac{1}{(2\pi\hbar)^{2n}} \int e^{i\frac{\Psi_1(x,y,\omega_1,\omega_2,\sigma,\lambda)}{\hbar}} (f|_{\Omega}(x) - f|_{\Omega}(y))^2 A_1'(x,y,\omega_1,\omega_2,\sigma,\lambda) \mathbb{1}_{\{|\sigma| \le -2\lambda\}} dx dy d\omega_1 d\omega_2 d\sigma d\lambda,$$

$$I_{2,\hbar} = \frac{1}{(2\pi\hbar)^{2n}} \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} (f|_{\Omega}(x) - f|_{\Omega}(y))^2 A_2(x,y,\omega_1,\xi,\lambda) \mathbb{1}_{\{|\sigma(x,y,\xi,\omega_1,\lambda)| \le -2\lambda\}} dx dy d\omega_1 d\xi d\lambda.$$

$$(4.28)$$

We tune a smooth cutoff so that

• the amplitude A_1' belongs to $S^1(\mathbb{R}^{4n+2})$ with $A_1'(x,y,\omega_1,\omega_2,\sigma,\lambda)=0$ on the set $\{|\mathcal{Z}(x,\omega_1,\omega_2,\sigma,\lambda)|\leq \delta\}$ and A_1' is supported on the set $(x,\lambda)\in\Omega'\times[-\underline{\ell},\underline{\ell}]$ and $|x-y|\leq\underline{\ell}$ with $\underline{\ell}\ll\delta$.

• the amplitude $A_2 \in S^1(\mathbb{R}^{4n+1})$ and it is supported in

$$(x,\lambda) \in \Omega' \times \mathbb{R}, \ (y,\xi) \in \mathbb{R}^{2n}, \ |y-x|+|\lambda|+|\xi| \le 2\delta \ll \underline{c}.$$

Finally, the principal part of A_2 satisfies on the diagonal x = y and at $\xi = 0$ (where $\mathcal{Z} = 0$ also)

$$A_2(x, x, 0, \omega_1, \lambda)|_{\hbar=0} = A_1(x, x, \omega_1, \omega_1, \lambda, 0) \left(\frac{\mathrm{d}\sigma \mathrm{d}\omega_2}{\mathrm{d}\xi_1 \mathrm{d}\xi^{\perp}} \right) \Big|_{y=x, \xi=0}$$

since $(\sigma, \omega_2) = (0, \omega_1)$ when $(y, \xi) = (x, 0)$, cf. (4.27). Moreover, we compute the Jacobian

$$\frac{\mathrm{d}\sigma\mathrm{d}\omega_2}{\mathrm{d}\xi_1\mathrm{d}\xi^{\perp}}\Big|_{y=x,\xi=0} = \left| \det \begin{pmatrix} 2R & \dots & 0 & \dots \\ \vdots & \ddots & & \\ 0 & & I_{n-1}/R & \\ \vdots & & \ddots & \end{pmatrix} \right| = 2R^{2-n}.$$

Hence, by (4.22) with $R(x,0) = |V(x)|^{1/2}$, we obtain

$$A_2(x, x, 0, \omega, 0) = R(x, 0)^{n-2} g(0)/2 = |V(x)|^{n/2-1}/2, \qquad x \in \Omega', \omega \in \mathbb{S}^{n-1}.$$
(4.29)

4.2.3 Step 3: the main symbol is in the class \mathcal{F}

We now prove the following fact.

Proposition 4.10. The symbol

$$F: (x, y, \xi) \in \mathbb{R}^{3n} \times (0, 1] \mapsto \int_{\mathbb{S}^{n-1} \times \mathbb{R}} A_2(x, y, \xi, \omega, \lambda) \mathbb{1}_{\{|\sigma(x, y, \xi, \omega, \lambda)| > -2\lambda > 0\}} d\omega d\lambda.$$

is in the class $\mathcal{F}(\mathbb{R}^{2n}, \mathbb{R}^n)$ (cf. Definition 4.4) and it satisfies (4.13).

According to (4.28) and the previous definition, we can rewrite

$$I_{2,\hbar} = \frac{-1}{(2\pi\hbar)^{2n}} \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} (f|_{\Omega}(x) - f|_{\Omega}(y))^2 F(x,y,\xi) dxdyd\xi + ||[f|_{\Omega},K]||^2$$

where

$$K: (x,y) \mapsto \frac{1}{(2\pi\hbar)^n} \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} a(x,y,\xi) \mathrm{d}x \mathrm{d}y \mathrm{d}\xi \quad \text{and} \quad a: \mathbb{R}^{3n} \mapsto \int A_2(x,y,\xi,\omega,\lambda) \mathbb{1}_{\{\lambda < 0\}} \mathrm{d}\lambda \mathrm{d}\omega.$$

The amplitude a belongs to $S^1(\mathbb{R}^{3n})$ ($\|a\|_{\mathscr{C}^k}$ are bounded uniformly in \hbar by differentiating under the integral). Consequently, by Lemma 4.8, $\|[f|_{\Omega}, K]\|^2 = \mathcal{O}(\hbar^{1-n})$

We will show in the next section (Proposition 4.11) that the integral $I_{1,\hbar} = \mathcal{O}(\hbar^{1-n})$ as $\hbar \to 0$. This allows to conclude that

$$\|[\widetilde{\Pi}, f|_{\Omega}]\|_{J^2}^2 = \frac{1}{(2\pi\hbar)^{2n}} \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} (f|_{\Omega}(x) - f|_{\Omega}(y))^2 F(x, y, \xi) dx dy d\xi + \mathcal{O}(\hbar^{1-n})$$

which completes the proof of Proposition 4.9.

Proof of Proposition 4.10.

First observe that according to (4.27), there exist two smooth functions $\nu:(x,y,\xi,\omega,\lambda)\in\mathbb{R}^{4n+1}\mapsto\mathbb{S}^{n-1}$ and $r:(x,y,\xi,\omega,\lambda)\in\mathbb{R}^{4n+1}\mapsto\mathbb{R}_+$ such that

$$\sigma = 2r\nu \cdot \xi, \qquad (r,\nu) = (R,\omega) + \mathcal{O}(|x-y| + |\xi|). \tag{4.30}$$

Then, we claim that for $\{(x,y,\xi)\in\mathbb{R}^{3n}:x\in\Omega',|x-y|,|\xi|<2\delta\}$, we can make a (smooth) change of coordinates

$$(\lambda, \omega) \in \mathbb{R} \times \mathbb{S}^{n-1} \mapsto (\varsigma, \nu) \in \mathbb{R} \times \mathbb{S}^{n-1}, \qquad \varsigma = -\lambda/r$$

in the integral F. We denote by $J:(x,y,\xi,\nu,\varsigma)\in\mathbb{R}^{4n+1}\to\mathbb{R}_+$ the corresponding Jacobian. Observe that if $(y,\xi)=(x,0),\ \varsigma=-\lambda/\sqrt{\lambda-V(x)}$ and, by solving this quadratic equation and choosing the appropriate root, we obtain

$$\lambda = -\varsigma \sqrt{-V(x)} + \mathcal{O}(\varsigma^2), \qquad \omega = \nu.$$

Thus, in this case.

$$J(x, x, 0, \nu, \varsigma) = \sqrt{-V(x)} + \mathcal{O}(\varsigma). \tag{4.31}$$

Generally, $|\varsigma| \leq \underline{c}$ (as $r \geq \underline{c}$ for $x \in \Omega'$ and $|y - x|, |\lambda|, |\xi| \leq 2\delta$ with $\delta \ll \underline{c}$), so a similar computation using (4.30) shows that $J(x, y, \xi, \nu, \varsigma)$ is non-degenerate. Hence, this change of coordinates is admissible and there is an amplitude $B \in S^1(\mathbb{R}^{4n+1})$ so that

$$F = \int_{\mathbb{S}^{n-1} \times \mathbb{R}} B(x, y, \xi, \nu, \varsigma) \mathbb{1}_{\{|\nu \cdot \xi| > \varsigma > 0\}} d\nu d\varsigma.$$

$$(4.32)$$

Moreover, by (4.30), the principal part of B is given by, if $(y, \xi, \zeta) = (x, 0, 0)$ with $x \in \Omega'$, $\nu \in \mathbb{S}^{n-1}$,

$$B(x, x, 0, \nu, 0)|_{\hbar=0} = A_2(x, x, 0, \nu, 0)|_{\hbar=0} J(x, x, 0, \nu, 0)$$
$$= |V(x)|^{\frac{n-1}{2}}/2$$
(4.33)

using (4.29) and (4.31).

Using spherical coordinates, writing $\xi = R\theta$ with $(R, \theta) \in \mathbb{R}_+ \times \mathbb{S}^{n-1}$ and making a change of variable $\varsigma \leftarrow R\varsigma$ in (4.32), we obtain for $(x, y, \xi) \in \mathbb{R}^{3n}$

$$F(x,y,\xi) = R \int_{\mathbb{S}^{n-1} \times \mathbb{R}} B(x,y,\xi,\nu,R\varsigma) \mathbb{1}_{\{|\nu \cdot \vartheta| > \varsigma > 0\}} d\nu d\varsigma.$$
 (4.34)

Then, since B is smooth, by (4.33), the principal part of F satisfies on the diagonal $\{x = y\}$,

$$F(x, x, \xi) = R \int_{\mathbb{S}^{n-1} \times \mathbb{R}} (B(x, x, 0, \nu, 0) + \mathcal{O}(R)) \mathbb{1}_{\{|\nu \cdot \vartheta| > \varsigma > 0\}} d\nu d\varsigma$$

$$= \frac{|\xi| |V(x)|^{\frac{n-1}{2}}}{2} \left\{ \int_{\mathbb{S}^{n-1}} |\nu_1| d\nu + \mathcal{O}(|\xi|) \right\}$$

$$= c_{n-1} |\xi| |V(x)|^{\frac{n-1}{2}} + \mathcal{O}(|\xi|^2)$$

uniformly for $x \in \Omega'$. Here, we used that for $n \geq 2$,

$$\frac{1}{2} \int_{\mathbb{S}^{n-1}} |\nu_1| d\nu = \frac{|\mathbb{S}^{n-2}|}{n-1} = |B_{n-1}|$$

given the relationship between the volume of the unit sphere $\mathbb{S}^{n-1} = \partial B_n$ and the unit ball $B_n \subset \mathbb{R}^n$ (in particular, $|\mathbb{S}^0| = |B_1| = 2$). In dimension n = 1, the situation is special as $\nu \in \{\pm 1\}$, so that $c_0 = 1$. This establishes (4.13).

For $\theta \in \mathbb{S}^{n-1}$, let $\mathcal{R}_{\theta} : \mathbb{S}^{n-1} \to \mathbb{S}^{n-1}$ be the rotation so that $\mathcal{R}_{\theta}^{*}(\theta) = e_{1}$ (the first coordinate vector in \mathbb{R}^{n}). By (4.34), using the invariance under rotation of the Haar measure $d\nu$ on the sphere \mathbb{S}^{n-1} , we obtain

$$F(x,y,\xi) = R \int_{\mathbb{S}^{n-1} \times \mathbb{R}} B(x,y,\xi,\mathcal{R}_{\theta}\nu,R\varsigma) \mathbb{1}_{\{|\nu_1|>\varsigma>0\}} d\nu d\varsigma, \qquad \xi = R\theta, (R,\theta) \in \mathbb{R}_+ \times \mathbb{S}^{n-1}.$$

$$(4.35)$$

This function is clearly smooth in $(x, y, \xi) \in \mathbb{R}^{2n} \times \{\mathbb{R}^n \setminus 0\}$. In fact, for any $k \in \mathbb{N}_0$, as $R(\xi) \to 0$,

$$\|\partial_{\xi}^k \mathcal{R}_{\theta(\xi)}\| = \mathcal{O}_k(R^{-k}), \qquad \|\partial_{\xi}^k R(\xi)\| = \mathcal{O}_k(R^{1-k}).$$

Indeed, $\theta \in \mathbb{S}^{n-1} \to \mathcal{R}_{\theta}$ is smooth, with $\partial_{\xi} R(\xi) = \theta(\xi)$ for $\xi \neq 0$ and $\|\partial_{\xi}^{k} \theta(\xi)\| = \mathcal{O}_{k}(R^{-k})$. Thus, by differentiating (4.35) under the integral. we conclude that for any $k \in \mathbb{N}_{0}$, $\|\partial_{\xi}^{k} F(x, y, \xi)\| = \mathcal{O}_{k}(R^{1-k})$. as $R(\xi) \to 0$. This completes the proof that $F \in \mathcal{F}(\mathbb{R}^{2n}, \mathbb{R}^{n})$.

4.2.4 Step 4: Control of $I_{1,\hbar}$

At this stage, it remains to show that $I_{1,\hbar} = \mathcal{O}(\hbar^{1-n})$, where we recall that $I_{1,\hbar}$ is given by (4.28). By (4.28), we can work with (σ, λ) fixed and it suffices to obtain the following bounds (We drop the dependency of Ψ_1, \mathcal{Z} , etc on (σ, λ) for notational convenience. The proof of Proposition 4.11 relies on the stationary phase as formulated in Proposition A.1 which allows to deal with the parameters (σ, λ) in a uniform way.).

Proposition 4.11. Let (Ψ_1, \mathcal{Z}) be as in Section 4.2.2 and assume that $A \in S^1(\mathbb{R}^{4n})$ is supported on the set $\{x \in \Omega', |x-y| \leq \underline{\ell}, |\mathcal{Z}(x,\omega_1,\omega_2)| > \delta\}$ with $\underline{\ell} \ll \delta \ll 1$. Then, as $\hbar \to 0$,

$$\frac{1}{(2\pi\hbar)^{2n}} \int_{\omega_1,\omega_2 \in \mathbb{S}^{n-1}} e^{i\frac{\Psi_1(x,y,\omega_1,\omega_2)}{\hbar}} (f|_{\Omega}(x) - f|_{\Omega}(y))^2 A(x,y,\omega_1,\omega_2) dx dy d\omega_1 d\omega_2 = \mathcal{O}(\hbar^{1-n}).$$

Proof. Expanding the square, we need to deal with two integrals of the type

$$\frac{1}{(2\pi\hbar)^{2n}} \int e^{i\frac{\Psi_1(x,y,\omega_1,\omega_2)}{\hbar}} A_1(x,y,\omega_1,\omega_2) \mathbb{1}_{\Omega}(x) \mathrm{d}x \mathrm{d}y \mathrm{d}\omega_1 \mathrm{d}\omega_2 \tag{4.36}$$

$$\frac{1}{(2\pi\hbar)^{2n}} \int e^{i\frac{\Psi_1(x,y,\omega_1,\omega_2)}{\hbar}} A_2(x,y,\omega_1,\omega_2) \mathbb{1}_{\Omega}(x) \mathbb{1}_{\Omega}(y) dx dy d\omega_1 d\omega_2$$
(4.37)

where the amplitudes A_1, A_2 belong to $S^1(\mathbb{R}^{4n})$. There is a minus sign, but we do not expect a cancelation in this regime.

According to (4.25),

$$\partial_{y}\Psi_{1}(x,y,\omega_{1},\omega_{2}) = -\mathcal{Z}(x,\omega_{1},\omega_{2}) + \mathcal{O}(x-y). \tag{4.38}$$

Hence, if $A \in S^1(\mathbb{R}^{4n})$ is supported in $\{|x-y| \leq \underline{\ell}, |\mathcal{Z}(x,\omega_1,\omega_2)| > \delta\}$ with $\underline{\ell} \ll \delta \ll \underline{c}$, $\partial_y \Psi_1$ does not vanish on $\operatorname{supp}(A)$ and we can write

$$A(x, y, \omega_1, \omega_2) = \frac{1}{2\pi} J(x, y, \omega_1, \omega_2) \cdot \partial_y \Psi_1(x, y, \omega_1, \omega_2),$$

where $J: \mathbb{R}^{4n} \to \mathbb{R}^n$ is also in S^1 with the same support as A.

In these circumstances, we can repeat this procedure and integrate by parts with respect to y arbitrarily often in (4.36), and obtain

$$(4.36) = \mathcal{O}(\hbar^{\infty}).$$

For (4.37), we integrate by parts once and find that there are $B_1, B_2 \in S^1(\mathbb{R}^{4n})$ so that

$$(4.37) = \frac{1}{(2\pi\hbar)^{2n-1}} \int_{\{\hat{y}\in\partial\Omega\}} e^{i\frac{\Psi_1(x,\hat{y},\omega_1,\omega_2)}{\hbar}} B_1(x,\hat{y},\omega_1,\omega_2) \mathbb{1}_{\Omega}(x) dx d\hat{y} d\omega_1 d\omega_2$$

$$(4.39)$$

$$+\frac{1}{(2\pi\hbar)^{2n-1}}\int e^{i\frac{\Psi_1(x,y,\omega_1,\omega_2)}{\hbar}}B_2(x,y,\omega_1,\omega_2)\mathbb{1}_{\Omega}(x)\mathbb{1}_{\Omega}(y)\mathrm{d}x\mathrm{d}y\mathrm{d}\omega_1\mathrm{d}\omega_2. \tag{4.40}$$

Here, $B_1(\cdot) = \nu_{\Omega}(\hat{y}) \cdot J(\cdot) A_2(\cdot)$ where ν_{Ω} is the exterior normal to $\partial \Omega$, and we integrate the variable \hat{y} with respect to the volume measure over $\partial \Omega$ – it is worth remembering here that Ω has a smooth boundary.

The integral (4.40) has the same form as the original integral (4.37) with an extra power of \hbar . Thus, if we show that (4.39) = $\mathcal{O}(\hbar^{-n+1})$, by induction, we can conclude that also (4.37) = $\mathcal{O}(\hbar^{-n+1})$.

Then we consider the integral (4.39). Similarly to (4.25), the phase $\partial_x \Psi_1$ does not vanish on supp (B_1) . Thus, integrating by part with respect to x, we obtain

$$(4.39) = \frac{1}{(2\pi\hbar)^{2(n-1)}} \int_{\{\hat{x}, \hat{y} \in \partial\Omega\}} e^{i\frac{\Psi_1(\hat{x}, \hat{y}, \omega_1, \omega_2)}{\hbar}} B_3(\hat{x}, \hat{y}, \omega_1, \omega_2) d\hat{x} d\hat{y} d\omega_1 d\omega_2 + \frac{Q_{\hbar}}{2\pi\hbar}$$
(4.41)

where $B_3 \in S^1(\mathbb{R}^{4n})$ and Q_{\hbar} is an integral of the same type as (4.39). Thus, by the same reasoning as above, it suffices to show that the boundary integral in (4.41) is $\mathcal{O}(\hbar^{1-n})$ to conclude that the original integral is also of order $\mathcal{O}(\hbar^{1-n})$.

In dimension n = 1, obviously (4.41) = $\mathcal{O}(1)$, so this reasoning already shows that the integral in question is bounded, as required.

In what follows, we assume that $n \geq 2$ and focus on the boundary integral (4.41) for a fixed $\hat{x} \in \partial\Omega$. Recall that the amplitude B_3 belongs to S^1 and it is supported on $\{\hat{x}, \hat{y} \in \partial\Omega : |\hat{x} - \hat{y}| \leq \underline{\ell}, |\mathcal{Z}(\hat{x}, \omega_1, \omega_2)| > \delta\}$. Let $\Pi_{\hat{x}}$ denote the orthogonal projection on $T_{\hat{x}}(\partial\Omega)$. On the above set $(\underline{\ell})$ is small,

$$(\hat{x} - \hat{y}) = \Pi_{\hat{x}}(\hat{x} - \hat{y}) + \mathcal{O}(|\hat{x} - \hat{y}|^2)$$

then, by (4.23), the phase satisfies

$$\Psi_{1}(\hat{x}, \hat{y}, \omega_{1}, \omega_{2}) = (\hat{x} - \hat{y}) \cdot \mathcal{Z}(\hat{x}, \omega_{1}, \omega_{2}) + \mathcal{O}(|\hat{x} - \hat{y}|^{2})
= \Pi_{\hat{x}}(\hat{x} - \hat{y}) \cdot \Pi_{\hat{x}} \mathcal{Z}(\hat{x}, \omega_{1}, \omega_{2}) + \mathcal{O}(|\hat{x} - \hat{y}|^{2}).$$
(4.42)

We split the integral (4.41) in several parts depending on $\Pi_{\hat{x}}^{\perp}(\omega_j)$ for $j \in \{1, 2\}$. Let

$$\mathcal{S}_{\hat{x},\ell}^{\pm} := \left\{ \omega \in \mathbb{S}^{n-1} : \pm \Pi_{\hat{x}}^{\perp}(\omega) > \ell \right\}$$

and $\chi^j_{\hat{x},\underline{\ell}}:\mathbb{S}^{n-1}\to [0,1]$ for $j\in\{0,+,-\}$ be smooth cutoffs such that

$$\mathbb{1}_{\mathcal{S}_{\hat{x},2\ell}^{\pm}} \leq \chi_{\hat{x},\underline{\ell}}^{\pm} \leq \mathbb{1}_{\mathcal{S}_{\hat{x},\ell}^{\pm}} \qquad \qquad \chi_{\hat{x},\underline{\ell}}^{0} + \chi_{\hat{x},\underline{\ell}}^{-} + \chi_{\hat{x},\underline{\ell}}^{+} = \mathbb{1}_{\mathbb{S}^{n-1}}.$$

We first consider the integrals

$$(\hat{x}, \omega_2) \mapsto \frac{1}{(2\pi\hbar)^{2(n-1)}} \int \chi_{\hat{x},\underline{\ell}}^{\pm}(\omega_1) e^{i\frac{\Psi_1(\hat{x},\hat{y},\omega_1,\omega_2)}{\hbar}} B_3(\hat{x},\hat{y},\omega_1,\omega_2) d\hat{y}d\omega_1. \tag{4.43}$$

Since the map $\mathbb{S}^{n-1} \ni \omega \mapsto \Pi_{\hat{x}}\omega$ is a diffeomorphism from $\mathcal{S}_{\hat{x},\underline{\ell}}^{\pm}$ to its image, we can make a change of variables

$$\hat{y} \to u = \Pi_{\hat{x}}(\hat{y} - \hat{x}), \qquad \omega_1 \to v = \Pi_{\hat{x}}(\omega_1),$$

in (4.43). We obtain

$$(4.43) = \frac{1}{(2\pi\hbar)^{2(n-1)}} \int e^{i\frac{\Psi_2(\hat{x},\omega_2,u,v)}{\hbar}} B_4(\hat{x},\omega_2,u,v) du dv$$

where the amplitude B_4 belongs to S^1 and the new phase Ψ_2 satisfies, by (4.42) and (4.26),

$$\Psi_2(\hat{x}, \omega_2, u, v) = u \cdot (R_+ v - R_- \Pi_{\hat{x}}(\omega_2)) + \mathcal{O}(u^2), \qquad R_{\pm} = \sqrt{\lambda - V(\hat{x}) \pm \sigma}.$$

Any critical point of Ψ_2 with respect to (u, v) satisfies u = 0 and the Hessian is non-degenerate with determinant $R_+^{2(n-1)}$ and $R_{\pm} \geq \sqrt{\underline{c}}$ (there is at most one critical point given by $v = \Pi_{\hat{x}}(\omega_2)R_-/R_+$). Thus, by Proposition A.1 with d = 2(n-1), one has the following estimate, uniformly in $(\hat{x}, \omega_2) \in \partial\Omega \times \mathbb{S}^{n-1}$ (as well as the auxiliary parameters $(\sigma, \lambda) \in \mathbb{R}^2$):

$$(4.43) = \mathcal{O}(\hbar^{n-1}).$$

By the same argument, with the same uniformity,

$$\int \chi_{\hat{x},\underline{\ell}}^{\pm}(\omega_2) e^{i\frac{\Psi_1(\hat{x},\hat{y},\omega_1,\omega_2)}{\hbar}} B_3(\hat{x},\hat{y},\omega_1,\omega_2) d\hat{y} d\omega_2 = \mathcal{O}(\hbar^{n-1}).$$

Hence, we are left to deal with the integral

$$\hat{x} \mapsto \frac{1}{(2\pi\hbar)^{2(n-1)}} \int \chi_{\hat{x},\underline{\ell}}^0(\omega_1) \chi_{\hat{x},\underline{\ell}}^0(\omega_2) e^{i\frac{\Psi_1(\hat{x},\hat{y},\omega_1,\omega_2)}{\hbar}} B_3(\hat{x},\hat{y},\omega_1,\omega_2) d\hat{y} d\omega_1 d\omega_2.$$

Since $\hat{x} \mapsto \Pi_{\hat{x}}$ is smooth (we assume that $\partial \Omega$ is smooth) and $\partial_{\hat{y}} = \Pi_{\hat{y}}(\partial_y) + \mathcal{O}(x-y)$, according to (4.38),

$$\partial_{\hat{y}} \Psi_1(\hat{x}, \hat{y}, \omega_1, \omega_2) = \Pi_{\hat{x}} \mathcal{Z}(\hat{x}, \omega_1, \omega_2) + \mathcal{O}(\underline{\ell}) \qquad \text{for } \{\hat{y} \in \partial\Omega : |\hat{y} - \hat{x}| \leq \underline{\ell}\}.$$

Thus, on the support of the previous integral, $\{|\Pi_{\hat{x}}^{\perp}(\omega_1)|, |\Pi_{\hat{x}}^{\perp}(\omega_2)| \leq \underline{\ell}, |\mathcal{Z}(\hat{x}, \omega_1, \omega_2)| > \delta\}$ with $\underline{\ell} \ll \delta$. Since \mathcal{Z} is a linear combination of (ω_1, ω_2) , see (4.24), we have $\Pi_{\hat{x}}\mathcal{Z}(\hat{x}, \omega_1, \omega_2) = \mathcal{Z}(\hat{x}, \omega_1, \omega_2) + \mathcal{O}(\underline{\ell})$ and then

$$|\partial_{\hat{y}}\Psi_1(\hat{x},\hat{y},\omega_1,\omega_2)| \ge \delta/2.$$

This shows that the phase has no critical point in the previous integral. Hence, by Lemma A.1 (non-stationary phase version), we conclude that uniformly in $\hat{x} \in \partial\Omega$ (as well as the auxiliary parameters $(\sigma, \lambda) \in \mathbb{R}^2$),

$$\int \chi^0_{\hat{x},\underline{\ell}}(\omega_1)\chi^0_{\hat{x},\underline{\ell}}(\omega_2)e^{i\frac{\Psi_1(\hat{x},\hat{y},\omega_1,\omega_2)}{\hbar}}B_3(\hat{x},\hat{y},\omega_1,\omega_2)\mathrm{d}\hat{y}\mathrm{d}\omega_1\mathrm{d}\omega_2 = \mathcal{O}(\hbar^\infty).$$

Altogether, this establishes that the integrals (4.41), (4.39) as well as the original integral are all $\mathcal{O}(\hbar^{1-n})$ with the required uniformity in $(\sigma, \lambda) \in \mathbb{R}^2$. Going back to (4.28), we conclude that $I_{1,\hbar} = \mathcal{O}(\hbar^{1-n})$ and this also finalizes the proof of Proposition 4.9.

4.3 Case of a contractible open set

It remains to study integrals of the form given by the right-hand-side of Proposition 4.9. We first do so in the case where Ω is topologically simple: if it is diffeomorphic to the unit ball, then we can proceed by a change of variables.

We denote $B = B_n = \{x \in \mathbb{R}^n : |x| < 1\}.$

Proposition 4.12. Let $\Omega \in \mathbb{R}^n$ be a bounded open set with a smooth boundary; suppose that there is a C^{∞} -diffeomorphism $\varphi : \mathbb{R}^n \to \mathbb{R}^n$ with $\varphi^{-1}(\Omega) = B$. Let $F \in \mathcal{F}(\mathbb{R}^{2n}, \mathbb{R}^n)$ according to Definition 4.4 and assume that as $r \to 0$,

$$F(x, x, r\omega) = rR(x) + \mathcal{O}(r^2)$$
(4.44)

uniformly for $(x,\omega) \in \Omega' \times \mathbb{S}^{n-1}$. Let $f \in C^{\infty}(\mathbb{R}^n)$ and denote $f = f\mathbb{1}_{\Omega}$. Then, as $\hbar \to 0$,

$$\int e^{i\frac{(x-y)\cdot\xi}{\hbar}} F(x,y,\xi) (f|_{\Omega}(x) - f|_{\Omega}(y))^2 d\xi dx dy = (2\pi\hbar)^{n+1} \left(\frac{\log \hbar^{-1}}{-\pi^2} \int_{\partial\Omega} R(\hat{x}) f(\hat{x})^2 d\hat{x} + \mathcal{O}(1)\right)$$

where $d\hat{x}$ denotes the volume measure on the boundary $\partial\Omega$.

Without loss of generality, we assume that $\varphi = I$ on $\mathbb{R}^n \setminus \Omega'$. Throughout this section, we denote

$$L_{\hbar} := \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} F(x,y,\xi) (f|_{\Omega}(x) - f|_{\Omega}(y))^2 d\xi dx dy. \tag{4.45}$$

The proof of Proposition 4.12 will be divided in three steps.

- In Section 4.3.1, using the map φ as a change of coordinates, we show that we can reduce (4.45) to the case where $\Omega = B$.
- In Section 4.3.2, we obtain the asymptotics of (4.45) when $\Omega = B$ by using the expression of the Fourier transform $\widehat{\mathbb{1}_B}$ in terms of a Bessel function, cf. Lemma 4.16.
- Combining these results, we complete the proof in Section 4.3.3.

These computations rely importantly on the estimates from Section 4.1 for integrals involving $\mathbb{1}_{\Omega}$.

4.3.1 Change of variables

Proposition 4.13. Under the assumptions of Proposition 4.12, define a map

$$\Phi: (x,y,\xi) \in \mathbb{R}^{3n} \mapsto \left(\varphi(x),\varphi(y),D\varphi(\tfrac{x+y}{2})^{-*}\xi\right)$$

and let $G := J_{\Phi} F \circ \Phi$. Then G is also in the class \mathcal{F} and, as $\hbar \to 0$,

$$\int e^{i\frac{(x-y)\cdot\xi}{\hbar}} F(x,y,\xi) (f|_{\Omega}(x) - f|_{\Omega}(y))^2 d\xi dx dy = \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} G(x,y,\xi) (g_B(x) - g_B(y))^2 d\xi dx dy + \mathcal{O}(\hbar^{n+1})$$

where $g_B = f|_{\Omega}(\varphi) = f(\varphi) \mathbb{1}_B$.

Proof. By construction, the map $\Phi: \mathbb{R}^{3n} \to \mathbb{R}^{3n}$ is a C^{∞} -diffeomorphism and it is straightforward to check that G is also in the class \mathcal{F} , cf. Definition 4.4 (since $\varphi = I$ on $\mathbb{R}^n \setminus \Omega'$, G has compact support). In particular, for every $k \in \mathbb{N}_0$ with $k \leq n+3$,

$$\|\partial_{\xi}^{k}G(x,y,\xi)\| \leq |\xi|^{1-k} \varkappa_{1}(\xi) \varkappa_{2}(x,y) \tag{4.46}$$

where $\varkappa_1 \in C_c^{\infty}(\mathbb{R}^n)$ and $\varkappa_2 \in C_c^{\infty}(\mathbb{R}^{2n})$.

By a Taylor expansion, we have for $(x, y, \xi) \in \mathbb{R}^{3n}$,

$$(\varphi(x) - \varphi(y)) \cdot D\varphi(\frac{x+y}{2})^{-*}\xi = (x-y) \cdot \xi + Q(\frac{x+y}{2}, x-y)\xi$$

where $Q: \mathbb{R}^{2n} \to \mathbb{R}^n$ is C_c^{∞} and vanishes at zero along with its two first derivatives:

$$|Q(v,u)| \le C|u|^3. (4.47)$$

Let us denote

$$\check{G}(\frac{x+y}{2}, x - y, \xi) = G(x, y, \xi),$$
 and $\Gamma(\frac{x+y}{2}, x - y) = (g_B(x) - g_B(y))^2.$

Let $\chi \in C_c^{\infty}(\mathbb{R}^n)$ be equal to 1 near 0. By a change of variable using Φ .

$$\int e^{i\frac{(x-y)\cdot\xi}{\hbar}} F(x,y,\xi) (f|_{\Omega}(x) - f|_{\Omega}(y))^{2} d\xi dx dy = \int e^{i\frac{u\cdot\xi + Q(v,u)\xi}{\hbar}} \check{G}(v,u,\xi) \Gamma(v,u) d\xi dv du$$

$$= \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} G(x,y,\xi) (g_{B}(x) - g_{B}(y))^{2} d\xi dx dy$$

$$+ I_{\hbar} + \mathcal{O}(\hbar^{n+1})$$
(4.48)

where

$$I_{\hbar} = \int (1 - \chi(\frac{\xi}{\hbar})) e^{i\frac{u\xi}{\hbar}} \left(e^{i\frac{Q(v,u)\xi}{\hbar}} - 1 \right) \breve{G}(v,u,\xi) \Gamma(v,u) d\xi dv du.$$

Here, we used Lemma 4.6 together with the fact that $G \in \mathcal{F}$ to introduce the cutoff $1 - \chi(\frac{\xi}{\hbar})$ in the integral (4.48), up to an error $\mathcal{O}(\hbar^{n+1})$.

Now, our goal is now to show that $I_{\hbar} = \mathcal{O}(\hbar^{n+1})$ as $\hbar \to 0$. The argument is similar to the proof of Lemma 4.7, albeit the induction process is more technical. We will integrate by parts repeatedly in the variable u, stopping every time a derivative hits either $\chi(\frac{\xi}{\hbar})$ or $e^{i\frac{Q}{\hbar}}$. To this end, we introduce the following integrals for $k \in \mathbb{N}_0$ with $k \leq n+2$:

$$J_{k,\hbar} = 2i \int e^{i\frac{u\xi}{\hbar}} u \cdot \partial_{\xi} \chi(\frac{\xi}{\hbar}) \left(e^{i\frac{Q(v,u)\xi}{\hbar}} - 1 \right) \frac{u^{\otimes k} \cdot \partial_{\xi}^{k} \check{G}(v,u,\xi)}{|u|^{2(k+1)}} \Gamma(u,v) d\xi du dv,$$

$$I_{k,\hbar} = \int (1 - \chi(\frac{\xi}{\hbar})) e^{i\frac{u\xi + Q(v,u)\xi}{\hbar}} \frac{Q(v,u)u}{|u|^{2}} \frac{u^{\otimes k} \cdot \partial_{\xi}^{k} \check{G}(v,u,\xi)}{|u|^{2k}} \Gamma(u,v) d\xi du dv.$$

Provided we can show that all these integrals are finite, one has

 I_{\hbar} $= i\hbar \int e^{i\frac{u\xi}{\hbar}} \frac{u \cdot \partial_{\xi}}{|u|^{2}} \left\{ (1 - \chi(\frac{\xi}{\hbar})) \left(e^{i\frac{Q(v,u)\xi}{\hbar}} - 1 \right) \check{G}(v,u,\xi) \right\} \Gamma(u,v) d\xi du dv$ $= i\hbar \int (1 - \chi(\frac{\xi}{\hbar})) e^{i\frac{u\xi}{\hbar}} \left(e^{i\frac{Q(v,u)\xi}{\hbar}} - 1 \right) \frac{u \cdot \partial_{\xi} \check{G}(v,u,\xi)}{|u|^{2}} \Gamma(u,v) d\xi du dv - I_{0,\hbar} - J_{0,\hbar}$ $= (i\hbar)^{2} \int (1 - \chi(\frac{\xi}{\hbar})) e^{i\frac{u\xi}{\hbar}} \left(e^{i\frac{Q(v,u)\xi}{\hbar}} - 1 \right) \frac{u^{\otimes 2} \cdot \partial_{\xi}^{2} \check{G}(v,u,\xi)}{|u|^{4}} \Gamma(u,v) d\xi du dv - I_{0,\hbar} - J_{0,\hbar} - i\hbar (I_{1,\hbar} + J_{1,\hbar})$ $= \cdots$ $= (i\hbar)^{n+3} \int (1 - \chi(\frac{\xi}{\hbar})) e^{i\frac{u\xi}{\hbar}} \left(e^{i\frac{Q(v,u)\xi}{\hbar}} - 1 \right) \frac{u^{\otimes (n+3)} \cdot \partial_{\xi}^{n+3} \check{G}(v,u,\xi)}{|u|^{2(n+3)}} \Gamma(u,v) d\xi du dv - \sum_{k=0}^{n+2} (i\hbar)^{k} (I_{k,\hbar} + J_{k,\hbar}).$ (4.40)

Let us first bound the $J_{k,\hbar}$ integrals. Using (4.46) and that $\left|e^{i\frac{Q(v,u)\xi}{\hbar}}-1\right| \leq \frac{C}{\hbar}|u|^3|\xi|$ for $(u,v)\in\mathbb{R}^{2n}$, for any $k\in\mathbb{N}$ with $k\leq n+2$,

$$|J_{k,\hbar}| \leq \frac{C}{\hbar} \int |\xi| |\partial_{\xi} \chi(\frac{\xi}{\hbar})| \frac{\|\partial_{\xi}^{k} \check{G}(v, u, \xi)\|}{|u|^{k-2}} \Gamma(u, v) d\xi du dv$$

$$\leq C \hbar^{n+1-k} \int |\partial_{\xi} \chi(\xi)| |\xi|^{2-k} \frac{(g_{B}(x) - g_{B}(y))^{2}}{|x - y|^{k-2}} \varkappa_{2}(x, y) d\xi du dv.$$

where we rescaled $\frac{\xi}{\hbar} \leftarrow \xi$ since $|\partial_{\xi} \chi|$ has compact support in $\mathbb{R}^n \setminus \{0\}$. By Proposition 4.2, these integrals are finite so that for every $k \in \mathbb{N}_0$ with $k \leq n+2$,

$$J_{k,\hbar} = \mathcal{O}(\hbar^{1+n-k}). \tag{4.50}$$

To estimate $I_{k,\hbar}$, we reverse the change of variables Φ , and obtain

$$I_{k,\hbar} = \int (1 - \chi(\frac{A(x,y)\xi}{\hbar}))e^{i\frac{(x-y)\cdot\xi}{\hbar}} L_k(x,y,\xi) (f|_{\Omega}(x) - f|_{\Omega}(y))^2 dxdyd\xi$$
(4.51)

where $A: \mathbb{R}^{n \times n} \to \operatorname{GL}_n$ is a smooth map, and L_k is the image by Φ^{-1} of the function

$$(x,y,\xi) \in \mathbb{R}^{3n} \mapsto \frac{Q(\frac{x+y}{2},x-y)(x-y)}{|x-y|^2} \frac{(x-y)^{\otimes k} \cdot \partial_{\xi}^{k} \check{G}(x,y,\xi)}{|x-y|^{2k}} |D\phi(x)| |D\phi(y)| \det(A(x,y))^{-1}.$$

Using (4.46) and (4.47), together with the fact that Φ maps the diagonal x = y to itself, we verify that for every $j \in \mathbb{N}$,

$$\left|\partial_{\xi}^{j} L_{k}(x, y, \xi)\right| \leq C|x - y|^{2-k}|\xi|^{1-k-j} \varkappa(\xi, x, y).$$

where \varkappa is compactly supported in a neighbourhood of x = y. Hence, by Proposition 4.7, for every $k \ge n+2$,

$$I_{k,\hbar} = \mathcal{O}(\hbar^{1+n-k}).$$

Going back to (4.49) and using the previous estimates with k = 0, we obtain

$$I_{\hbar} = i\hbar \int \overline{\chi}(\frac{\xi}{\hbar}) e^{i\frac{u\xi}{\hbar}} \left(e^{i\frac{Q(v,u)\xi}{\hbar}} - 1 \right) \frac{u \cdot \partial_{\xi} \check{G}(v,u,\xi)}{|u|^2} \Gamma(u,v) d\xi du dv + \mathcal{O}(\hbar^{1+n}).$$

The leading term is of the form as I_{\hbar} , so we can repeat the process. By induction, after $k \in \mathbb{N}$ integrations by part, we obtain

$$I_{\hbar} = (i\hbar)^k \int \overline{\chi}(\frac{\xi}{\hbar}) e^{i\frac{u\xi}{\hbar}} \left(e^{i\frac{Q(v,u)\xi}{\hbar}} - 1 \right) \frac{u^{\otimes k} \cdot \partial_{\xi}^k \breve{G}(v,u,\xi)}{|u|^{2k}} \Gamma(u,v) d\xi du dv - \sum_{j < k} (i\hbar)^j (I_{j,\hbar} + J_{j,\hbar}).$$

The error terms are all controlled as above. Hence, taking k = n + 3, we obtain

$$|I_{\hbar}| \leq \hbar^{n+3} \int \overline{\chi}(\frac{\xi}{\hbar}) e^{i\frac{u\xi}{\hbar}} \left| e^{i\frac{Q(v,u)\xi}{\hbar}} - 1 \right| \frac{\|\partial_{\xi}^{n+3} G(v,u,\xi)\|}{|u|^{n+3}} \Gamma(u,v) d\xi du dv + \mathcal{O}(\hbar^{1+n})$$

$$\leq C\hbar^{n+2} \int \overline{\chi}(\frac{\xi}{\hbar}) |\xi|^{-1-n} \frac{(f|\Omega(x) - f|\Omega(y))^2}{|x - y|^n} \varkappa_2(x,y) dx dy d\xi + \mathcal{O}(\hbar^{1+n})$$

$$= \mathcal{O}(\hbar^{1+n})$$

where we used again Proposition 4.2 to bound the last integral.

In the end, going back to formula (4.48), we conclude that

$$\int e^{i\frac{(x-y)\cdot\xi}{\hbar}} F(x,y,\xi) (f|_{\Omega}(x) - f|_{\Omega}(y))^2 d\xi dx dy = \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} G(x,y,\xi) (g_B(x) - g_B(y))^2 d\xi dx dy + O(\hbar^{n+1})$$
 as claimed.

4.3.2 Asymptotics in case of the ball

The goal of this section is to obtain the following asymptotics.

Proposition 4.14. Let $f \in C^{\infty}(\mathbb{R}^n)$, $F \in \mathcal{F}(\mathbb{R}^{2n}, \mathbb{R}^n)$ and denote

$$q(r) = \frac{1}{r} \int_{\mathbb{S}^{n-1}} F(\omega, \omega, r\omega) f(\omega)^2 d\omega, \qquad r > 0.$$
 (4.52)

This function is continuous on \mathbb{R}_+ , with compact support, and we assume that $q(r) \to Q$ as $r \to 0$ and that

$$\int_0^1 \left| \frac{q(r) - Q}{r} \right| \mathrm{d}r < \infty. \tag{4.53}$$

Then, by (4.45) with $\Omega = B$,

$$L_{\hbar} = -4\hbar^{n+1} ((2\pi)^{n-1} Q \log \hbar^{-1} + \mathcal{O}(1)).$$

We first simplify the integral L_{\hbar} of (4.45), using the estimates from Section 4.1.

Lemma 4.15. Let $f \in C^{\infty}(\mathbb{R}^n)$, $F \in \mathcal{F}(\mathbb{R}^{2n}, \mathbb{R}^n)$ and $g(x, \xi) := F(x, x, \xi)f(x)^2$ for $(x, \xi) \in \mathbb{R}^{2n}$. Then $g \in \mathcal{F}(\mathbb{R}^n, \mathbb{R}^n)$ and

$$L_{\hbar} = -2 \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} g(\frac{x+y}{2},\xi) \mathbb{1}_B(x) \mathbb{1}_B(y) d\xi dx dy + \mathcal{O}(\hbar^{n+1}).$$

Proof. By symmetry of the integral in question, we can assume that

$$L_{\hbar} = 2 \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} F(x,y,\xi) (f_B(x) - f_B(y)) f_B(x) d\xi dx dy.$$

Then, we can write

$$\frac{L_{\hbar}}{2} = \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} F(x,y,\xi) (f(x) - f(y)) f_B(x) d\xi dx dy$$
(4.54)

$$+ \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} G(x,y,\xi) \mathbb{1}_{B}(x) \mathbb{1}_{B^c}(y) d\xi dx dy$$

$$(4.55)$$

where $G(x, y, \xi) := F(x, y, \xi) f(x) f(y)$.

We begin by showing that (4.54) is negligible. Since $(x,y) \mapsto F(x,y,\xi)$ is $C_c^{\infty}(\mathbb{R}^{2n})$ for $\xi \in \mathbb{R}^n$, by a Taylor expansion,

$$(f(x) - f(y))F(x, y, \xi) = (x - y) \cdot a(\frac{x+y}{2}, \xi) + b(x, y, \xi)$$

where $a \in \mathcal{F}$, $b \in \mathcal{F}$ and

$$a(x,\xi) = \partial f(x)F(x,x,\xi), \qquad b(x,y,\xi) = \mathcal{O}(|x-y|^3).$$

In particular, the function $(x, y, \xi) \mapsto b(x, y, \xi) f_B(x)$ satisfies the assumptions of Lemma 4.7. This implies that

$$(4.54) = 2 \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} (x-y) \cdot a(\frac{x+y}{2},\xi) f_B(x) d\xi dx dy + \mathcal{O}(\hbar^{n+1})$$

Then, we perform an integration by part with respect to ξ and a change of variables $\frac{x+y}{2} \leftarrow v$, $(x-y) \leftarrow u$, we obtain

$$\int e^{i\frac{(x-y)\cdot\xi}{\hbar}}(x-y)\cdot a(\frac{x+y}{2},\xi)f_B(x)d\xi dxdy = (i\hbar)\int e^{i\frac{(x-y)\cdot\xi}{\hbar}}a'(\frac{x+y}{2},\xi)f_B(x)d\xi dxdy$$
$$= (i\hbar)\int e^{i\frac{u\cdot\xi}{\hbar}}a'(v,\xi)f_B(v+\frac{u}{2})d\xi dudv$$

where $a' = \operatorname{div}_{\xi} a$ is L^{∞} with compact support. Using the Fourier transform, computing the integral over u, and then over v, we obtain

$$\int e^{i\frac{(x-y)\cdot\xi}{\hbar}}(x-y)\cdot a(\frac{x+y}{2},\xi)f_B(x)d\xi dxdy = i\hbar(2\pi)^{\frac{n}{2}}\int e^{-i\frac{v\cdot\xi}{\hbar}}a'(v,\xi)\overline{\widehat{f_B}(\frac{2\xi}{\hbar})}d\xi dv$$

$$= i\hbar(2\pi)^n\int \widehat{a'}(\frac{\xi}{\hbar},\xi)\overline{\widehat{f_B}(\frac{2\xi}{\hbar})}d\xi \tag{4.56}$$

where for $(\zeta, \xi) \in \mathbb{R}^{2n}$ with $\xi \neq 0$,

$$\widehat{a}'(\zeta,\xi) = \frac{1}{(2\pi)^{\frac{n}{2}}} \int e^{-iv\cdot\zeta} a'(v,\xi) dv = \operatorname{div}_{\xi} \left(\frac{1}{(2\pi)^{\frac{n}{2}}} \int e^{-iv\cdot\zeta} a(v,\xi) dv \right).$$

Since $a \in \mathcal{F}$, $\zeta \mapsto \widehat{a'}(\zeta, \xi)$ is in the Schwartz class for $\xi \neq 0$ and $\xi \mapsto \widehat{a'}(\zeta, \xi)$ is L^{∞} . In the end, by scaling and Cauchy-Schwarz,

$$\int \widehat{a'}(\frac{\xi}{\hbar},\xi) \overline{\widehat{f_B}(\frac{2\xi}{\hbar})} d\xi = \mathcal{O}(\hbar^n ||a'||_{L^2 \times L^{\infty}} ||f_B||_{L^2}).$$

and we conclude that $(4.54) = \mathcal{O}(\hbar^{n+1})$.

Going back to (4.55), we have

$$G(x,y,\xi) = g(\frac{x+y}{2},\xi) + e(x,y,\xi)$$
 where $g \in \mathcal{F}$, $e \in \mathcal{F}$ and $e(x,y,\xi) = \mathcal{O}(|x-y|^2)$,

so that the function

$$(x,y,\xi)\mapsto e(x,y,\xi)\mathbb{1}_B(x)\mathbb{1}_{B^c}(y)$$

satisfies the assumptions of Lemma 4.7 – the condition (4.12) for $j \le n+2$ follow directly from (4.5) with $\Omega = B$. Altogether, this implies that

$$\frac{L_{\hbar}}{2} = \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} g(\frac{x+y}{2},\xi) \mathbb{1}_{B}(x) \mathbb{1}_{B^c}(y) d\xi dx dy + \mathcal{O}(\hbar^{n+1}). \tag{4.57}$$

Moreover, as in (4.56), we have

$$\int e^{i\frac{(x-y)\cdot\xi}{\hbar}} g(\frac{x+y}{2},\xi) \mathbb{1}_B(x) d\xi dx dy = (2\pi)^n \int \widehat{g}(\frac{\xi}{\hbar},\xi) \overline{\widehat{\mathbb{1}_B}(\frac{2\xi}{\hbar})} d\xi$$

where for $(\zeta, \xi) \in \mathbb{R}^{2n} \mapsto \widehat{g}(\zeta, \xi)$ satisfies $|\widehat{g}(\zeta, \xi)| \leq |\xi| \varkappa(\zeta)$ where \varkappa is in the Schwartz class; since $g \in \mathcal{F}$. Using this bound and the fact that $\widehat{\mathbb{1}}_B \in L^{\infty}$, we obtain

$$\int e^{i\frac{(x-y)\cdot\xi}{\hbar}} g(\frac{x+y}{2},\xi) \mathbb{1}_B(x) d\xi dx dy = \mathcal{O}(\hbar^{n+1}).$$

Hence, the claim follows from (4.57).

The proof relies on the following explicit asymptotics.

Lemma 4.16. Let J_{ν} denote the Bessel function of the first kind for $\nu \in \mathbb{R}_{+}$. As $\xi \to +\infty$,

$$\widehat{\mathbb{1}}_B(\xi) = \frac{J_{n/2}(|\xi|)}{|\xi|^{n/2}} = \frac{2\cos(|\xi| + c_n) + \mathcal{O}(|\xi|^{-1})}{\sqrt{2\pi}|\xi|^{(n+1)/2}}.$$
(4.58)

Proof. The relationship between Bessel functions and the Fourier transform of $\mathbb{1}_B$ where B is the unit ball in \mathbb{R}^n is classical: see e.g. [19], formula (A.5). The asymptotics of Bessel functions are also classical, see [49, formula (10.17.3)].

Proof of Proposition 4.14.

Starting from Lemma 4.15 and making a change of variables $\frac{x+y}{2} \leftarrow v$, $(x-y) \leftarrow u$, we have

$$L_{\hbar} = -2 \int e^{i\frac{u\cdot\xi}{\hbar}} g(v,\xi) \mathbb{1}_B(v+\frac{u}{2}) \mathbb{1}_B(v-\frac{u}{2}) d\xi du dv + \mathcal{O}(\hbar^{n+1}).$$

Since $g \in \mathcal{F}$, by Lemma 4.6, we can introduce a cutoff $\overline{\chi}(\frac{\xi}{\hbar})$ in the previous integral, up to an error $\mathcal{O}(\hbar^{n+1})$. Moreover, computing this Fourier transform as a convolution (with our convention, $(2\pi)^{\frac{n}{2}}\widehat{uv} = \widehat{u} * \widehat{v}$ for functions $u, v \in L^2(\mathbb{R}^n)$),

$$\int e^{i\frac{u\cdot\xi}{\hbar}} \mathbb{1}_B(v+\frac{u}{2}) \mathbb{1}_B(v-\frac{u}{2}) du = 2^n \int e^{-2iv\cdot\zeta} \widehat{\mathbb{1}_B}(\zeta+\frac{\xi}{\hbar}) \widehat{\mathbb{1}_B}(\frac{\xi}{\hbar}-\zeta) d\zeta = \int e^{-iv\cdot\zeta} \widehat{\mathbb{1}_B}(\frac{\zeta}{2}+\frac{\xi}{\hbar}) \widehat{\mathbb{1}_B}(\frac{\zeta}{2}-\frac{\xi}{\hbar}) d\zeta$$

This implies that

$$L_{\hbar} = -2 \int \overline{\chi}(\frac{\xi}{\hbar}) e^{-iv\cdot\zeta} g(v,\xi) \widehat{\mathbb{1}}_{B}(\frac{\zeta}{2} + \frac{\xi}{\hbar}) \widehat{\mathbb{1}}_{B}(\frac{\zeta}{2} - \frac{\xi}{\hbar}) dv d\zeta d\xi + \mathcal{O}(\hbar^{n+1}).$$

This integral is well-defined since $\mathbb{1}_B \in L^2(\mathbb{R}^n)$ and g has compact support on \mathbb{R}^{2n} . Moreover we can take the Fourier transform of the smooth function $v \in \mathbb{R}^n \mapsto g(v,\xi)$ for a fixed ξ , we obtain

$$L_{\hbar} = -2(2\pi)^{\frac{n}{2}} \int \overline{\chi}(\frac{\xi}{\hbar}) \widehat{g}(\zeta, \xi) \widehat{\mathbb{1}_{B}}(\frac{\zeta}{2} + \frac{\xi}{\hbar}) \widehat{\mathbb{1}_{B}}(\frac{\zeta}{2} - \frac{\xi}{\hbar}) d\zeta d\xi + \mathcal{O}(\hbar^{n+1})$$

where for $k \in \mathbb{N}$, $\zeta \in \mathbb{R}^n$, r > 0,

$$\sup_{\xi \in \mathbb{R}^n} |\widehat{g}(\zeta, \xi)| \le \frac{C_k}{(1 + |\zeta|^2)^k} \quad \text{and} \quad \sup_{\omega \in \mathbb{S}^{n-1}} \|\widehat{g}(\cdot, r\omega)\|_{L^1} \le Cr. \tag{4.59}$$

Now, using the asymptotics (4.58), for every $\zeta \in \mathbb{R}^n$, $\xi = r\omega$ with $(r, \omega) \in \mathbb{R}_+ \times \mathbb{S}^{n-1}$, as $r \to \infty$,

$$\widehat{\mathbb{1}_B}(\zeta + \xi)\widehat{\mathbb{1}_B}(\zeta - \xi) = \frac{2\cos(r + \omega \cdot \zeta + c_n)\cos(r - \omega \cdot \zeta + c_n) + \mathcal{O}(r^{-1})}{\pi r^{n+1}}$$
$$= \frac{\cos(2r + 2c_n) + \cos(2\zeta \cdot \omega) + \mathcal{O}(r^{-1})}{\pi r^{n+1}}.$$

The error term is uniform for $|\zeta| \leq r^{\alpha}$ for any $0 < \alpha < 1$ and $\omega \in \mathbb{S}^{n-1}$, so by (4.59), we can substitute these asymptotics into the previous formula for L_{\hbar} (up to a negligible error). This yields 3 terms,

$$L_{\hbar} = -\frac{1}{\pi^2} (2\pi\hbar)^{n+1} (I_{1,\hbar} + I_{2,\hbar} + I_{3,\hbar}) + \mathcal{O}(\hbar^{n+1})$$
(4.60)

where

$$L_{1,\hbar} = (2\pi)^{-\frac{n}{2}} \int \overline{\chi}(\frac{r}{\hbar}) \widehat{g}(\zeta, r\omega) \cos(\zeta \cdot \omega) \frac{\mathrm{d}r}{r^2} \mathrm{d}\zeta \mathrm{d}\omega$$
$$L_{2,\hbar} = (2\pi)^{-\frac{n}{2}} \int \overline{\chi}(\frac{r}{\hbar}) \widehat{g}(\zeta, r\omega) \cos(2r/\hbar + 2c_n) \frac{\mathrm{d}r}{r^2} \mathrm{d}\zeta \mathrm{d}\omega$$

and $L_{3,\hbar}$ is controlled using (4.59) as

$$L_{3,\hbar} = \mathcal{O}\left(\hbar \int \overline{\chi}(\frac{r}{\hbar}) \sup_{\omega \in \mathbb{S}^{n-1}} \|\widehat{g}(\cdot, r\omega)\|_{L^{1}} \frac{\mathrm{d}r}{r^{3}}\right) = \mathcal{O}\left(\hbar \int \overline{\chi}(\frac{r}{\hbar}) \frac{\mathrm{d}r}{r^{2}}\right).$$

Then, since $\int \overline{\chi}(r) \frac{\mathrm{d}r}{r^2} < \infty$,

$$L_{3,\hbar} = \mathcal{O}(1). \tag{4.61}$$

We now turn to the highly oscillating term $L_{2,\hbar}$ that we can rewrite (by Fourier's inversion formula)

$$L_{2,\hbar} = \int \overline{\chi}(\frac{r}{\hbar})g(0,r\omega)\cos(2r/\hbar + 2c_n)\frac{\mathrm{d}r}{r^2}\mathrm{d}\omega.$$

Let $j(r) := \int g(0, r\omega) d\omega$, which we may view as an even function in $\mathcal{F}(\mathbb{R})$. So

$$L_{2,\hbar} = \Re\left(e^{2ic_n} \int \overline{\chi}(\frac{r}{\hbar}) j(r) e^{i\frac{2r}{\hbar}} \frac{\mathrm{d}r}{r^2}\right).$$

and we can make an integration by parts,

$$L_{2,\hbar} = -\frac{\hbar}{2} \Im \left(e^{2ic_n} \int \partial_r \left\{ \overline{\chi}(\frac{r}{\hbar}) j(r) \frac{1}{r^2} \right\} e^{i\frac{2r}{\hbar}} dr \right).$$

Then, we control $L_{2,\hbar}$ as follows,

$$\left| \int \partial_r \overline{\chi}(\frac{r}{\hbar}) j(r) r^{-2} dr \right| \le C \int |\chi'(r)| \frac{dr}{r} < \infty,$$

similarly

$$\hbar \int \left| \overline{\chi}(\frac{r}{\hbar}) j(r) r^{-3} \right| dr \le C \hbar \int \overline{\chi}(\frac{r}{\hbar}) \frac{dr}{r^2} = C \int \overline{\chi}(r) \frac{dr}{r^2} < \infty$$

and

$$\hbar \int \left| \overline{\chi}(\frac{r}{\hbar}) j'(r) r^{-2} \right| dr \le C \int \overline{\chi}(r) \frac{dr}{r^2} < \infty.$$

We conclude that

$$L_{2,\hbar} = \mathcal{O}(1). \tag{4.62}$$

We finally turn to the main $L_{1,\hbar}$ which we compute using Fourier's inversion formula,

$$L_{1,\hbar} = \int \overline{\chi}(\frac{r}{\hbar}) g(\omega, r\omega) \frac{\mathrm{d}r}{r^2} \mathrm{d}\zeta \mathrm{d}\omega$$

where we used the invariance of the Haar measure $d\omega$ on \mathbb{S}^{n-1} . Using the notation (4.52), this yields

$$L_{1,\hbar} = \int \overline{\chi}(\frac{r}{\hbar})q(r)\frac{\mathrm{d}r}{r}.$$

Since $q(r) \to Q$ as $r \to 0$, if $Q \neq 0$, this integral cannot be bounded as $\hbar \to 0$. However, it diverges logarithmically under the assumption (4.53). Indeed, we can split if C > 0 is sufficiently large (depending only on the cutoff),

$$L_{1,\hbar} = Q \left(\int_{C\hbar}^{1} \frac{\mathrm{d}r}{r} + \int_{0}^{C\hbar} \overline{\chi}(\frac{r}{\hbar}) \frac{\mathrm{d}r}{r} \right) + \int_{1}^{\infty} \frac{q(r)}{r} \mathrm{d}r + \mathcal{O} \left(\int_{0}^{1} \left| \frac{q(r) - Q}{r} \right| \mathrm{d}r \right)$$

The last three terms are $\mathcal{O}(1)$ since $\int_0^C \overline{\chi}(r) \frac{\mathrm{d}r}{r} < \infty, q \in C_c$ and (4.53) holds. This shows that

$$L_{1,\hbar} = Q \log \hbar^{-1} + \mathcal{O}(1).$$

Combining this asymptotics with (4.61), (4.62) into (4.60), we conclude that as $\hbar \to 0$,

$$L_{\hbar} = -\frac{1}{\pi^2} (2\pi\hbar)^{n+1} (Q \log \hbar^{-1} + \mathcal{O}(1)).$$

4.3.3 Proof of Proposition 4.12

We now put together the different steps of the proof of Proposition 4.12. Recall that $\varphi : \mathbb{R}^n \to \mathbb{R}^n$ is a C^{∞} -diffeomorphism such that $\varphi^{-1}(\Omega) = B$. First by Proposition 4.13 with $g = f(\varphi)$

$$L_{\hbar} = \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} F(x,y,\xi) (f|_{\Omega}(x) - f|_{\Omega}(y))^{2} d\xi dx dy$$
$$= \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} G(x,y,\xi) (g|_{B}(x) - g|_{B}(y))^{2} d\xi dx dy + \mathcal{O}(\hbar^{n+1})$$

and on the diagonal,

$$G(x, x, \xi) = F(\varphi(x), \varphi(x), D\varphi(x)^{-*}\xi) J_{\varphi}(x), \qquad (x, \xi) \in \mathbb{R}^{2n}.$$

Here we used that by construction, $J_{\Phi}(x, y, \xi) = J_{\varphi}(x)J_{\varphi}(y)J_{\varphi}^{-1}(\frac{x+y}{2}).$

Then, as $G \in \mathcal{F}(\mathbb{R}^{2n}, \mathbb{R}^n)$, by Proposition 4.14, we obtain

$$L_{\hbar} = (2\pi\hbar)^{n+1} \left(\frac{1}{\pi^2} Q \log \hbar^{-1} + \mathcal{O}(1)\right)$$

where, if it exists,

$$Q = \lim_{r \to 0^+} \left(\frac{1}{r} \int_{\mathbb{S}^{n-1}} G(\omega, \omega, r\omega) g(\omega)^2 d\omega \right).$$

It remains to argue under the assumption (4.44) this limits is well-defined and (4.53) holds. We have as $r \to 0$,

$$G(w, w, rw) = rR(\varphi(\omega))J_{\varphi}(\omega)|D\varphi(\omega)^{-*}\omega| + \mathcal{O}(r^{2})$$

$$= rR(\hat{\varphi}(\omega))J_{\hat{\varphi}}(\omega) + \mathcal{O}(r^{2})$$
(4.63)

where the map $\hat{\varphi}: \partial B \to \partial \Omega$ is the C^{∞} -diffeomorphism induced by φ on the boundary and the errors are controlled uniformly over $\omega \in \partial B$. To obtain (4.63), note that for any $\omega \in \partial B$, we can decompose $T_{\omega}(\mathbb{R}^n) = T_{\omega}(\partial B) \oplus \mathbb{R}\omega$ and, with $\hat{x} = \varphi(\omega)$, $T_{\hat{x}}(\mathbb{R}^n) = T_{\hat{x}}(\partial \Omega) \oplus \mathbb{R}\nu(\hat{x})$ where $\nu(\hat{x})$ is the (unit) normal to $\partial \Omega$ at \hat{x} . In this decomposition, since $\varphi^{-1}(\Omega) = B$ the matrix of the differential $D\varphi$ of the map φ has the following from

$$D\varphi = \begin{pmatrix} D\hat{\varphi} & 0 \\ \star & \alpha \end{pmatrix}, \quad \text{where} \quad \alpha(\omega) = \nu(\hat{x}) \cdot D\varphi(\omega)\omega, \quad \hat{x} = \varphi(\omega). \tag{4.64}$$

In particular $\alpha > 0$ on ∂B and, by taking determinants, this implies that for $\omega \in \partial B$,

$$J_{\varphi}(\omega) = J_{\hat{\varphi}}(\omega)\alpha(\omega) \tag{4.65}$$

Moreover by definition of ν , for any $\omega \in \partial B$ and $v \in T_{\omega}(\partial B) = \omega^{\perp}$,

$$0 = \nu(\varphi(\omega)) \cdot D\varphi(\omega)u,$$

which shows that $D\varphi(\omega)^*\nu(\varphi(\omega))$ is proportional to ω . According to (4.64), this implies that for $\omega \in \partial B$,

$$D\varphi(\omega)^*\nu(\varphi(\omega)) = \alpha(\omega)\omega$$

and since $|\nu(\varphi(\omega))| = 1$, we conclude that

$$1 = \alpha(\omega)|D\varphi(\omega)^{-*}\omega|$$

Combined with (4.65), this proves formula (4.63). Then, we deduce from this expansion that

$$Q = \int_{\partial B} R(\hat{\varphi}(\omega)) f(\hat{\varphi}(\omega))^2 J_{\hat{\varphi}}(\omega) d\omega = \int_{\partial \Omega} R(\hat{x}) f(\hat{x})^2 d\hat{x}$$

by a change of variable. In addition, we have

$$\int_0^1 \int_{\mathbb{S}^{n-1}} \left| \frac{G(w, w, rw) - rR(\hat{\varphi}(\omega)) J_{\hat{\varphi}}(\omega)}{r^2} \right| g(\omega)^2 dr d\omega < \infty.$$

which guarantees that the condition (4.53) holds. This completes the proof.

4.4 Partition of unity; Proof of Theorem 1.

We now combine our previous results, with a decomposition using a partition of unity, to prove Theorem 1. Recall that $\Omega \in \mathcal{D}$ is an open set with a smooth boundary and, according to Proposition 4.9, $(2\pi\hbar)^{n-1} \| [\widetilde{\Pi}, f|_{\Omega}] \|_{\mathbf{J}^2}^2 = \mathcal{Q}(f) + \mathcal{O}(1)$ as $\hbar \to 0$ where \mathcal{Q}_{Ω} is the quadratic form

$$Q_{\Omega}: f \mapsto \frac{-1}{(2\pi\hbar)^{n+1}} \int e^{i\frac{(x-y)\cdot\xi}{\hbar}} F(x,y,\xi) (f|_{\Omega}(x) - f|_{\Omega}(y))^2 d\xi dx dy$$

acting on smooth functions.

Moreover, by Proposition 4.12 and (4.13), in case Ω is contractible (C^{∞} -diffeormorphic to B_n), one has $\mathcal{Q}_{\Omega}(f) = \log \hbar^{-1} \mathcal{V}(f) + \mathcal{O}(1)$ as $\hbar \to 0$ where \mathcal{V} is the quadratic form

$$\mathcal{V}_{\Omega}: f \mapsto \frac{\mathbf{c}_{n-1}}{\pi^2} \int_{\partial \Omega} |V(\hat{x})|^{\frac{n-1}{2}} f(\hat{x})^2 d\hat{x}$$

$$\tag{4.66}$$

acting on smooth functions.

We claim that there is a smooth partition of unity $(\chi_j)_{1 \leq j \leq J}$ such that

- $\chi_j|_{\Omega} = \chi_j|_{\Omega_j}$ where Ω_j is a contractible set, for $j \in [J]$.
- $(\chi_i + \chi_j)|_{\Omega} = (\chi_i + \chi_j)|_{\Omega_{i,j}}$ where $\Omega_{i,j}$ is a contractible set, for $i, j \in [J]$.

These sets are arbitrary smooth contractible sets. The condition $\chi|_{\Omega} = \chi|_{\Omega_i}$ is equivalent to

$$\Omega \cap (\operatorname{supp} \chi) = \Omega_j \cap (\operatorname{supp} \chi),$$

so we can always choose a partition sufficiently fine so that these conditions hold. This partition is finite since Ω is compact.

Thus, it follows that

$$\begin{aligned} \mathcal{Q}_{\Omega}(1) &= \mathcal{Q}_{\Omega}\left(\sum_{j=1}^{J} \chi_{j}\right) = \sum_{i,j=1}^{J} \mathcal{Q}_{\Omega}(\chi_{j} + \chi_{i}) - \sum_{j=1}^{J} \mathcal{Q}_{\Omega}(\chi_{j}) \\ &= \sum_{i,j=1}^{J} \mathcal{Q}_{\Omega_{ij}}(\chi_{j} + \chi_{i}) - \sum_{j=1}^{J} \mathcal{Q}_{\Omega_{j}}(\chi_{j}) \\ &= \log \hbar^{-1} \left(\sum_{i,j=1}^{J} \mathcal{V}_{\Omega_{ij}}(\chi_{j} + \chi_{i}) - \sum_{j=1}^{J} \mathcal{V}_{\Omega_{j}}(\chi_{j})\right) + \mathcal{O}(1) \end{aligned}$$

Using that $\mathcal{V}_{\Omega_{ij}}(\chi_j + \chi_i) = \mathcal{V}_{\Omega}(\chi_j + \chi_i)$ and $\mathcal{V}_{\Omega_j}(\chi_j) = \mathcal{V}_{\Omega}(\chi_j)$ for $i, j \in [J]$, we conclude that

$$Q_{\Omega}(1) = \log \hbar^{-1} \mathcal{V}_{\Omega}(\sum_{j=1}^{J} \chi_j) + \mathcal{O}(1).$$

Similarly, we have for any $f \in C^{\infty}(\mathbb{R}^n)$, $\mathcal{Q}_{\Omega}(f) = \log \hbar^{-1} \frac{c_{n-1}}{\pi^2} \mathcal{V}_{\Omega}(f) + \mathcal{O}(1)$ as $\hbar \to 0$.

According to Proposition 2.5, this shows that for any open set $\Omega \in \mathcal{D}$ with smooth boundary and any $f \in C^{\infty}(\mathbb{R}^n)$,

$$(2\pi\hbar)^{n-1} \|[\Pi, f|_{\Omega}]\|_{J^2}^2 = (2\pi\hbar)^{n-1} \|[\widetilde{\Pi}, f|_{\Omega}]\|_{J^2}^2 + \mathcal{O}(1) = \log \hbar^{-1} \mathcal{V}_{\Omega}(f) + \mathcal{O}(1) \quad \text{as } \hbar \to 0.$$

This completes the proof (with $\mu = 0$ and V(x) < 0 for $x \in \Omega$).

4.5 Central limit theorem; Proof of Theorem 2.

The Gaussian fluctuations of the counting statistics, after rescaling, are due to the determinantal structure of the free fermions point process \mathbf{X} and the fact that for a non-trivial smooth set Ω , var $\mathbf{X}(\Omega) \to \infty$ as $\hbar \to \infty$. In the random matrix context, this observation is due to [13] and it has been extended to general determinantal processes in [63]. By [63, Thm 1], we deduce from Theorem 1 that for any open set $\Omega \in \mathcal{D}$ with smooth boundary and any $f \in C^{\infty}(\mathbb{R}^n)$, (in distribution) as $\hbar \to 0$,

$$\frac{\mathbf{X}(f|_{\Omega}) - \mathbb{E}[\mathbf{X}(f|_{\Omega})]}{\sqrt{-(2\pi\hbar)^{1-n}\log\hbar}} \Rightarrow \mathcal{N}_{0,\mathcal{V}_{\Omega}(f)}.$$

In particular, if Ω has disjoint components $\{\Omega_1, \dots, \Omega_k\}$, $\mathcal{V}_{\Omega} = \sum_{j=1}^k \mathcal{V}_{\Omega_j}$ according to (4.66), then

$$\left(\frac{\mathbf{X}(\Omega_1) - \mathbb{E}[\mathbf{X}(\Omega_1)]}{\sqrt{-(2\pi\hbar)^{1-n}\log\hbar}}, \cdots, \frac{\mathbf{X}(\Omega_k) - \mathbb{E}[\mathbf{X}(\Omega_k)]}{\sqrt{-(2\pi\hbar)^{1-n}\log\hbar}}\right) \Rightarrow \mathcal{N}_{0,\Sigma}$$

where $\Sigma = \operatorname{diag}(\mathcal{V}_{\Omega_1}, \cdots, \mathcal{V}_{\Omega_k}).$

Note that this argument cannot be directly applied to a collection of sets $\Omega_1, \dots, \Omega_k \in \mathcal{D}$ (open, with smooth boundaries) with intersecting boundaries. However, we can use the off-diagonal decay of the regularized kernel $\tilde{\Pi}$ and the estimates from Lemma 4.3 to show that the cross terms are negligible.

Lemma 4.17. Let $\Omega \in \mathcal{D}$ be open. The kernel (2.3) satisfies for $(x,y) \in \Omega^2$,

$$|\widetilde{\Pi}(x,y)|^2 \le \frac{C\hbar^{1-n}}{(\hbar + |x-y|)^{n+1}}$$

Proof. Let $\chi: \mathbb{R}^n \to [0,1]$ be a smooth cutoff such that $\chi = 1$ on Ω and $\chi = 0$ on Ω' where Ω' is a neighborhood of Ω in \mathcal{D} . We proceed as in Section 4.2.1, we make a change of variable $\xi = r\omega$ with $r > \underline{c}$ (cf. (2.1)), $\omega \in \mathbb{S}^{n-1}$ and we apply the stationary phase method to the integral (2.3) in the variables (t, r) with (x, y, ω, λ) fixed. The critical point is non-degenerate, given by (4.18) with $\sigma = 0$, so we obtain

$$\chi(x)\widetilde{\Pi}(x,y)\chi(y) = \frac{1}{(2\pi\hbar)^n} \int e^{i\frac{\Psi_1(x,x-y,\omega,\lambda)}{\hbar}} b(x,y,\omega,\lambda) \mathbb{1}\{\lambda \le 0\} d\omega d\lambda$$
 (4.67)

where $b \in S^1(\mathbb{R}^{3n+1})$ is supported on $\{|\lambda|, |x-y| \leq \underline{\ell}\}$, and the phase Ψ_1 vanishes on the diagonal $\{x=y\}$. This already implies the following saturated bound, locally uniformly in the bulk:

$$|\widetilde{\Pi}(x,y)| \le C\hbar^{-n}$$
.

Then, we can write

$$\Psi_1(x, u, \omega, \lambda) = u \cdot \zeta(x, u, \omega, \lambda)$$

where the map $\zeta : \mathbb{R}^{3n+1} \to \mathbb{R}^n$ is smooth on supp(b) and make a change of coordinates $(\lambda, \omega) \to \zeta(x, u, \omega, \lambda)$ in (4.67). As in Section 4.2.2 (albeit using a simpler analysis), we claim that

$$\zeta(x, u, \omega, \lambda) = R(x, \lambda)\omega + \mathcal{O}(u), \qquad R = \sqrt{\lambda - V(x)}$$

and the error is smooth, so the matrix

$$\frac{\partial \zeta}{\partial \lambda \partial \omega} = R^{-1} \mathbf{I}_n + \mathcal{O}(u)$$

is non-degenerate since $R \geq \underline{c}$ on $\mathrm{supp}(b)$ and $\underline{\ell} \ll \underline{c}$. Hence, there is $g \in S^1(\mathbb{R}^{3n})$, supported on $\{|x-y| \leq \underline{\ell}\}$, so that

$$(4.67) = \frac{1}{(2\pi\hbar)^n} \int e^{i\frac{(x-y)\cdot\zeta}{\hbar}} g(x,y,\zeta) \mathbb{1}\{\lambda(x,y,\zeta) \le 0\} d\zeta$$

where the map $\lambda: \mathbb{R}^{3n} \to \mathbb{R}$ is smooth with on the diagonal $\{x = y\}$,

$$\lambda(x, x, \zeta) = |\zeta|^2 + V(x).$$

By perturbation, $\operatorname{Hess}_{\zeta}\lambda(x,y,\zeta)$ is positive-definite for $x\in\Omega', |x-y|\leq\underline{\ell}$ so that $\{\zeta\in\mathbb{R}^n:\lambda(x,u,\zeta)\leq0\}$ is a strongly convex body. Thus, by [31, Corollary 7.7.15], if $u\in\mathbb{R}^n\mapsto\Lambda(x,y,u)$ denotes the Fourier transform of $\zeta\in\mathbb{R}^n\mapsto\mathbbm{1}\{\lambda(x,y,\zeta)\leq0\}$, then

$$\sup_{x,y\in\Omega'} |\Lambda(x,y,u)| \le \frac{C}{1+|u|^{\frac{n+1}{2}}}.$$

This is to be compared with the case of a ball (Lemma 4.16). This implies that

$$\chi(x)\widetilde{\Pi}(x,y)\chi(y) = \hbar^{-n}\Lambda(x,y,\cdot) * \widehat{g}(x,y,\cdot)|_{\frac{x-y}{\hbar}}$$

where $u \in \mathbb{R}^n \mapsto \widehat{g}(x,y,u)$ is a Schwartz function. We conclude that

$$|\chi(x)\widetilde{\Pi}(x,y)\chi(y)| \leq C\hbar^{-\frac{n-1}{2}}|x-y|^{-\frac{n+1}{2}}.$$

which is the appropriate away from the diagonal.

Remark 4.18. The bound from Lemma 4.17 corresponds to $|\widetilde{\Pi}(x,y)| \leq C\hbar^{-n}|\widehat{\mathbb{1}}\{\cdot \leq 1\}(\frac{x-y}{\hbar})|$ for $(x,y) \in \Omega^2$ (inside the bulk). This is consistent with the scaling limit of the regularized kernel $\widetilde{\Pi}$. Namely, we showed in [19] that for $x \in \mathcal{D}$, $\hbar^n \widetilde{\Pi}(x + \hbar u, x + \hbar v) \to K(u - v)$ where $K = C_n \widehat{\mathbb{1}}\{\cdot \leq c_n\}$ for some constants C_n, c_n depending only on the dimension n.

Given two open sets $\Omega_1, \Omega_2 \in \mathcal{D}$ with smooth boundary such that $|\partial \Omega_1 \cap \partial \Omega_2| = 0$, Lemmas 4.17 and 4.3 imply that as $\hbar \to \infty$,

$$\big|\operatorname{tr}([\widetilde{\Pi}, \mathbbm{1}_{\Omega_1}][\widetilde{\Pi}, \mathbbm{1}_{\Omega_2}])\big| \leq C \hbar^{1-n} \int \frac{|(\mathbbm{1}_{\Omega_1}(x) - \mathbbm{1}_{\Omega_1}(y))(\mathbbm{1}_{\Omega_2}(x) - \mathbbm{1}_{\Omega_2}(y))|}{(\hbar + |x - y|)^{n+1}} \mathrm{d}x \mathrm{d}y = o\big(\log(\hbar^{-1})\hbar^{1-n}\big).$$

Then, expanding the square, under the assumptions of Theorem 2, it holds for $\alpha \in \mathbb{R}^k$,

$$\frac{[\widetilde{\Pi}, \sum_{j=1}^k \alpha_k \mathbb{1}_{\Omega_k}]\|_{J^2}^2}{(2\pi\hbar)^{n-1} (\log \hbar^{-1})} = \sum_{j=1}^k \frac{\alpha_k [\widetilde{\Pi}, \mathbb{1}_{\Omega_k}]\|_{J^2}^2}{(2\pi\hbar)^{n-1} (\log \hbar^{-1})} + o(1) \quad \text{as } \hbar \to 0.$$

Replacing Π by the projector Π using Proposition 2.5, we conclude by Theorem 1 that as $\hbar \to 0$,

$$\frac{\|[\Pi, \sum_{j=1}^k \alpha_k \mathbb{1}_{\Omega_k}]\|_{\mathcal{J}^2}^2}{(2\pi\hbar)^{n-1}(\log \hbar^{-1})} = \sum_{j=1}^k \alpha_k \mathcal{V}_{\Omega_k}(1) + o(1).$$

Again, applying Soshnikov's result [63, Thm 1], by the Cramér–Wold argument, this completes the proof of Theorem 2.

One has the following more precise covariance estimate in the case of a relatively non-singular intersection.

Remark 4.19. If $H^{\beta}(\partial\Omega_1\cap\partial\Omega_2)<+\infty$ for some $\beta< n-1$, then the last covariance bound improves to

$$Cov(\mathbf{X}(\mathbb{1}_{\Omega_1}), \mathbf{X}(\mathbb{1}_{\Omega_2})) = \mathcal{O}(\hbar^{1-n})$$

using the second part of Lemma 4.3. At this scale, we cannot use the regularised projector from Proposition 2.5 to compute the limit of the covariance. This difficulty already appears in the variance estimates in [19].

5 Trace norm of commutators & entropy estimates

5.1 Proof of Theorem 4: Multi-scale argument

The goal of this section is to prove Theorem 4, that is, to show that for any open set $\Omega \subseteq \mathcal{D}$ with a smooth boundary

$$\|[\Pi, \mathbb{1}_{\Omega}]\|_{\mathbb{T}^1} = \mathcal{O}(\hbar^{1-n} |\log \hbar|^2).$$
 (5.1)

We begin by a simple truncation step.

Lemma 5.1. Let $\vartheta : \mathbb{R} \to [0,1]$ be any smooth function which equal to 1 on $[-\underline{c}/2,\underline{c}/2]$. Let $\Gamma_{\pm} = \vartheta \mathbb{1}_{\mathbb{R}_{\pm}}(H)$ and $\Theta_{\pm} = \vartheta \mathbb{1}_{\mathbb{R}_{+}}(w)$. It holds

$$\|[\Pi, \mathbb{1}_{\Omega}]\|_{J^{1}} \leq \|\Gamma_{-}\Theta_{+}\Gamma_{+}\Theta_{-}\|_{J^{1}} + \|\Gamma_{+}\Theta_{+}\Gamma_{-}\Theta_{-}\|_{J^{1}} + \|\Gamma_{+}\Theta_{-}\Gamma_{-}\Theta_{+}\|_{J^{1}} + \|\Gamma_{-}\Theta_{-}\Gamma_{+}\Theta_{+}\|_{J^{1}} + \mathcal{O}(\hbar^{1-n}).$$

Proof. First observe that since the operator H and the map w are bounded from below

$$\begin{cases}
\Pi = \Gamma_{-} + f_{1}(H) \\
1 - \Pi = \Gamma_{+} + 1 - f_{2}(H)
\end{cases}
\begin{cases}
\mathbb{1}_{\Omega} = \Theta_{-} + f_{1}(w) \\
1 - \Pi = \Theta_{+} + 1 - f_{2}(w)
\end{cases}$$
(5.2)

where $f_j: \mathbb{R} \to [0,1]$ are smooth functions, supported in $(-\infty,\underline{c}]$ with $f_1\mathbb{1}_{\mathbb{R}_+} = 0$ and $(1-f_2)\mathbb{1}_{\mathbb{R}_-} = 0$. Second, observe that

$$\begin{split} [\Pi, \mathbb{1}_{\Omega}] &= \mathbb{1}_{\Omega^c} \Pi \mathbb{1}_{\Omega} - \mathbb{1}_{\Omega} \Pi \mathbb{1}_{\Omega^c} \\ &= \Pi \mathbb{1}_{\Omega^c} \Pi \mathbb{1}_{\Omega} + (1 - \Pi) \mathbb{1}_{\Omega^c} \Pi \mathbb{1}_{\Omega} - (1 - \Pi) \mathbb{1}_{\Omega} \Pi \mathbb{1}_{\Omega^c} - \Pi \mathbb{1}_{\Omega} \Pi \mathbb{1}_{\Omega^c}. \end{split}$$

Then, using that $\mathbb{1}_{\Omega}\mathbb{1}_{\Omega^c}=0$, we can change the first and fourth terms as follows:

$$[\Pi, \mathbb{1}_{\Omega}] = -\Pi \mathbb{1}_{\Omega^{c}} (1 - \Pi) \mathbb{1}_{\Omega} + (1 - \Pi) \mathbb{1}_{\Omega^{c}} \Pi \mathbb{1}_{\Omega} - (1 - \Pi) \mathbb{1}_{\Omega} \Pi \mathbb{1}_{\Omega^{c}} + \Pi \mathbb{1}_{\Omega} (1 - \Pi) \mathbb{1}_{\Omega^{c}}. \tag{5.3}$$

Every term on the RHS of (5.3) can be handled in the same way, so we focus on the last one. Using (5.2), since $f_1(H)(1-\Pi)=0$ and $\Gamma_-(1-f_2(H))=0$, one has

$$\Pi \mathbb{1}_{\Omega} (1 - \Pi) \mathbb{1}_{\Omega^c} = \Gamma_- \mathbb{1}_{\Omega} \Gamma_+ \mathbb{1}_{\Omega^c} + [f_1(H), \mathbb{1}_{\Omega}] (1 - \Pi) \mathbb{1}_{\Omega^c} + \Gamma_- [\mathbb{1}_{\Omega}, 1 - f_2(H)] \mathbb{1}_{\Omega^c}$$

According to Proposition 2.6, both $||[f_j(H), \mathbb{1}_{\Omega}]||_{\mathbb{I}^1} = \mathcal{O}(\hbar^{1-n})$ for $j \in \{1, 2\}$, so

$$\Pi \mathbb{1}_{\Omega} (1 - \Pi) \mathbb{1}_{\Omega^c} = \Gamma_- \mathbb{1}_{\Omega} \Gamma_+ \mathbb{1}_{\Omega^c} + \mathcal{O}_{\mathbb{I}^1} (\hbar^{1-n}).$$

Similarly,

$$\Gamma_{-}\mathbb{1}_{\Omega}\Gamma_{+}\mathbb{1}_{\Omega^{c}} = \Gamma_{-}\Theta_{-}\Gamma_{+}\Theta_{+} + \Gamma_{-}[f_{1}(\mathbf{w}), \Gamma_{+}]\mathbb{1}_{\Omega^{c}} + \Gamma_{-}\Theta_{-}[\Gamma_{+}, 1 - f_{2}(\mathbf{w})].$$

Observe that according to Lemma 2.9 and (1.10),

$$[f_1(\mathbf{w}), \Gamma_{\pm}] = [f_1(\mathbf{w}), f_j(H)] - [f_1(\mathbf{w}), \Pi] = \mathcal{O}_{J^1}(\hbar^{1-n}).$$

Thus, we conclude that

$$\Pi \mathbb{1}_{\Omega} (1 - \Pi) \mathbb{1}_{\Omega^c} = \Gamma_- \Theta_- \Gamma_+ \Theta_+ + \mathcal{O}_{J^1} (\hbar^{1-n}).$$

Again, every term in the bound from Lemma 5.1 can be handled in the same way. We proceed using a dyadic argument to *isolate the contributions from different scales* that we will estimate based on Proposition 3.1. To setup the dyadic decompositions, on top of the notations from Section 2.1, we use the following convention throughout this section.

Notations 5.1.

- Let $w : \mathbb{R}^n \to \mathbb{R}$ be a smooth function such that $\Omega = \{w < 0\} \in \mathcal{D}$ and $\partial_x w(x) \neq 0$ for all $x \in \partial \Omega = \{w = 0\}$. In particular, we assume that $\partial_x w \neq 0$ on $\{|w| \leq \underline{c}\}$ by choosing the constant \underline{c} sufficiently small.
- Let $\vartheta : \mathbb{R} \to [0,1]$ supported in $[-\underline{c},\underline{c}]$ and equal to 1 on $[-\underline{c}/2,\underline{c}/2]$. Let $\chi : \mathbb{R} \to [0,1]$ supported in [-1,1] and equal to 1 on $[-\underline{c},\underline{c}]$.
- Let $L := |\log_2(\hbar)|$ and let $\eta_k := \hbar 2^k$ for $k \in \mathbb{N}_{< L}$.
- Let $\chi_0 := \chi(\hbar^{-1}\cdot)$ and

$$\chi_k : x \mapsto \chi_0(2^{-k}x) - \chi_0(2^{1-k}x), \qquad k \in \mathbb{N}_{$$

We also let $\chi_L: x \mapsto \chi(x) - \chi_0(2^{1-L}x)$. We use a similar notation for $(\vartheta_k)_{k=0}^L$.

- We use the shorthand notation, $\chi_k^{\pm} = \chi_k \mathbb{1}_{\mathbb{R}_{\pm}}$ for $k \in [0, L]$; similarly for $(\vartheta_k^{\pm})_{k=0}^L$.
- We let $\psi_k^{\pm} := \vartheta \cdot (\chi_k^{\pm} * \rho_h)$ for $k \in [1, L]$, as in the Notation of Section 2.1. In particular, $\chi_k^{\pm}, \psi_k^{\pm} : \mathbb{R} \to [0, 1]$ are in S^{η_k} for $k \in [1, L]$.
- For $k \in [1, L]$, let $\check{\vartheta}_k^{\pm}$ be in S^{η_k} such that

$$\chi_k^{\pm} = \chi_k^{\pm} \check{\chi}_k^{\pm}, \qquad \chi_k^{\mp} \check{\chi}_m^{\pm} = 0, \qquad \text{for } k, m \in [1, L].$$
(5.4)

This yields a dyadic decomposition of the operators appearing in Lemma 5.1.

Lemma 5.2. Using the notation 5.1, we have $\Theta_{\pm} = \sum_{k=0}^{L} \vartheta_k(w)$ and $\Gamma_{\pm} = \sum_{k=1}^{L} \widetilde{\chi}_k^{\pm}(H) + \Upsilon_{\pm}$ where the errors satisfy

$$\|\Upsilon_{\pm}\|_{J^{1}} = \mathcal{O}(\hbar^{1-n})$$
 $\|\Upsilon_{\pm}\|_{L^{2} \to L^{2}} \le 1.$

Proof. The decomposition of Θ_{\pm} is obvious, so we focus on the decomposition of Γ_{+} (Γ_{-} is handled similarly). By construction $\sum_{k=0}^{L} \chi_{k}^{+} = \mathbb{1}_{\mathbb{R}_{+}}$ on $[-\underline{c},\underline{c}]$. So, since ρ is a Schwartz mollifier, we claim that for every $m \in \mathbb{N}$, there is $\varkappa_{m} : \mathbb{R} \to [-1,1]$, smooth with compact support so that

$$\sum_{k=1}^{L} \chi_k^+ * \rho_{\hbar} = \mathbb{1}_{\mathbb{R}_+} + \varkappa_m(\hbar^{-1} \cdot) + \mathcal{O}(\hbar^m) \quad \text{with a smooth error on } [-\underline{\mathbf{c}}, \underline{\mathbf{c}}].$$

By Lemma 2.7, this implies that $\sum_{k=1}^{L} \tilde{\chi}_{k}^{\pm}(H) = \Gamma_{+} + \Upsilon_{+}$ where the operator Υ_{+} satisfies

$$\|\Upsilon_\pm\|_{J^1}=\mathcal{O}(\hbar^{1-n}) \qquad \qquad \|\Upsilon_\pm\|_{L^2\to L^2}\leq 1.$$

We are now ready to proceed to obtain the estimate (5.1).

Proof of Theorem 4. According to Lemma 5.1, it suffices to show that $\|\Gamma_{+}\Theta_{+}\Gamma_{-}\Theta_{-}\|_{J^{1}} \leq C\hbar^{1-n}L^{2}$; since the other terms are handled similarly. By Lemma 5.2,

$$\Gamma_{+}\Theta_{+}\Gamma_{-}\Theta_{-} = \sum_{i,k\geq 1} \sum_{j,m\geq 0} \widetilde{\chi}_{i}^{+}(H)\vartheta_{j}^{+}(\mathbf{w})\widetilde{\chi}_{k}^{-}(H)\vartheta_{m}^{-}(\mathbf{w}) + \mathcal{O}_{\mathbf{J}^{1}}(\hbar^{1-n})$$

where the indices range up to L. The idea is to split this sum in two parts:

$$\mathbf{I} := \sum_{\substack{i,k \geq 1; j,m \geq 0 \\ i+j \leq k+m}} \widetilde{\chi}_i^+(H) \vartheta_j^+(\mathbf{w}) \widetilde{\chi}_k^-(H) \vartheta_m^-(\mathbf{w}) \qquad \text{and} \qquad \mathbf{II} := \sum_{\substack{i,k \geq 1; j,m \geq 0 \\ i+j > k+m}} \widetilde{\chi}_i^+(H) \vartheta_j^+(\mathbf{w}) \widetilde{\chi}_k^-(H) \vartheta_m^-(\mathbf{w})$$

Both sums will be handled in a similar way, so we focus on I for now. Using (5.4), one has

$$\widetilde{\chi}_i^+(H)\vartheta_j^+(\mathbf{w})\widetilde{\chi}_k^-(H)\vartheta_m^-(\mathbf{w}) = \widetilde{\chi}_i^+(H)\vartheta_j^+(\mathbf{w})\big[\big[\widetilde{\chi}_k^-(H),\vartheta_m^-(\mathbf{w})\big],\widecheck{\vartheta}_m^-(\mathbf{w})\big].$$

Then, we can bound

$$\|\widetilde{\chi}_{i}^{+}(H)\vartheta_{j}^{+}(\mathbf{w})\widetilde{\chi}_{k}^{-}(H)\vartheta_{m}^{-}(\mathbf{w})\|_{\mathbf{J}^{1}} \leq \|\widetilde{\chi}_{i}^{+}(H)\vartheta_{j}^{+}(\mathbf{w})\|_{\mathbf{J}^{2}} \begin{cases} \|\widetilde{\chi}_{k}^{-}(H)\vartheta_{m}^{-}(\mathbf{w})\|_{\mathbf{J}^{2}} & \text{if } \eta_{k}\eta_{m} \leq \hbar \\ \|\left[\left[\widetilde{\chi}_{k}^{-}(H),\vartheta_{m}^{-}(\mathbf{w})\right],\widecheck{\vartheta}_{m}^{-}(\mathbf{w})\right]\right\|_{\mathbf{J}^{2}} & \text{if } \eta_{k}\eta_{m} > \hbar \end{cases}$$

According to Proposition 3.1, there exists a constant C > 0 so that for any $k \in [1, L]$ and $m \in [0, L]$,

$$\begin{cases} \left\| \left[\left[\widetilde{\chi}_k^{\pm}(H), \vartheta_m^{\pm}(\mathbf{w}) \right], \widecheck{\vartheta}_m^{\pm}(\mathbf{w}) \right] \right\|_{\mathbf{J}^2}^2 \leq C \hbar^{3-n} \eta_k^{-2} \eta_m^{-2} = C \hbar^{-1-n} 2^{-2(k+m)} & \text{if } \eta_k \eta_m > \hbar, \\ \left\| \widetilde{\chi}_k^{\pm}(H) \vartheta_m^{\pm}(\mathbf{w}) \right\|_{\mathbf{J}^2}^2 \lesssim \hbar^{-n} \eta_k \| \vartheta_m^{\pm}(\mathbf{w}) \|_{L^2}^2 \leq C \hbar^{-n} \eta_k \eta_m = C \hbar^{2-n} 2^{k+m}. \end{cases}$$

Here we used that $\|\vartheta_m^{\pm}(\mathbf{w})\|_{L^2}^2 \lesssim \eta_m$ for $m \in [0, L]$. Altogether, this implies that

$$\begin{split} \|\mathbf{I}\|_{\mathbf{J}^{1}} &\leq \sum_{\substack{i,k \geq 1; j,m \geq 0\\ i+j \leq k+m}} \|\widetilde{\chi}_{i}^{+}(H)\vartheta_{j}^{+}(\mathbf{w})\widetilde{\chi}_{k}^{-}(H)\vartheta_{m}^{-}(\mathbf{w})\|_{\mathbf{J}^{1}} \\ &\leq C^{2} \bigg(\hbar^{2-n} \sum_{\substack{i+j \leq k+m\\ h2^{k+m} < 1}} 2^{\frac{i+j+k+m}{2}} + \hbar^{\frac{1}{2}-n} \sum_{\substack{i+j \leq k+m\\ 1 < h2^{k+m}}} 2^{\frac{i+j}{2}-(k+m)} \bigg). \end{split}$$

Since all indices range up to L, these sums are controlled by

$$\sum_{\substack{i+j \leq k+m \\ \hbar 2^{k+m} \leq 1}} 2^{\frac{i+j+k+m}{2}} \leq L^2 \sum_{\hbar 2^m \leq 1} 2^m \leq 2\hbar^{-1} L^2 \quad \text{and} \quad \sum_{\substack{i+j \leq k+m \\ 1 \leq \hbar 2^{k+m}}} 2^{\frac{i+j}{2} - (k+m)} \leq L^2 \sum_{\hbar 2^m \geq 1} 2^{-\frac{m}{2}} \leq 2\hbar^{\frac{1}{2}} L^2.$$

Hence, we conclude that

$$\|\mathbf{I}\|_{\mathbf{J}^1} = \mathcal{O}(L^2 \hbar^{1-n}). \tag{5.5}$$

We have a similar control for II, which we expect by symmetry. Let us first record that

$$\sum_{i,k \geq 1} \sum_{j,m \geq 0} \| [\widetilde{\chi}_i^+(H), \vartheta_j^+(\mathbf{w})] \|_{\mathbf{J}^2} \| [\widetilde{\chi}_k^-(H), \vartheta_m^-(\mathbf{w})] \|_{\mathbf{J}^2} = \left(\sum_{k \geq 1, m \geq 0} \| [\widetilde{\chi}_k^+(H), \vartheta_m^+(\mathbf{w})] \|_{\mathbf{J}^2} \right)^2$$

and, as above (by Proposition 3.1),

$$\sum_{k \geq 1, m \geq 0} \big\| \big[\widetilde{\chi}_k^+(H), \vartheta_m^+(\mathbf{w}) \big] \big\|_{\mathbf{J}^2} \leq C \bigg(\hbar^{1 - \frac{n}{2}} \sum_{\hbar 2^{k+m} < 1} 2^{\frac{k+m}{2}} + \hbar^{-\frac{n}{2}} \sum_{\hbar 2^{k+m} > 1} 2^{-\frac{k+m}{2}} \bigg) = \mathcal{O} \big(L \hbar^{\frac{1-n}{2}} \big).$$

This estimate shows that

$$II = \sum_{\substack{i,k \ge 1; j,m \ge 0\\ i+j > k+m}} \vartheta_j^+(\mathbf{w}) \widetilde{\chi}_i^+(H) \vartheta_m^-(\mathbf{w}) \widetilde{\chi}_k^-(H) + \mathcal{O}_{\mathsf{J}^1} (L^2 \hbar^{1-n}).$$

Now, using that

$$\vartheta_j^+(\mathbf{w})\widetilde{\chi}_i^+(H)\vartheta_m^-(\mathbf{w})\widetilde{\chi}_k^-(H) = \left[\widecheck{\vartheta}_j^+(\mathbf{w}), \left[\vartheta_j^-(\mathbf{w}), \widetilde{\chi}_i^-(H)\right]\right]\vartheta_m^-(\mathbf{w})\widetilde{\chi}_k^-(H)$$

we can proceed exactly as above to show that $\|II\|_{J^1} = \mathcal{O}(L^2\hbar^{1-n})$. This completes the proof.

5.2 Bounds for the entanglement entropy: Proof of Theorem 3

The estimates of Theorem 3 for the (entanglement) entropy follow by interpolation from the bounds for the Hilbert-Schmidt and trace norm of the commutator $[\Pi, \mathbb{1}_{\Omega}]$ of Theorems 1 and 4. Our goal is to prove the following general inequalities.

Lemma 5.3. Let **X** be a determinantal process on a Polish space \mathcal{X} associated with a (self-adjoint) operator $0 < \Pi \le 1$ locally trace-class. Then, for any open set $\Omega \subseteq \mathcal{X}$ with smooth boundary,

$$2\|[\Pi, \mathbb{1}_{\Omega}]\|_{J^{2}}^{2} \leq \mathcal{S}_{\Omega}(\mathbf{X}) \leq 4\|[\Pi, \mathbb{1}_{\Omega}]\|_{J^{2}}^{2} \log \left(\frac{\|[\Pi, \mathbb{1}_{\Omega}]\|_{J^{1}}}{\|[\Pi, \mathbb{1}_{\Omega}]\|_{J^{2}}^{2}}\right).$$

Since we establish that $\|[\Pi, \mathbb{1}_{\Omega}]\|_{J^2}^2 \simeq c\hbar^{1-n}\log(\hbar^{-1})$ for some constant c > 0 if $\Omega \in \mathcal{D}$ is an open set with smooth boundary and $\|[\Pi, \mathbb{1}_{\Omega}]\|_{J^1} \leq C\log(\hbar^{-1})\|[\Pi, \mathbb{1}_{\Omega}]\|_{J^2}^2$ as $\hbar \to 0$, Theorem 3 follows directly from Lemma 5.3. We now turn to its proof.

Let $s:[0,1]\to\mathbb{R}_+$, given by $s:\lambda\mapsto -\lambda\log(\lambda)-(1-\lambda)\log(1-\lambda)$. Recall that for a general determinantal point process \mathbf{X} , defined on a Polish space \mathcal{S} and associated with a locally trace-class operator $0<\Pi\leq 1$, the (entanglement) entropy of any open set $\Omega\in\mathcal{X}$ with smooth boundary is

$$S_{\Omega}(\mathbf{X}) = \sum_{n \in \mathbb{N}} s(\lambda_n), \tag{5.6}$$

where $\{\lambda_n\}$ denote the non-trivial eigenvalues of operator $\Pi|_{\Omega} = \mathbb{1}_{\Omega}\Pi\mathbb{1}_{\Omega} - \{\lambda_n\}$ is a countable sequence in (0,1] by the spectral theorem for locally compact operators.

The main step of the proof is to describe the relationship between the spectra of $\Pi \mathbb{1}_{\Omega} \Pi$ and the square-commutator $-[\Pi, \mathbb{1}_{\Omega}]^2$. This follows from a general result.

Lemma 5.4. Let P,Q be two self-adjoint projections on a separable Hilbert space. The non-negative operators PQP, $-[P,Q]^2$ commute. The spectral data of $-[P,Q]^2$, on the orthogonal of its kernel, consists exactly of the eigenpairs $(\lambda(1-\lambda),u)$ and $(\lambda(1-\lambda),[P,Q]u)$, where (λ,u) is an eigenpair of PQP with $\lambda>0$. Thus, given $f:[0,\frac{1}{4}]\to\mathbb{R}_+$ continuous with f(0)=0 and $g:\lambda\in[0,1]\mapsto f(\lambda(1-\lambda))$, one has

$$\operatorname{tr} f(-[P,Q]^2) = 2 \operatorname{tr} g(PQP).$$
 (5.7)

Proof. Using that $P^2 = P$ and $Q^2 = Q$, we simply compute

$$P[P,Q]^{2} = [P,Q]^{2}P = PQPQP - PQP = PQP(PQP - 1).$$
(5.8)

So $(P, -[P, Q]^2)$ commute. Then, using that $(P, -[P, Q]^2)$ and $(Q, -[P, Q]^2)$ commute, we also have

$$[PQP, [P, Q]^2] = PQ[P, Q]^2P - P[P, Q]^2QP = P[Q, [P, Q]^2]P = 0.$$

Thus $(PQP, -[P, Q]^2)$ also commute as claimed.

Since $(P, -[P, Q]^2)$ commute, the spectral data of $-[P, Q]^2$ can be decomposed into the spectral data of $-P[P, Q]^2P$ and $-(1-P)[P, Q]^2(1-P)$. Then, (5.8) shows that any eigenpair (λ, u) of PQP with $\lambda > 0$ gives an eigenpair $(\lambda(1-\lambda), u)$ of $-P[P, Q]^2P$. This means that for $\sigma > 0$, the eigenspace of $-P[P, Q]^2P$ associated with σ is $F_{\sigma} = \ker(PQP - \lambda)$ if $\sigma = \lambda(1-\lambda)$. By the spectral theorem, this gives the complete spectral data of $-P[P, Q]^2P$, on the orthogonal of its kernel.

Now, let $\sigma > 0$ be an eigenvalue of $-(1-P)[P,Q]^2(1-P)$ and let E_{σ} the associated eigenspace and T := [P,Q]. Since $E_{\sigma} \subset \ker(P)$, one has for $u \in E_{\sigma}$,

$$Tu = PQu$$
, $-[P,Q]^2u = \sigma u$ and $-P[P,Q]^2P(Tu) = -PQ[P,Q]^2u = \sigma PQu = \sigma Tu$

using the commutation relations. This shows that (σ, Tu) is an eigenpair of $-P[P,Q]^2P$. Moreover, $T^2 = -\sigma I$ on E_{σ} , so that $T: E_{\sigma} \to F_{\sigma}$ is 1-1.

In conclusion, $\ker(-[P,Q]^2 - \sigma) = \operatorname{span}\{u, Tu : u \in \ker(PQP - \lambda)\}\$ if $\sigma = \lambda(1 - \lambda)$ with $\lambda > 0$ so that choosing an orthonormal basis $\{(\lambda_k, u_k)\}_{k \in \mathbb{N}}$ of eigenfunction of PQP, by the spectral theorem $(PQP \geq 0)$, for any $f : \mathbb{R}_+ \to \mathbb{R}_+$, continuous with f(0) = 0, one has

$$\operatorname{tr} f(-[P,Q]^2) = 2 \sum_{\lambda_k > 0} f(\lambda_k (1 - \lambda_k)) = 2 \operatorname{tr} g(PQP)$$

where $g(\lambda) = f(\lambda(1-\lambda))$. Note that since $PQP \leq 1$, one can consider only test functions f defined on $[0, \frac{1}{4}]$, both sides of (5.7) are non-negative and possibly infinite.

Remark 5.5. Considering Widom's conjecture, in light of Lemma 5.4, the spectral asymptotics of Conjecture 1 are equivalent to, given $f:[0,\frac{1}{4}] \to \mathbb{R}_+$ continuous with f(0)=0,

$$\frac{\operatorname{tr} f(-[\Pi, \mathbb{1}_{\Omega}]^2)}{(2\pi\hbar)^{1-n} \log \hbar^{-1}} \to 8C_{\Omega} \int_{[0, \frac{1}{4}]} \frac{f(\sigma)}{\sigma \sqrt{1 - 4\sigma}} d\sigma.$$

We now turn to show the inequalities for the entropy.

Proof of Lemma 5.3. Considering (5.6), the map $s = f(\lambda(1-\lambda))$ where $f:[0,\frac{1}{4}] \to \mathbb{R}_+$ is given by

$$f: \sigma \mapsto -\frac{1-\sqrt{1-4\sigma}}{2}\log(\frac{1-\sqrt{1-4\sigma}}{2}) - \frac{1+\sqrt{1-4\sigma}}{2}\log(\frac{1+\sqrt{1-4\sigma}}{2}).$$

This function is continuous with f(0) = 0 (one has $f(\sigma) \sim -\sigma \log(\sigma)$ as $\sigma \to 0$) so that, according to Lemma 5.4, we can rewrite

$$S_{\Omega}(\mathbf{X}) = \operatorname{tr} f(-[\Pi, \mathbb{1}_{\Omega}]^2).$$

The proof is based on the basic inequalities, for $\sigma \in [0, \frac{1}{4}]$,

$$c\sigma \le f(\sigma) \le -2\sigma \log \sigma. \tag{5.9}$$

Observe that these function are all smooth, increasing, concave on $(0, \frac{1}{4})$ and take the values 0, respectively log 2 at the endpoints; so the inequality (5.9) follows from the fact that $f'(\sigma) \sim \log(1/\sigma)$ as $\sigma \to 0$.

Thus, by monotonicity, with $g: \sigma \mapsto -2\sigma \log \sigma$, we obtain

$$c\|[\Pi,\mathbb{1}_\Omega]\|_{J^2}^2 \leq \mathcal{S}_\Omega \leq \operatorname{tr} g(-[\Pi,\mathbb{1}_\Omega]^2).$$

We now relate the upper-bound to the trace-norm of the commutator $[\Pi, \mathbb{1}_{\Omega}]$. We can assume that $\|[\Pi, \mathbb{1}_{\Omega}]\|_{J^1} < \infty$, otherwise there is nothing to prove, in which case $\|[\Pi, \mathbb{1}_{\Omega}]\|_{J^1}^2 = \sum_{n \in \mathbb{N}} \sigma_n < \infty$ where $\{\sigma_n\}$ denotes the non-zero eigenvalues of $-[\Pi, \mathbb{1}_{\Omega}]^2$. Then, by convexity of $\sigma \mapsto \log(\sigma^{-1/2})$ on \mathbb{R}_+ and Jensen's inequality,

$$\sum_{n \in \mathbb{N}} \sigma_n \log(\sigma_n^{-1/2}) \le \left(\sum_{n \in \mathbb{N}} \sigma_n\right) \log \left(\frac{\sum_{n \in \mathbb{N}} \sigma_n^{1/2}}{\sum_{n \in \mathbb{N}} \sigma_n}\right)$$

or equivalently,

$$\frac{1}{4}\operatorname{tr} g(-[\Pi, \mathbb{1}_{\Omega}]^2) \le \|[\Pi, \mathbb{1}_{\Omega}]\|_{J^2}^2 \log \left(\frac{\|[\Pi, \mathbb{1}_{\Omega}]\|_{J^1}}{\|[\Pi, \mathbb{1}_{\Omega}]\|_{J^2}^2}\right).$$

This completes the proof.

A Stationary phase with mild amplitudes

We first recall a standard version of the stationary phase method where the amplitude is controlled in \mathscr{C}^k , independently of semiclassical parameter \hbar , and the integral depends on a parameter $z \in \mathcal{C}$. We refer for instance to theorem 7.7.5 in [31] or to our previous work [19, Proposition A.15].

Proposition A.1 (Stationary phase lemma). For $d, q \in \mathbb{N}$, let $\Omega \in \mathbb{R}^d$ and $\mathfrak{V} \in \mathbb{R}^q$ be open. Let $\Phi: \Omega \times \mathfrak{V} \to \mathbb{R}$ be a smooth function² such that $\partial_x \Phi(x,z) = 0$ has a unique solution $(x_z,z) \in \Omega \times \mathfrak{V}$ with $\Phi(x_z,z) = 0$ and the Hessian $\nabla_x^2 \Phi(x_z,z)$ is non-degenerate for $z \in \mathfrak{V}$. Let $a \in S^1(\Omega \times \mathfrak{V})$ be classical symbol. Then, there exists another classical symbol $b \in S^1(\mathfrak{V})$ so that

$$\int_{\Omega} e^{i\frac{\Phi(x,z)}{\hbar}} a(x,z) dx = (2\pi\hbar)^{\frac{d}{2}} b(z,\hbar)$$

and the principal part of b is given by $b_0(z) = a_0(x_z, z) / \sqrt{\det \nabla_x^2 \Phi(x_z, z)}$ for $z \in A$. On the other hand, if $x \mapsto \Phi(x, z)$ has no critical points in Ω , then

$$\int_{\Omega} e^{i\frac{\Phi(x,z)}{\hbar}} a(x,z) dx = \mathcal{O}(\hbar^{\infty})$$

where the error term is controlled in $S^1(\mho)$.

The goal of this section is to generalize the above expansion when the symbol $a \in S^{\delta}$ is sufficiently regular $(\delta \geq \hbar^{\frac{1}{2}})$ with an explicit control of the error. We begin by the simple case where the phase Φ has no critical point within the support of the symbol a.

Lemma A.2. For $d, q \in \mathbb{N}$, let $\Omega \in \mathbb{R}^d$ and $\mho \in \mathbb{R}^q$ be open. Let $\Phi : \Omega \times \mho \to \mathbb{R}$ be a smooth function such that $\partial_x \Phi(x, z) = 0$ has a unique solution $(x_z, z) \in \Omega \times \mho$ and the Hessian $\nabla_x^2 \Phi(x_z, z)$ is non-degenerate for all $z \in \mho$. Let $\delta \geq \hbar^{\frac{1}{2}}$ and $a \in \mathscr{C}_c(\Omega \times \mho)$ with $a \in S_x^{\delta}$ and suppose that for $z \in \mho$, a(x, z) = 0 for $x \in B(x_z, \epsilon)$ for some $\epsilon \geq \delta$. Then, for every $k \in \mathbb{N}$,

$$\sup_{z \in \mathcal{O}} \left| h^{-\frac{d}{2}} \int e^{i\frac{\Phi(x,z)}{h}} a(x,z) dx \right| \le C_k \left(\frac{h}{\delta \epsilon} \right)^k.$$

In particular, if $a \in S^{\delta}(\Omega \times \mho)$ with $\delta \geq \hbar^{\frac{1}{2}}$ and $\epsilon \delta \gg \hbar$, then as $\hbar \to 0$

$$\int e^{i\frac{\Phi(x,z)}{\hbar}} a(x,z) dx = \mathcal{O}_{\mathscr{C}^{\infty}}(\hbar^{\infty}). \tag{A.1}$$

Proof. To ease notation, we treat the case without parameter: $\mho = \emptyset$, $\Phi : \Omega \to \mathbb{R}$ is a smooth function with a unique non-degenerate critical point $x_{\emptyset} \in \Omega$ and $a \in S^{\delta}(\Omega)$ with a = 0 on $B(x_{\emptyset}, \epsilon)$ with $\epsilon \geq \delta \geq \hbar^{1/2}$. By assumptions, $x \mapsto \frac{a(x)}{|\nabla \Phi(x)|^2}$ is a smooth function, so that integrating by parts, we have

$$\int e^{i\frac{\Phi(x)}{\hbar}}a(x)\mathrm{d}x = i\hbar \int e^{i\frac{\Phi(x)}{\hbar}}\frac{a(x)\Psi(x) + \nabla a(x) \cdot \nabla \Phi(x)}{|\nabla \Phi(x)|^2}\mathrm{d}x$$

where $\Psi: x \mapsto \Delta\Phi(x) + |\nabla\Phi(x)|^2 \nabla\Phi(x) \cdot \nabla |\nabla\Phi(x)|^{-2}$ is also smooth on Ω . We can perform this operation k times, and we obtain differential operators $L^{j,k}$ of degree $\leq j$ (taking values into symmetric j-tensors) whose coefficients are smooth functions (depending on Φ) so that

$$\int e^{i\frac{\Phi(x)}{\hbar}} a_{\hbar}(x) dx = (i\hbar)^k \int e^{i\frac{\Phi(x)}{\hbar}} \sum_{0 \le j \le k} \frac{\nabla \Phi(x)^{\otimes j} \cdot L^{j;k} a(x)}{|\nabla \Phi(x)|^{2k}} dx. \tag{A.2}$$

²In fact, it is not necessary to assume that the phase Φ and the symbol a are smooth with respect to the parameter $z \in \mho$; continuity suffices.

Since x_{\emptyset} is the unique non-degenerate critical point of Φ , $|\nabla \Phi(x)| \approx |x - x_{\emptyset}|$ for $x \in \Omega$, so we have the following estimates, for every $0 \leq j \leq k$:

$$\frac{|\nabla \Phi(x)^{\otimes j} \cdot L^{j;k} a(x)|}{|\nabla \Phi(x)|^{2k}} \le C_k \delta^{-j} |x - x_{\emptyset}|^{-2k+j}, \qquad x \in \Omega.$$

Then, using that a = 0 on $B(x_{\emptyset}, \epsilon)$ with $\epsilon \geq \delta$, we obtain for every $k \geq d$ and for every $0 \leq j \leq k$,

$$\int \frac{|\nabla \Phi(x)^{\otimes j} \cdot L^{j;k} a(x)|}{|\nabla \Phi(x)|^{2k}} dx \le C_k \delta^{-j} \epsilon^{-2k+j+d} \le C_k \delta^{-k} \epsilon^{-k+d}.$$

We conclude that for every $k \geq d$,

$$\left| \hbar^{-\frac{d}{2}} \int_{\Omega} e^{i\frac{\Phi(x)}{\hbar}} a(x) dx \right| \le C_k \hbar^{k-d/2} \delta^{-k} \epsilon^{-k+d} = C_k \left(\frac{\hbar}{\delta \epsilon} \right)^{k-d} \left(\frac{\hbar^{1/2}}{\delta} \right)^d$$

which proves the claim in case $\mho = \emptyset$ using the condition $\delta \geq \hbar^{1/2}$. The argument remains the same if (Φ, a) depend continuously on a parameter $z \in \mho$ and the estimates are uniform provided that \mho is relatively compact. In particular, if $\epsilon \delta \gg \hbar$, we obtain uniformly for $z \in \mho$,

$$\int e^{i\frac{\Phi(x,z)}{\hbar}} a(x,z) dx = \mathcal{O}(\hbar^{\infty}).$$

If we apply ∂_z^{α} to the LHS, we obtain a similar integral with another symbol $a_{\alpha} \in S^{\delta}$ times $\hbar^{-|\alpha|}$, so this proves (A.1).

Proposition A.2 already implies that such oscillatory integrals are uniformly bounded.

Corollary A.3. Let Φ satisfy the assumptions of Lemma A.2. Then, for any $a \in S^{\delta}(\Omega \times \mho)$ with $\delta \geq \hbar^{\frac{1}{2}}$, as $\hbar \to 0$,

$$\hbar^{-\frac{d}{2}} \int_{\Omega} e^{i\frac{\Phi(x,z)}{\hbar}} a(x,z) dx = \mathcal{O}_{S^{\delta}}(1).$$

Proof. By a change of variable, we can assume without loss of generality that $x_z = 0$ for $z \in \mathcal{V}$ and $\delta = \hbar^{\frac{1}{2}}$. Let $\chi \in C_c^{\infty}(\mathbb{R}^d)$ be a cutoff with $\chi = 1$ on the unit ball and we split

$$\hbar^{-\frac{d}{2}} \int_{\Omega} e^{i\frac{\Phi(x,z)}{\hbar}} a(x,z) dx = \hbar^{-\frac{d}{2}} \int_{\Omega} e^{i\frac{\Phi(x,z)}{\hbar}} a(x,z) \chi(x/\delta) dx + \hbar^{-\frac{d}{2}} \int_{\Omega} e^{i\frac{\Phi(x,z)}{\hbar}} b(x,z) dx$$

where $b:(x,z)\mapsto a(x,z)(1-\chi(\delta^{-1}x))$ is also in S^{δ} . Hence, by Lemma A.2, the second integral is $\mathcal{O}(1)$, while the first integral is also $\mathcal{O}(1)$ by a direct volume estimate.

Then, we can improve this a-priori estimate by giving the asymptotic expansion of such oscillatory integrals using the stationary phase method with precise estimates on the error term.

Proposition A.4. For $d, q \in \mathbb{N}$, let $\Omega \in \mathbb{R}^d$ and $\mathfrak{V} \in \mathbb{R}^q$ be open. Let $\Phi : \Omega \times \mathfrak{V} \to \mathbb{R}$ be a smooth function such that $\partial_x \Phi(x,z) = 0$ has a unique solution $(x_z,z) \in \Omega \times \mathfrak{V}$ with $\Phi(x_z,z) = 0$ and the Hessian $\nabla_x^2 \Phi(x_z,z)$ is non-degenerate for $z \in \mathfrak{V}$. Let $a \in S^{\delta}(\Omega \times \mathfrak{V})$ with $\delta \geq \hbar^{\frac{1}{2}}$. Then for every $\ell \in \mathbb{N}_0$, as $\hbar \to 0$,

$$(2\pi i\hbar)^{-\frac{d}{2}} \int e^{i\frac{\Phi(x,z)}{\hbar}} a(x,z) dx = \sum_{0 \le j < \ell} \hbar^j L_x^j a(x_z,z) + \mathcal{O}_{S^{\delta}}((\hbar\delta^{-2})^{\ell})$$
(A.3)

where, for $j \in \mathbb{N}_0$, L_x^j is a differential operator (acting on x) of degree 2j whose coefficients depend only on the phase Φ . In addition, if $\operatorname{supp}(a) \subseteq \mathcal{A}$, then for every $\ell \in \mathbb{N}_0$ and $\epsilon \in [\delta, 1]$, as $\hbar \to 0$,

$$(2\pi i\hbar)^{-\frac{d}{2}} \int e^{i\frac{\Phi(x,z)}{\hbar}} a(x,z) dx = \mathcal{O}_{S^{\delta}} \left(\left(\frac{\hbar}{\delta \epsilon}\right)^{\ell} \right) \quad uniformly \ on \ \{z \in \mho : dist ((x_z,z),\mathcal{A}) \ge \epsilon \}.$$
 (A.4)

Proof. To compute the integral (A.3), we apply Morse lemma (with parameters) to the phase Φ , see [31, p502]. By Lemma A.2 applied with a fixed ϵ , we can assume that a is supported on an arbitrary small neighborhood $\mathcal{U} \subset \mathbb{R}^{d \times q}$ of the critical point (x_z, z) , up to a negligible error (in the sense of (A.1)). Then, by a C^{∞} diffeormorphism φ , there is coordinate system such that the critical points are (0, z) and the phase

$$\Phi(x,z) = \frac{x \cdot Hx}{2}, \qquad H = \begin{pmatrix} I_{d-\alpha} & 0\\ 0 & -I_{\alpha} \end{pmatrix},$$

where $\alpha \in [0,d]$ is the Morse index, that is, the number of negative eigenvalues of the non-degenerate matrix matrix $\nabla_x^2 \Phi(x_z,z)$ – by continuity α is constant (independent of $z \in \mathcal{V}$ and so is the phase Φ in this new coordinate system). Moreover, the symbol $b = a(\varphi) \in S^{\delta}$ with the same δ . Consequently, it suffices to show that the integral

$$R_0: z \mapsto (2\pi i\hbar)^{-\frac{d}{2}} \int e^{i\frac{xHx}{2\hbar}} b(x,z) \mathrm{d}x$$

has an expansion of the type (A.3) as $\hbar \to 0$. First, by Corollary A.3, since the phase is independent of z, by differentiating under the integral, $R_0 \in S^{\delta}(\mho)$ and it satisfies (A.4), by Lemma A.2.

Let $L := i(\nabla_x \cdot H \nabla_x)$ and $b_1 := \delta^2 L b$. Observe that if we integrate by parts twice, for any $\lambda > 0$,

$$-\partial_{\lambda} \left[\lambda^{\frac{d}{2}} \int e^{i\lambda \frac{xHx}{2\hbar}} b(x,z) dx \right] = \hbar \lambda^{\frac{d}{2}-2} \int e^{i\lambda \frac{xHx}{2\hbar}} Lb(x,z) dx. \tag{A.5}$$

Moreover, by the standard stationary phase (Proposition A.1 – for a fixed \hbar with b independent of λ):

$$\lim_{\lambda \to +\infty} \left(\frac{\lambda}{2\pi i \hbar} \right)^{\frac{d}{2}} \int e^{i\lambda \frac{xHx}{2\hbar}} b(x,z) dx = i^{-\alpha} b(0,z)$$

$$= i^{-\alpha} |\det \nabla_x^2 \Phi(x_z, z)|^{-1/2} a(x_z, z)$$
(A.6)

going back to the original coordinate system. We have $b_1 \in S^{\delta}$, so by Corollary A.3,

$$\sup_{z \in \mathcal{W}} \left| h^{-\frac{d}{2}} \lambda^{\frac{d}{2}} \int e^{i\lambda \frac{xHx}{2\hbar}} b_1(x, z) \mathrm{d}x \right| = O(1)$$

and

$$R_1(z) := (2\pi i\hbar)^{-\frac{d}{2}} \int_1^\infty \lambda^{\frac{d}{2}-2} \left\{ \int e^{i\lambda \frac{xHx}{2\hbar}} b_1(x,z) dx \right\} d\lambda$$

is also in S^{δ} (just like the symbol b_1 by differentiating under the integral). Thus, integrating both sides of (A.5) for $\lambda \in [1, \infty)$, by (A.6), we obtain

$$R_0(z) = i^{-\alpha}b(0,z) + \hbar\delta^{-2}R_1(z).$$

This proves the claim for $\ell = 1$ with $L_0 = i^{-\alpha}$ (in this coordinate system). The general expansion follows by induction: for $\ell \in \mathbb{N}$, let $b_{\ell} := \delta^{2\ell} L^{\ell} b$ and

$$R_{\ell}(z) := (2\pi i\hbar)^{-\frac{d}{2}} \int_{1}^{\infty} \lambda^{\frac{d}{2} - 2} g_{\ell}(\lambda) \left\{ \int e^{i\lambda \frac{xHx}{2\hbar}} b_{\ell}(x, z) dx \right\} d\lambda \tag{A.7}$$

with $g_1 = 1$ and for $\ell \in \mathbb{N}$, $g_{\ell+1} : [1, \infty) \to [0, 1]$ is the solution of

$$g'_{\ell+1}(\lambda) = \lambda^{-2} g_{\ell}(\lambda) \,, \qquad g_{\ell+1}(1) = 0.$$

In particular, g_{ℓ} is non-decreasing and $g_{\ell}(\infty) = \lim_{\lambda \to \infty} g_{\ell}(\lambda)$ exists for every $\ell \geq 2$. By an integration by parts, using (A.5)–(A.6),

$$R_{\ell}(z) = g_{\ell+1}(\infty)i^{-\alpha}b_{\ell}(0,z) - (2\pi i\hbar)^{-\frac{d}{2}} \int_{1}^{\infty} g_{\ell+1}(\lambda)\partial_{\lambda} \left\{ \lambda^{\frac{d}{2}} \int e^{i\lambda \frac{xHx}{2\hbar}} b_{\ell}(x,z) dx \right\} d\lambda$$
$$= g_{\ell+1}(\infty)i^{-\alpha}b_{\ell}(0,z) + \hbar\delta^{-2}R_{\ell+1}(z).$$

By induction, $g_{\ell+1}(\infty) = \int_1^\infty \frac{g'_{\ell}(\sigma)}{\sigma} d\sigma = \int_1^\infty \frac{g_{\ell-1}(\sigma)}{\sigma^3} d\sigma = \cdots = \frac{1}{(\ell-1)!} \int_1^\infty \frac{g_1(\sigma)}{\sigma^{\ell+1}} d\sigma = \frac{1}{\ell!}$, so we conclude that for $\ell \in \mathbb{N}$,

$$R_0(z) = i^{-\alpha} \sum_{k=0}^{\ell} (\hbar \delta^{-2})^k b_k(0, z) / k! + (\hbar \delta^{-2})^{\ell+1} R_{\ell+1}(z).$$

Since $b_k = \delta^{2k} L^k b$ for $k \in \mathbb{N}_0$, this proves (A.3) with $L^k = i^{-\alpha} \frac{(i\nabla \cdot H\nabla)^k}{k!}$ in this coordinate system. Going back to the original coordinates, L^k are differential operators of degree 2k whose coefficients depend only on the phase Φ with $L^0 = i^{-\alpha} |\det \nabla_x^2 \Phi|^{-1/2}$ as a multiplication operator.

Remark A.5. We give a slight generalization of Proposition A.4. Let $g: \Omega \times \mho \to \mathbb{R}$, \mathcal{C}^1 , such that $\nabla g \neq 0$ on $\{|g| < 1\}$. Let $a \in S^{\delta}(\Omega \times \mho)$ with $\delta \geq \hbar^{\frac{1}{2}}$, assume that $\operatorname{supp}(a) \subset \{|g| < 1\}$ and it holds for $\ell \in \mathbb{N}$, $\epsilon \in [\delta, 1]$,

$$a = \mathcal{O}_{S^{\delta}}(\left(\frac{\hbar}{\delta \epsilon}\right)^{\ell})$$
 uniformly on $\{|g(x,z)| > \epsilon\}.$ (A.8)

Then, setting $f(z) := Cg(x_z, z)$ for $z \in \mathcal{V}$ and some constant $C \ge 1$, it holds for $\ell \in \mathbb{N}, \epsilon \in [\delta, 1]$,

$$(2\pi i\hbar)^{-\frac{d}{2}} \int e^{i\frac{\Phi(x,z)}{\hbar}} a(x,z) \mathrm{d}x = \mathcal{O}_{S^{\delta}} \left(\left(\frac{\hbar}{\delta \epsilon}\right)^{\ell} \right) \quad \text{uniformly on } \{ |\mathrm{f}(z)| > \epsilon \}.$$

This follows directly by splitting the integral

$$(2\pi i\hbar)^{-\frac{d}{2}} \left\{ \int e^{i\frac{\Phi(x,z)}{\hbar}} a(x,z) \chi_{\epsilon}(\mathbf{g}(x,z)) \mathrm{d}x + \int e^{i\frac{\Phi(x,z)}{\hbar}} a(x,z) \left\{ 1 - \chi_{\epsilon}(\mathbf{g}(x,z)) \right\} \mathrm{d}x \right\}$$

where $\chi : \mathbb{R} \to [0,1]$ is a smooth cutoff supported in B_1 and equals to 1 on a neighborhood of 0. In particular, since $\epsilon \geq \delta$, both symbols are in S^{δ} and the first integral satisfies (A.4) with $\mathcal{A} = \{|g(x,z)| \leq \epsilon\}$. Moreover, since g is not degenerate,

$$\{z \in \mho : |f(z)| > \epsilon\} \subset \{z \in \mho : \operatorname{dist}((x_z, z), \mathcal{A}) \ge \epsilon\}$$

if C is sufficiently large. By Lemma A.2, the second integral is also $\mathcal{O}((\frac{\hbar}{\delta\epsilon})^{\infty})$.

In the case of the "standard" phase $\Phi(x,\xi)=x\cdot\xi$, which is relevant for pseudo-differential calculus, we can also perform a stationary phase even in the case where the scalings in x,ξ are different and the amplitude does not necessarily belong to $S^{\delta}(\mathbb{R}^{2n})$ with $\delta\geq\hbar^{\frac{1}{2}}$. Adapting the proof of Proposition A.4, we obtain the following statement.

Proposition A.6. Let $a: \mathbb{R}^{2n+m} \to \mathbb{R}$ with $a \in S_x^{\varepsilon_1} \times S_\xi^{\varepsilon_2} \times S_z^{\eta}$ with $\varepsilon_1, \varepsilon_2, \eta \in [\hbar, 1]$ and assume that $\varepsilon_1 \varepsilon_2 \geq \hbar$. Then for every $\ell \in \mathbb{N}_0$, as $\hbar \to 0$,

$$\frac{1}{(2\pi i\hbar)^n} \int e^{i\frac{x\cdot\xi}{\hbar}} a(x,\xi,z) dx d\xi = \sum_{0 \le k < \ell} \frac{1}{k!} (i\hbar\partial_x \cdot \partial_\xi)^k a(x,\xi,z) \big|_{x=\xi=0} + \mathcal{O}_{S^\eta} \big(\big(\frac{\hbar}{\varepsilon_1 \varepsilon_2}\big)^\ell \big). \tag{A.9}$$

Proof. First, we record from the proof of Proposition A.4 that if the phase is (independent of z) and given by $\Phi: X \mapsto \frac{X \cdot HX}{2}$ where H is a constant non-degenerate matrix (det $H \neq 0$), then the operator

$$L^{k} := i^{-\alpha} (\det H)^{-1/2} \frac{(i\nabla \cdot H\nabla)^{k}}{k!}, \quad \text{for } k \in \mathbb{N}_{0}.$$

Moreover, in the special case $H = \begin{pmatrix} 0 & I_n \\ I_n & 0 \end{pmatrix}$ and $X = (x, \xi)$ for $n \in \mathbb{N}$ (so that $\Phi(X) = x \cdot \xi$, if $\varepsilon_1, \varepsilon_2 \geq \sqrt{\hbar}$, then the expansion (A.9) follows directly from Proposition A.4. Note that by differentiating under the integral, since $a \in S_z^{\eta}$, the error term is controlled in the class S_z^{η} .

We deal with the general case by a change of variable. Without loss of generality, we assume that $\varepsilon_1 \geq \varepsilon_2$ and that the symbol a is independent of the parameter z. Let $(\gamma, \delta) = (\sqrt{\varepsilon_2/\varepsilon_1}, \sqrt{\varepsilon_1\varepsilon_2})$, let $\chi \in \mathscr{C}_c^{\infty}(\mathbb{R}^n, [0, 1])$ be such that $\chi = 1$ on B_n and, let $\theta^k(\xi) = \frac{1-\chi(\xi)}{|\xi|^{2k}} \xi^{\otimes k}$ for $\ell \in \mathbb{N}_0$. Note that $\gamma, \delta \in [\sqrt{h}, 1]$ and let $\chi_{\gamma} = \chi(\cdot/\gamma)$ and $\theta_{\gamma}^k = \theta^k(\cdot/\gamma)$. We split

$$\int e^{i\frac{x\cdot\xi}{\hbar}}a(x,\xi)\mathrm{d}x\mathrm{d}\xi = \int e^{i\frac{x\cdot\xi}{\hbar}}a(x,\xi)\chi_{\gamma}(\xi)\mathrm{d}x\mathrm{d}\xi + \int e^{i\frac{x\cdot\xi}{\hbar}}a(x,\xi)\theta_{\gamma}^{0}(\xi)\mathrm{d}x\mathrm{d}\xi.$$

We can perform repeated integration by parts in the second integral (writing $\frac{-i\hbar\xi\cdot\partial_x}{|\xi^2|}e^{i\frac{x\cdot\xi}{\hbar}}=e^{i\frac{x\cdot\xi}{\hbar}}$), we obtain for any $k\in\mathbb{N}$,

$$\int e^{i\frac{x\cdot\xi}{\hbar}}a(x,\xi)\mathrm{d}x\mathrm{d}\xi = \int e^{i\frac{x\cdot\xi}{\hbar}}a(x,\xi)\chi_{\gamma}(\xi)\mathrm{d}x\mathrm{d}\xi + \frac{(i\hbar)^k}{(\varepsilon_1\gamma)^k}\int e^{i\frac{x\cdot\xi}{\hbar}}a_k(x,\xi)\cdot\theta_{\gamma}^k(\xi)\mathrm{d}x\mathrm{d}\xi.$$

where $a_k = \varepsilon_1^k \partial_x^k a$ is in S^{ε_1} and $\varepsilon_1 \gamma = \delta$. Since a_k is uniformly bounded and compactly supported and $\|\theta_{\gamma}^k(\xi)\|_{L^1} = \mathcal{O}_k(1)$ if k > n, this shows that

$$\int e^{i\frac{x\cdot\xi}{\hbar}}a(x,\xi)\mathrm{d}x\mathrm{d}\xi = \int e^{i\frac{x\cdot\xi}{\hbar}}a(x,\xi)\chi_{\gamma}(\xi)\mathrm{d}x\mathrm{d}\xi + \mathcal{O}(\hbar^{\infty}).$$

Then, by rescaling (the phase and measure are invariant under this change of variables), letting

$$\tilde{a}(x,\xi) := a(x/\gamma,\xi\gamma)\chi(\xi),$$

we have $\widetilde{a} \in S^{\delta}(\mathbb{R}^{2n})$ (in particular this symbol has a fixed compact support and $\delta \geq \sqrt{\hbar}$) and

$$\frac{1}{(2\pi i\hbar)^n} \int e^{i\frac{x\cdot\xi}{\hbar}} a(x,\xi) dx d\xi = \frac{1}{(2\pi i\hbar)^n} \int e^{i\frac{x\cdot\xi}{\hbar}} \widetilde{a}(x,\xi) dx d\xi + \mathcal{O}(\hbar^{\infty})$$

$$= \sum_{0 \le k \le \ell} \hbar^k L^k \widetilde{a}(x,\xi)|_{x=\xi=0} + \mathcal{O}((\hbar\delta^{-2})^{\ell})$$

by applying Proposition A.4. Since $L^k \widetilde{a} = L^k a = \frac{1}{k!} (i\hbar \partial_x \cdot \partial_\xi)^k a$ in this case, this completes the proof.

More specifically, we also need the following consequence of Proposition A.6.

Corollary A.7. Let $n, m \in \mathbb{N}$. Let $a : \mathbb{R}^{2n+m} \to \mathbb{R}$, bounded with compact support, and assume that $(x,\xi) \mapsto \int a(x,\xi,z) dz$ is in $S_x^{\varepsilon_1} \times S_\xi^{\varepsilon_2}$ for some $\varepsilon_1, \varepsilon_2 \in [\hbar,1]$ with $\varepsilon_1 \varepsilon_2 \geq \hbar$. Then for every $\ell \in \mathbb{N}_0$, as $\hbar \to 0$,

$$\frac{1}{(2\pi i\hbar)^n} \int e^{i\frac{x\cdot\xi}{\hbar}} a(x,\xi,z) dx d\xi dz = \sum_{0 \le k < \ell} \frac{1}{k!} \int (i\hbar\partial_x \cdot \partial_\xi)^k a(x,\xi,z) \big|_{x=\xi=0} dz + \mathcal{O}\Big(\big(\frac{\hbar}{\varepsilon_1 \varepsilon_2}\big)^\ell \Big).$$

Proof. By assumption, we can apply Proposition A.6 to the symbol $(x, \xi) \mapsto \int a(x, \xi, z) dz$. This yields the required expansion. Note that since a is smooth with a fixed compact support, one can differentiate under the integral with respect to z.

Corollary A.8. Let $m, \ell \in \mathbb{N}$ and $\varepsilon_1, \varepsilon_2, \delta \in [\hbar, 1]$ with $\varepsilon_1 \leq \delta$ and $\varepsilon_1 \varepsilon_2 \geq \hbar$. Let $(t, \sigma, z) \mapsto a(t, \sigma, z)$ be in $S_t^{\varepsilon_1}(\mathbb{R}) \times S_{\sigma}^{\varepsilon_2}(\mathbb{R})$ and supported in $\{z \in \mathcal{V}\}$ with $\mathcal{V} \in \mathbb{R}^m$ open (\mathcal{V}) is allowed to depend on the scales $\varepsilon_1, \varepsilon_2, \delta$. Assume that $a(t, \sigma, z) = \mathcal{O}_{S_*^{\varepsilon_1} \times S_{\sigma}^{\varepsilon_2}}(\left(\frac{\hbar}{\varepsilon \delta}\right)^{\ell})$ uniformly for $t \in [-\delta, \delta], \sigma \in \mathbb{R}, z \in \mathcal{V}$, then as $\hbar \to 0$,

$$\frac{1}{2\pi i\hbar} \int e^{i\frac{t\sigma}{\hbar}} a(t,\sigma,z) dt d\sigma dz = \mathcal{O}\left(|\mho| \left(\frac{\hbar}{\varepsilon\delta}\right)^{\ell}\right), \qquad \varepsilon = \min(\varepsilon_1, \varepsilon_2).$$

Let $g : \mathbb{R}^n \to \mathbb{R}$, C^1 , such that $\{|g| \leq 1\}$, $\partial_x g \neq 0$ on $\{g = 0\}$ is compact, and $a \in S_t^{\varepsilon_1}(\mathbb{R}) \times S_{\sigma}^{\varepsilon_2}(\mathbb{R})$ is supported in $\{|g| \leq 1\}$. In particular, if it holds for $\ell \geq 2$ and $\delta \in [\varepsilon_1, 1]$

$$a(t,\sigma,z) = \mathcal{O}_{S_t^{\varepsilon_1} \times S_\sigma^{\varepsilon_2}} \left(\left(\frac{\hbar}{\varepsilon \delta} \right)^{\ell} \right) \text{ uniformly for } t \in [-\delta,\delta], \sigma \in \mathbb{R}, z \in \{ |\mathbf{g}| \geq \delta \}$$

then

$$\frac{1}{2\pi i\hbar} \int e^{i\frac{t\sigma}{\hbar}} a(t,\sigma,z) dt d\sigma dz = \mathcal{O}((\frac{\hbar}{\varepsilon\varepsilon_1})^{\ell}).$$

Proof. We simply introduce a smooth cutoff $\chi: \mathbb{R} \to [0,1]$ such that $\chi = 0$ on $\mathbb{R} \setminus [-1,1]$ and $\chi^{\dagger} = 1 - \chi$ satisfies $\chi^{\dagger} = 0$ on $[-\frac{1}{2},\frac{1}{2}]$. Then, by repeated integrations by part, it holds for $z \in \mathcal{V}$,

$$\int e^{i\frac{t\sigma}{\hbar}} a(t,\sigma,z) dt d\sigma = \int e^{i\frac{t\sigma}{\hbar}} \chi(t/\delta) a(t,\sigma,z) dt d\sigma + \left(\frac{i\hbar}{\eta\delta}\right)^k \int e^{i\frac{t\sigma}{\hbar}} \frac{\chi^{\dagger}(t/\delta)}{(t/\delta)^k} \eta^k \partial_{\sigma}^k a(t,\sigma,z) dt d\sigma$$

By assumption, both symbols $(t, \sigma) \mapsto \chi(t/\delta)a(t, \sigma, z)$ and $(t, \sigma) \mapsto \frac{\chi^{\dagger}(t/\delta)}{(t/\delta)^k}\eta^k \partial_{\sigma}^k a(t, \sigma, z)$ are $\mathcal{O}_{S_t^{\varepsilon_1} \times S_{\sigma}^{\varepsilon_2}}(\left(\frac{\hbar}{\varepsilon \delta}\right)^k)$ uniformly for $z \in \mathcal{V}$, hence Proposition A.6 shows that the first integral is $\mathcal{O}(\hbar(\frac{\hbar}{\varepsilon \delta})^k)$ and the second is $\mathcal{O}(\hbar)$ uniformly for $z \in \mathcal{V}$. Then, by integrating over $z \in \mathcal{V}$, this completes the proof.

References

- [1] L. Bollmann and P. Müller. Widom's formula for discontinuous matrix-valued symbols. (arXiv:2311.06036), Nov. 2023.
- [2] P. Bourgade, P. Lopatto, and O. Zeitouni. Optimal rigidity and maximum of the characteristic polynomial of Wigner matrices. (arXiv:2312.13335), Dec. 2023.
- [3] P. Bourgade, K. Mody, and M. Pain. Optimal local law and central limit theorem for β -ensembles. Communications in Mathematical Physics, 390(3):1017–1079, 2022.
- [4] J. Boursier. Optimal local laws and CLT for the circular Riesz gas. (arXiv:2112.05881), Feb. 2023.
- [5] P. Calabrese and J. Cardy. Entanglement entropy and quantum field theory. *Journal of Statistical Mechanics: Theory and Experiment*, 2004(06):P06002, June 2004.
- [6] P. Calabrese, P. Le Doussal, and S. N. Majumdar. Random matrices and entanglement entropy of trapped Fermi gases. *Physical Review A*, 91(1):012303, Jan. 2015.
- [7] P. Calabrese, M. Mintchev, and E. Vicari. Exact relations between particle fluctuations and entanglement in Fermi gases. *EPL (Europhysics letters)*, 98(2):20003, 2012.

- [8] C. Callan and F. Wilczek. On geometric entropy. Physics Letters B, 333(1):55–61, July 1994.
- [9] L. Charles and B. Estienne. Entanglement entropy and Berezin-Toeplitz operators. *Communications in Mathematical Physics*, Oct. 2018.
- [10] C. Charlier. Asymptotics of Hankel determinants with a one-cut regular potential and Fisher–Hartwig singularities. *International Mathematics Research Notices*, 2019(24):7515–7576, 2019.
- [11] J. Chazarain. Spectre d'un hamiltonien quantique et mécanique classique. Communications in Partial Differential Equations, 5(6):595–644, 1980.
- [12] T. Claeys, B. Fahs, G. Lambert, and C. Webb. How much can the eigenvalues of a random Hermitian matrix fluctuate? *Duke Mathematical Journal*, 170(9):2085–2235, 2021.
- [13] O. Costin and J. L. Lebowitz. Gaussian Fluctuation in Random Matrices. *Physical Review Letters*, 75(1):69–72, July 1995.
- [14] D. S. Dean, P. Le Doussal, S. N. Majumdar, and G. Schehr. Noninteracting fermions at finite temperature in a d-dimensional trap: Universal correlations. *Physical Review A*, 94(6):063622, 2016.
- [15] P. Deift, A. Its, and I. Krasovsky. Asymptotics of Toeplitz, Hankel, and Toeplitz + Hankel determinants with Fisher-Hartwig singularities. *Annals of mathematics*, pages 1243–1299, 2011.
- [16] P. Deift, A. Its, and I. Krasovsky. Toeplitz matrices and toeplitz determinants under the impetus of the ising model: Some history and some recent results. *Communications on Pure and Applied Mathematics*, 66(9):1360–1438, 2013.
- [17] P. Deift and X. Zhou. A steepest descent method for oscillatory Riemann–Hilbert problems. Asymptotics for the MKdV equation. *Annals of Mathematics*, 137(2):295–368, 1993.
- [18] A. Deleporte and G. Lambert. Central limit theorem for smooth statistics of one-dimensional free fermions. (arXiv:2304.12275), May 2023.
- [19] A. Deleporte and G. Lambert. Universality for free fermions and the local Weyl law for semiclassical Schrödinger operators. J. Euro. Math. Soc., 2024.
- [20] M. Dimassi and J. Sjöstrand. Spectral Asymptotics in the Semi-Classical Limit. Number 268. Cambridge university press, 1999.
- [21] J. Dubail, J.-M. Stéphan, J. Viti, and P. Calabrese. Conformal field theory for inhomogeneous onedimensional quantum systems: The example of non-interacting Fermi gases. SciPost Physics, 2(1):002, 2017.
- [22] B. Fahs. Uniform Asymptotics of Toeplitz Determinants with Fisher-Hartwig Singularities. Communications in Mathematical Physics, 383(2):685–730, Apr. 2021.
- [23] F. Finster, M. Lottner, and A. Sobolev. The Fermionic Entanglement Entropy and Area Law for the Relativistic Dirac Vacuum State. arXiv preprint arXiv:2310.03493, 2023.
- [24] M. E. Fisher and R. E. Hartwig. Toeplitz Determinants: Some Applications, Theorems, and Conjectures. In K. E. Shuler, editor, *Advances in Chemical Physics*, volume 15, pages 333–353. Wiley, 1 edition, Jan. 1969.
- [25] S. Fournais and S. Mikkelsen. An optimal semiclassical bound on commutators of spectral projections with position and momentum operators. *Letters in Mathematical Physics*, 110(12):3343–3373, 2020.

- [26] Y. V. Fyodorov, G. A. Hiary, and J. P. Keating. Freezing Transition, Characteristic Polynomials of Random Matrices, and the Riemann Zeta Function. *Physical Review Letters*, 108(17):170601, Apr. 2012.
- [27] D. Gioev. Szegő limit theorem for operators with discontinuous symbols and applications to entanglement entropy. *International Mathematics Research Notices*, 2006, 2006.
- [28] D. Gioev and I. Klich. Entanglement Entropy of Fermions in Any Dimension and the Widom Conjecture. *Physical Review Letters*, 96(10):100503, Mar. 2006.
- [29] J. Gustavsson. Gaussian fluctuations of eigenvalues in the GUE. In *Annales de l'IHP Probabilités et Statistiques*, volume 41, pages 151–178, 2005.
- [30] B. Helffer and D. Robert. Comportement semi-classique du spectre des hamiltoniens quantiques elliptiques. In *Annales de l'institut Fourier*, volume 31, pages 169–223, 1981.
- [31] L. Hörmander. The Analysis of Linear Partial Differential Operators. I. Springer, Berlin, 2003.
- [32] J. B. Hough, M. Krishnapur, Y. Peres, and B. Virág. Determinantal processes and independence. *Probability surveys*, 3:206–229, 2006.
- [33] C. P. Hughes, J. P. Keating, and N. O'Connell. On the characteristic polynomial of a random unitary matrix. *Communications in Mathematical Physics*, 220(2):429–451, 2001.
- [34] B.-Q. Jin and V. E. Korepin. Quantum Spin Chain, Toeplitz Determinants and the Fisher—Hartwig Conjecture. *Journal of Statistical Physics*, 116(1):79–95, Aug. 2004.
- [35] J. Junnila, G. Lambert, and C. Webb. Multiplicative chaos measures from thick points of log-correlated fields. (arXiv:2209.06548), Mar. 2023.
- [36] J. Keating and F. Mezzadri. Random Matrix Theory and Entanglement in Quantum Spin Chains. Communications in Mathematical Physics, 252(1):543–579, Dec. 2004.
- [37] J. P. Keating and F. Mezzadri. Entanglement in quantum spin chains, symmetry classes of random matrices, and conformal field theory. *Physical Review Letters*, 94(5):050501, Feb. 2005.
- [38] V. E. Korepin. Universality of entropy scaling in one dimensional gapless models. *Physical Review Letters*, 92(9):096402, Mar. 2004.
- [39] I. Krasovsky. Correlations of the characteristic polynomials in the Gaussian Unitary Ensemble or a singular Hankel determinant. *Duke Mathematical Journal*, 139(3), 2007.
- [40] G. Lambert and E. Paquette. The law of large numbers for the maximum of almost Gaussian log-correlated fields coming from random matrices. *Probability Theory and Related Fields*, 173:157–209, 2019.
- [41] G. Lambert and E. Paquette. Strong approximation of Gaussian \$\beta\$-ensemble characteristic polynomials: The edge regime and the stochastic Airy function. (arXiv:2009.05003), July 2021.
- [42] G. Lambert and E. Paquette. Strong approximation of Gaussian β ensemble characteristic polynomials: The hyperbolic regime. The Annals of Applied Probability, 33(1):549–612, 2023.
- [43] H. J. Landau and H. Widom. Eigenvalue distribution of time and frequency limiting. *Journal of Mathematical Analysis and Applications*, 77(2):469–481, 1980.

- [44] H. Leschke, A. V. Sobolev, and W. Spitzer. Scaling of Rényi entanglement entropies of the free fermi-gas ground state: A rigorous proof. *Physical Review Letters*, 112(16):160403, Apr. 2014.
- [45] H. Leschke, A. V. Sobolev, and W. Spitzer. Asymptotic Growth of the Local Ground-State Entropy of the Ideal Fermi Gas in a Constant Magnetic Field. *Communications in Mathematical Physics*, 381(2):673–705, Jan. 2021.
- [46] R. Marino, S. N. Majumdar, G. Schehr, and P. Vivo. Number statistics for β -ensembles of random matrices: Applications to trapped fermions at zero temperature. *Physical Review E*, 94(3):032115, Sept. 2016.
- [47] M. L. Mehta. Random Matrices. Number 142 in Pure and Applied Mathematics. Elsevier, 2004.
- [48] K. Mody. The log-Characteristic Polynomial of Generalized Wigner Matrices is Log-Correlated. (arXiv:2302.05956), Jan. 2024.
- [49] F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, B. V. Saunders, H. S. Cohl, and M. A. Mc Clain, eds. NIST Digital Library of Mathematical Functions. dlmf.nist.gov.
- [50] S. O'Rourke. Gaussian Fluctuations of Eigenvalues in Wigner Random Matrices. *Journal of Statistical Physics*, 138(6):1045–1066, Mar. 2010.
- [51] E. Paquette and O. Zeitouni. The extremal landscape for the C\$\beta\$E ensemble. (arXiv:2209.06743), Oct. 2022.
- [52] P. Pfeiffer and W. Spitzer. Entanglement Entropy of Ground States of the Three-Dimensional Ideal Fermi Gas in a Magnetic Field. *Annales Henri Poincaré*, Nov. 2023.
- [53] D. Robert. Autour de l'approximation Semi-Classique. Number 68. Birkhauser, 1987.
- [54] T. Shirai and Y. Takahashi. Random point fields associated with certain Fredholm determinants. I. Fermion, Poisson and boson point processes. *Journal of Functional Analysis*, 205(2):414–463, 2003.
- [55] T. Shirai and Y. Takahashi. Random point fields associated with certain Fredholm determinants. II. Fermion shifts and their ergodic and Gibbs properties. *Annals of Probability*, 31(3):1533–1564, 2003.
- [56] B. Simon. Trace Ideals and Their Applications. Number 120. American Mathematical Soc., 2005.
- [57] N. R. Smith, P. Le Doussal, S. N. Majumdar, and G. Schehr. Counting statistics for noninteracting fermions in a d-dimensional potential. *Physical Review E*, 103(3):L030105, Mar. 2021.
- [58] N. R. Smith, P. Le Doussal, S. N. Majumdar, and G. Schehr. Full counting statistics for interacting trapped fermions. *SciPost Physics*, 11(6):110, 2021.
- [59] N. R. Smith, P. Le Doussal, S. N. Majumdar, and G. Schehr. Counting statistics for noninteracting fermions in a rotating trap. *Physical Review A*, 105(4):043315, Apr. 2022.
- [60] A. V. Sobolev. Pseudo-Differential Operators with Discontinuous Symbols: Widom's Conjecture, volume 222. American Mathematical Soc., 2013.
- [61] A. V. Sobolev. Functions of self-adjoint operators in ideals of compact operators. *Journal of the London Mathematical Society*, 95(1):157–176, 2017.

- [62] A. Soshnikov. Determinantal random point fields. Rossiiskaya Akademiya Nauk. Moskovskoe Matematicheskoe Obshchestvo. Uspekhi Matematicheskikh Nauk, 55(5(335)):107–160, 2000.
- [63] A. Soshnikov. Gaussian Limit for Determinantal Random Point Fields. *The Annals of Probability*, 30(1):171–187, Jan. 2002.
- [64] M. Srednicki. Entropy and area. Physical Review Letters, 71(5):666-669, Aug. 1993.
- [65] S. Torquato. Hyperuniform states of matter. Physics Reports, 745:1–95, June 2018.
- [66] S. Torquato, A. Scardicchio, and C. E. Zachary. Point processes in arbitrary dimension from fermionic gases, random matrix theory, and number theory. *Journal of Statistical Mechanics: Theory and Experiment*, 2008(11):P11019, 2008.
- [67] E. Vicari. Entanglement and particle correlations of Fermi gases in harmonic traps. *Physical Review* A, 85(6):062104, June 2012.
- [68] C. Webb. The characteristic polynomial of a random unitary matrix and Gaussian multiplicative chaos—the L2-phase. *Electron. J. Probab*, 20(104):21, 2015.
- [69] H. Widom. Toeplitz determinants with singular generating functions. American Journal of Mathematics, 95(2):333–383, 1973.
- [70] H. Widom. Asymptotic behavior of block Toeplitz matrices and determinants. Advances in Mathematics, 13(3):284–322, 1974.
- [71] H. Widom. Asymptotic behavior of block Toeplitz matrices and determinants. II. Advances in Mathematics, 21(1):1–29, 1976.
- [72] H. Widom. On a class of integral operators with discontinuous symbol. In *Toeplitz Centennial*, pages 477–500. Springer, 1982.