

MAXIMAL δ -SEPARATED SETS IN SEPARABLE METRIC SPACES AND WEAK FORMS OF CHOICE

MICHAŁ DYBOWSKI, PRZEMYSŁAW GÓRKA, AND PAUL HOWARD

ABSTRACT. We show that the statement “In every separable pseudometric space there is a maximal non-strictly δ -separated set.” implies the axiom of choice for countable families of sets. This gives answers to a question of Dybowski and Górka [2]. We also prove several related results.

1. INTRODUCTION

We will use the following standard notation: **ZF** for Zermelo-Fraenkel set theory, **ZFC** for Zermelo-Fraenkel set theory with the axiom of choice, **ZFA** for **ZF** weakened to permit the existence of atoms and **ZFA** + **AC** for **ZFA** with the addition of the axiom of choice.

All of our positive results about the implications between various consequences of the axiom of choice are theorems of **ZFA** and therefore theorems of the stronger theory **ZF**. Our independence results all hold in the theory **ZF** although many of them are first proved in **ZFA** and then transferred to **ZF** using a transfer theorem of Pincus.

Let $\delta > 0$, the subject of this paper is the deductive strength of the following theorem of **ZFA** + **AC**:

*For every separable pseudometric space (X, d) ,
 X has a maximal non-strictly δ -separated set.*

and three closely related theorems (see Definition 1.2, (6) - (9)). Our abbreviation for this theorem of **ZFA** + **AC** is **PN**(δ) (**P** for ‘pseudometric’ and **N** for ‘non-stricty’).

We begin with the definitions we will require.

Definition 1.1. (Metric space and set theoretic terminology)

- (1) We use ω for the set of natural numbers. That is, $\omega = \{0, 1, 2, \dots\}$.
- (2) $\mathbb{Z}$ denotes the set of integers and $\mathbb{Z}^+ = \{1, 2, 3, \dots\}$ is the set of positive integers.
- (3) A *metric space* is a pair (X, d) where X is a set and $d : X \times X \rightarrow [0, \infty)$ is a function which satisfies: For all x, y and z in X
 - (a) $d(x, y) = 0 \iff x = y$
 - (b) $d(x, y) = d(y, x)$
 - (c) $d(x, z) \leq d(x, y) + d(y, z)$
- (4) A *pseudometric space* is a pair (X, d) satisfying items (3b), (3c) above and for every $x \in X$ $d(x, x) = 0$.
- (5) An *ultrametric space* is a metric space (X, d) such that for every $x, y, z \in X$

$$d(x, z) \leq \max(d(x, y), d(y, z)).$$
- (6) Assume that (X, d) is either a metric space or a pseudometric space, that $Y \subseteq X$ and that $\delta > 0$ is a real number. Then
 - (a) Y is a *strictly δ -separated set* (in (X, d)) if for all distinct points x and y in Y , $d(x, y) > \delta$.

2020 *Mathematics Subject Classification.* 03E25 and 03E35 and 30L99.

Key words and phrases. Axiom of Choice, weak axioms of choice, well-ordered set, Fraenkel–Mostowski (FM) permutation model of ZFA, δ -separated sets, separable metric space.

- (b) Y is a *non-strictly δ -separated set* (in (X, d)) if for all distinct points x and y in Y , $d(x, y) \geq \delta$.

Definition 1.2. (Weak choice axioms)

- (1) **AC** (Form 1 in [5]) is the axiom of choice: For every set X of non-empty sets, there is a function $f : X \rightarrow \bigcup X$ such that $\forall y \in X, f(y) \in y$.
- (2) **CC** (Form 8 in [5]): For every countable set X of non-empty sets, there is a function $f : X \rightarrow \bigcup X$ such that $\forall y \in X, f(y) \in y$.
- (3) **CC**($\mathbb{R}$) (Form 94 in [5]): For every countable set X of non-empty sets of real numbers, there is a function $f : X \rightarrow \bigcup X$ such that $\forall y \in X, f(y) \in y$.
- (4) A relation R on set X is *entire (on X)* if for every $x \in X$ there exists $y \in X$ such that $x R y$.
- (5) **DC** (Form 43 in [5]) is “For every non-empty set X and every entire relation R on X there is a sequence $\{x_i\}_{i=1}^{\infty} \subset X$ such that $x_i R x_{i+1}$ for all $i \in \mathbb{Z}^+$ ”.

Each of the following eight statements has a parameter δ , where δ is a positive real number.

- (6) **PS**(δ): For every separable pseudometric space (X, d) , X has a maximal strictly δ -separated subset.
- (7) **MS**(δ): For every separable metric space (X, d) , X has a maximal strictly δ -separated subset.
- (8) **PN**(δ): For every separable pseudometric space (X, d) , X has a maximal non-strictly δ -separated subset.
- (9) **MN**(δ): For every separable metric space (X, d) , X has a maximal non-strictly δ -separated subset.
- (10) **PSE**(δ): For every separable pseudometric space (X, d) , if $S \subseteq X$ is strictly δ -separated set then there is an $S' \subseteq X$ such that $S \subseteq S'$ and S' is a maximal strictly δ -separated set.
- (11) **MSE**(δ): For every separable metric space (X, d) , if $S \subseteq X$ is strictly δ -separated set then there is an $S' \subseteq X$ such that $S \subseteq S'$ and S' is a maximal strictly δ -separated set.
- (12) **PNE**(δ): For every separable pseudometric space (X, d) , if $S \subseteq X$ is non-strictly δ -separated set then there is an $S' \subseteq X$ such that $S \subseteq S'$ and S' is a maximal non-strictly δ -separated set.
- (13) **MNE**(δ): For every separable metric space (X, d) , if $S \subseteq X$ is non-strictly δ -separated set then there is an $S' \subseteq X$ such that $S \subseteq S'$ and S' is a maximal non-strictly δ -separated set.

In [2] Dybowski and Górka studied **PS**(δ) and **MS**(δ). They proved that several strengthenings of **PS**(δ) are equivalent to **AC** in **ZF**. To be more specific, in their Proposition 4.2 they proved that the following statement is equivalent to **AC**:

$$\begin{aligned} &\text{For every pseudometric space } (X, d), \text{ and every } \delta > 0 \\ &X \text{ has a maximal strictly } \delta\text{-separated set.} \end{aligned} \tag{1}$$

(Note that the “separable” requirement has been removed.)

They also showed that if “pseudometric” is replaced by “metric” or by “ultrametric” in (1) each of the resulting statements is equivalent to **AC**.

Remark 1.3 (ZFA). For all $\delta \in \mathbb{R}^+$, **DC** $\implies$ **PS**(δ) + **PN**(δ) + **MN**(δ) + **MS**(δ).


Proof. The implication **DC** $\implies$ **PS**(δ) was proved before (see [2, Corollary 4.5]). The proof given can be modified to show that **DC** implies **PN**(δ). Since **PN**(δ) implies **MN**(δ) and **PS**(δ) implies **MS**(δ), it follows that **DC** also implies **MN**(δ) and **MS**(δ). $\square$

Finally they ask in Problem 4.6 about what else can be said concerning the strength, in set theory without **AC**, of the statements **PS**(δ) and **MS**(δ).

The purpose of this paper is to provide some answers to this question and to related questions involving items (6) through (13) in Definition 1.2 above. Our results are summarized in the following

diagram. (See the paragraph following the proof of Theorem 2.2 for a description of the abbreviations used in the diagram.)


We use a variation of the notation of [1], namely, if Ψ and Φ are sentences and occur in a box together then they are equivalent in **ZF**. If Ψ implies Φ in **ZF** and we don't know whether or not Φ implies Ψ in **ZF** we use $\Psi \rightarrow \Phi$. Sentence $\Psi \mapsto \Phi$ means that in **ZF**, Ψ implies Φ but Φ does not imply Ψ . $\Psi \dashv \vdash \Phi$ means Ψ does not imply Φ in **ZF** and we don't know whether or not Φ implies Ψ .



The diagram shows that there are several unsolved problems. For example

- (1) Does **PSE** imply **PNE**?
- (2) Does **CC** imply **MN** or **MNE**?
- (3) Does either of **MN** or **MSE** imply **MNE**?
- (4) Is **MN** provable in **ZF**?

We note that if the answer to question (4) is in the affirmative¹ then all of the other questions are answered and the diagram collapses to



¹It is worth to have in mind Theorem 5.8.

2. PRELIMINARY RESULTS

Proposition 2.1 (ZFA). If (X, d) is a separable metric space then $|X| \leq |\mathbb{R}|$.

Proof. For each point $x \in X$ we can define a (unique) sequence $\hat{x}$ of elements of D (the countable dense set) by letting $\hat{x}_n$ be the first element of D which is in $B(x, 1/(n+1))$. It is clear that the function $x \mapsto \hat{x}$ is one to one so $|X| \leq |D^{\mathbb{N}_0}| \leq |\mathbb{N}_0^{\mathbb{N}_0}| = |\mathbb{R}|$. $\square$

Theorem 2.2 (ZFA). Assume Ψ is one of **PS, MS, PN, MN, PSE, MSE, PNE** or **MNE** and δ_1 and δ_2 are positive real numbers. Then $\Psi(\delta_1) \implies \Psi(\delta_2)$.

Proof. We give an outline of the proof for the case that Ψ is **PSE** and leave the remaining similar proofs for the interested reader.

Assume **PSE**(δ_1) and let S be a strictly δ_2 -separated set in the pseudometric space (X, p) . Define a metric p' on X by $p'(x, y) = \frac{\delta_1}{\delta_2} p(x, y)$. Then (X, p') is a pseudometric space and S is a strictly δ_1 -separated set in (X, p') . Using **PSE**(δ_1) there is a maximal strictly δ_1 -separated set S' in (X, p') such that $S \subset S'$ and therefore S' is a maximal strictly δ_2 -separated set in (X, p) . $\square$

Having in mind Theorem 2.2, for the remainder of the paper we use the following notation. If Ψ is one of **PS, MS, PN, MN, PSE, MSE, PNE** or **MNE**, we will frequently write Ψ for any one of the equivalent statements “ $\forall \delta > 0, \Psi(\delta)$ ”, “ $\exists \delta > 0, \Psi(\delta)$ ” or, if δ is a specific positive real number, “ $\Psi(\delta)$ ”.

Proposition 2.3 (ZFA). Let (Ψ_1, Φ_1) be (**PS, PSE**) or (**PN, PNE**) and let (Ψ_2, Φ_2) be (**MS, MSE**) or (**MN, MNE**). Then, $\mathbf{CC} + \Psi_1 \implies \Phi_1 \implies \Psi_1$ and $\mathbf{CC}(\mathbb{R}) + \Psi_2 \implies \Phi_2 \implies \Psi_2$.

Proof. Let us notice that the implications $\Phi_1 \implies \Psi_1$ and $\Phi_2 \implies \Psi_2$ are clear. We shall prove only the implication $\mathbf{CC} + \Psi_1 \implies \Phi_1$ since $\mathbf{CC}(\mathbb{R}) + \Psi_2 \implies \Phi_2$ follows in the similar manner using the fact that all separable metric spaces can be bijectively embedded in $\mathbb{R}$ (see Proposition 2.1). We prove the implication $\mathbf{CC} + \Psi_1 \implies \Phi_1$ for pair $(\Psi_1, \Phi_1) = (\mathbf{PN}, \mathbf{PNE})$ since the remaining case follows in the similar way.

Let us assume **CC** and **PN** hold. Let (X, d) be a pseudometric separable space, $\delta > 0$ and $S \subset X$ be an arbitrary non-strictly δ -separated set. We define $N_\delta(S) = \bigcup_{x \in S} B(x, \delta)$. Let us consider pseudometric space $(X \setminus N_\delta(S), d)$. Under **CC** separability is hereditary² (see [1, Theorem 1.12]) so $X \setminus N_\delta(S)$ is separable. Thus by virtue of **PN** space $X \setminus N_\delta(S)$ contains a maximal non-strictly δ -separable set S_0 . We claim that $S' := S \cup S_0$ is the maximal non-strictly δ -separated set in X containing S .

Indeed, let $x, y \in S'$, $x \neq y$. If $x, y \in S$ or $x, y \in S_0$, then $d(x, y) \geq \delta$. If $x \in S$ and $y \in S_0$, then $y \notin N_\delta(S)$ so $y \notin B(x, \delta)$. It proves that S' is non-strictly δ -separated set.

Let us suppose that S' is not a maximal non-strictly δ -separated set i.e. there exists $x_0 \notin S'$ such that $d(x_0, x) \geq \delta$ for all $x \in S'$. In particular $d(x_0, x) \geq \delta$ for all $x \in S$ so $x_0 \notin N_\delta(S)$. Since $x_0 \in X \setminus N_\delta(S)$, $x_0 \notin S_0$ and S_0 is maximal non-strictly δ -separated set in $X \setminus N_\delta(S)$, then $d(x_0, x) < \delta$ for some $x \in S_0$ and it gives a contradiction. $\square$

Theorem 2.4 (ZFA). Let Ψ be one of **PS, MS, PN, MN, PSE, MSE, PNE** or **MNE**, then $\mathbf{DC} \implies \Psi$.

Proof. Let (Ψ_1, Ψ_2) be one of (**PS, PSE**), (**MS, MSE**), (**PN, PNE**) or (**MN, MNE**). By Remark 1.3 we have $\mathbf{DC} \implies \Psi_1$. Thus by virtue of Proposition 2.3 we get $\mathbf{DC} \implies \mathbf{DC} + \Psi_1 \implies \mathbf{CC} + \Psi_1 \implies \Psi_2$. $\square$

²For the implication $\mathbf{CC}(\mathbb{R}) + \Psi_2 \implies \Phi_2$ we use analogous fact that for spaces contained in $\mathbb{R}$ separability is hereditary under **CC**($\mathbb{R}$).

We now turn to a brief discussion of Fraenkel-Mostowski models and a transfer theorem of Pincus. Fraenkel-Mostowski permutation models of the theory **ZFA** provide a relatively easy way of proving independence results in the theory **ZFA**.³ These independence results usually have the form “There is a permutation model in which $\Phi = \Gamma \wedge \neg\Psi$ holds” where Γ and Ψ are consequences of **AC**. A sentence Φ is *transferable* if there is a meta theorem “If there is a Fraenkel-Mostowski model of **ZFA** in which Φ is true then there is a model of **ZF** in which Φ is true”. We will be using a transfer theorem of Pincus ([9, Theorem 4]). The part of the theorem that we need is

Theorem 2.5. Let Φ be a conjunction of **CC** and any (finite) number of injectively boundable statements. If Φ has a Fraenkel-Mostowski model then Φ has a **ZF** model.

For a definition of *injectively boundable* see [9] or [5, Note 103].

3. STRICT δ -SEPARATION IN SEPARABLE PSEUDOMETRIC SPACES

We first show that **PS** is provable in **ZFA**.

Theorem 3.1 (ZFA). For all $\delta \in \mathbb{R}^+$, **PS**(δ).

Proof. By Theorem 2.2, we only have to prove the Theorem for $\delta = 1$. As in the proof of Theorem 2.4 this proof depends on ideas from [2]. Let (X, p) be a separable pseudometric space with countable dense subset $D = \{d_i : i \in \mathbb{Z}^+\}$ and assume that (X, p) has no finite maximal strictly 1-separated set.

Lemma 3.2. If T is a finite subset of X and $N \in \mathbb{Z}^+$ is such that $\Omega := (d_N, 1/2) \setminus \bigcup_{t \in T} \overline{B}(t, 1) \neq \emptyset$, then there exists $i \in \mathbb{Z}^+$ such that $d_i \in \Omega$.

Proof. The set Ω is open and by our assumption non-empty. Since D is dense in (X, p) , there is an $i \in \mathbb{Z}^+$ such that $d_i \in \Omega$. This completes the proof of the Lemma. $\square$

We shall define by recursion a sequence of pairs of positive integers $(k_i, n_i)_{i=1}^\infty$ so that the set $\{d_{k_i} : i \in \mathbb{Z}^+\}$ is a maximal strictly 1-separated set in (X, p) . We give the construction of the set $(k_i, n_i)_{i=1}^\infty$ and prove by induction that $(k_i, n_i)_{i=1}^\infty$ has the following Properties for $i \in \mathbb{Z}^+$.

- (1) $d_{k_i} \in B(d_{n_i}, \frac{1}{2})$,
- (2) if $j \in \mathbb{Z}^+$ is such that $j < i$, then $n_j < n_i$,
- (3) if $j \in \mathbb{Z}^+$ is such that $j < i$, then $p(d_{k_i}, d_{k_j}) > 1$.

Construction of the sequence $(k_i, n_i)_{i=1}^\infty$:

- $(k_1, n_1) = (1, 1)$. It is clear that Properties (1) through (3) are true when $i = 1$.
- Assume that (k_i, n_i) has been defined for all $i < m$ and that for all $i < m$, Properties (1) through (3) are true. By Property (3) the set $\{d_{k_i} : i < m\}$ is strictly 1-separated, but, by our assumption, is not maximal. Therefore there exists $y \in X$ such that for all $i < m$, $p(y, d_{k_i}) > 1$ and thus there must be an $N \in \mathbb{Z}^+$ such that $y \in B(d_N, 1/2)$. Let

$$n_m = \min \left\{ l : B(d_l, 1/2) \setminus \bigcup_{i=1}^{m-1} \overline{B}(d_{k_i}, 1) \neq \emptyset \right\}. \quad (2)$$

By Lemma 3.2 there is $k \in \mathbb{Z}^+$ such that $d_k \in B(d_{n_m}, 1/2) \setminus \bigcup_{i=1}^{m-1} \overline{B}(d_{k_i}, 1)$. We let

$$k_m = \min \left\{ k : d_k \in B(d_{n_m}, 1/2) \setminus \bigcup_{i=1}^{m-1} \overline{B}(d_{k_i}, 1) \right\}. \quad (3)$$

Assuming that (k_i, n_i) satisfies items (1) through (3) (preceding the Construction) for all $i < m$ we show that (k_i, n_i) satisfies those properties for $i = m$.

Property (1) follows from Equation (3) in the Construction.

³The reader is referred to [7, Chapter 4] for a discussion of permutation models.

For Property (2) we first note that for all $j < m$, $n_j \neq n_m$. Indeed, since $p(d_{n_j}, d_{k_j}) < 1/2$ and $p(d_{n_m}, d_{k_m}) < 1/2$, the equality $n_j = n_m$ would lead to the contradiction

$$p(d_{k_j}, d_{k_m}) \leq p(d_{k_j}, d_{n_j}) + p(d_{n_j}, d_{k_m}) = p(d_{k_j}, d_{n_j}) + p(d_{n_m}, d_{k_m}) < 1.$$

To complete the proof assume that Property (2) is false and let i' be the least positive integer less than m for which $n_m \leq n_{i'}$. Then, since for all $j < m$ $n_j \neq n_m$, we have $n_m < n_{i'}$. We also know that $d_{k_m} \in B(d_{n_m}, 1/2) \setminus \bigcup_{i=1}^{i'-1} \overline{B}(d_{k_i}, 1)$. Therefore, $n_{i'} \leq n_m$ and we get a contradiction.

Property (3) follows from Equation (3) in the Construction.

To complete the proof of Theorem 3.1 we have to argue that $S = \{d_{k_i} : i \in \mathbb{Z}^+\}$ is a maximal strictly 1-separated set. The fact that S is strictly 1-separated follows from Property (3). To prove that S is maximal we assume the contrary. Then there exists $y \in X \setminus S$ such that for all $i \in \mathbb{Z}^+$, $p(y, d_{k_i}) > 1$. There is an $n \in \mathbb{Z}^+$ such that $y \in B(d_n, 1/2)$ and we assume that n is the least such positive integer. Since $\{n_i\}_{i \in \mathbb{Z}^+}$ is a strictly increasing sequence of integers there is the least $j \in \mathbb{Z}^+$ such that $n \leq n_j$. It is not possible that $n = n_j$ because, as above this leads to the contradiction

$$p(d_{k_j}, y) \leq p(d_{k_j}, d_{n_j}) + p(d_{n_j}, y) = p(d_{k_j}, d_{n_j}) + p(d_n, y) < 1$$

so $n < n_j$. On the other hand $y \in B(d_n, 1/2) \setminus \bigcup_{i=1}^{j-1} \overline{B}(d_{k_i}, 1)$ and thus by the definition of n_j in (2) we have $n_j \leq n$ which is an obvious contradiction. $\square$

Theorem 3.3 (ZFA). PSE $\implies$ CC.

Proof. By Theorem 2.2, it suffices to prove **PSE**(1) $\implies$ **CC**.

Assume that $\mathcal{Y} = \{Y_n : n \in \omega\}$ is a countable pairwise disjoint family of non-empty sets (and that the mapping $n \mapsto Y_n$ is one to one). Our plan is to apply **PSE**(1) to a certain separable pseudometric space (X, d) and a strictly 1-separated set in (X, d) to get a maximal strictly 1-separated set. From this we will be able to obtain a choice function for $\mathcal{Y}$.

To construct X we assume without loss of generality that

$$\left(\mathbb{R}^{\mathbb{Z}^+} \times \omega\right) \cap \left(\bigcup \mathcal{Y}\right) = \emptyset. \quad (4)$$

For $i \in \mathbb{Z}^+$ we define two sequences $a_i = (a_i(k))_{k \in \mathbb{Z}^+}$ and $a'_i = (a'_i(k))_{k \in \mathbb{Z}^+}$ (both in $\mathbb{R}^{\mathbb{Z}^+}$) by

$$a_i(k) = \begin{cases} \frac{i+1}{i} & \text{if } i = k \\ 0 & \text{otherwise} \end{cases}, \quad a'_i(k) = \begin{cases} \frac{1}{i} & \text{if } i = k \\ 0 & \text{otherwise} \end{cases}.$$

Let $\mathbf{z} \in \mathbb{R}^{\mathbb{Z}^+}$ be the zero sequence defined by $\mathbf{z}(k) = 0$ for all $k \in \mathbb{Z}^+$ and define A and A_E respectively by

$$A = \{a_i : i \in \mathbb{Z}^+\} \cup \{a'_i : i \in \mathbb{Z}^+\}, \quad A_E = \{a_i : i \in \mathbb{Z}^+\} \cup \{a'_i : i \in \mathbb{Z}^+\} \cup \{\mathbf{z}\}.$$

Let d_E be the usual Euclidean metric on A_E defined by

$$d_E(b, c) = \sqrt{\sum_{k=1}^{\infty} (c(k) - b(k))^2}.$$

For each $n \in \omega$, let $X_n = (A \times \{n\}) \cup Y_n$ and let $X = \bigcup_{n \in \omega} X_n$. Finally, define a metric d on X by

$$d(t, s) = \begin{cases} 3 & \text{if } \exists m, n \in \omega \text{ with } m \neq n \wedge t \in X_n \wedge s \in X_m \\ 0 & \text{if } \exists n \in \omega \text{ such that } t, s \in Y_n \\ d_E(x, y) & \text{if } \exists n \in \omega \exists x, y \in A \text{ such that } t = (x, n) \wedge s = (y, n) \\ d_E(x, \mathbf{z}) & \text{if } \exists n \in \omega \exists x \in A \text{ such that } (t = (x, n) \wedge s \in Y_n) \text{ or } \\ & (s = (x, n) \wedge t \in Y_n) \end{cases}$$

Our assumption that the elements of $\mathcal{Y}$ are pairwise disjoint and the assumption (4) ensure that d is well defined. Furthermore, one can easily convince oneself that (X, d) is a pseudometric space.

We now list some properties of (X, d) which will be useful in proving Lemmas 3.5 through 3.7. The proofs are straightforward.

Lemma 3.4.

- (1) For all $n \in \omega, i \in \mathbb{Z}^+, s \in Y_n, d((a'_i, n), s) = 1/i$.
- (2) If $i \neq j, d((a_i, n), (a_j, n)) = \sqrt{\left(\frac{i+1}{i}\right)^2 + \left(\frac{j+1}{j}\right)^2} > 1$.
- (3) For all $n \in \omega, i \in \mathbb{Z}^+, d((a_i, n), (a'_i, n)) = 1$.
- (4) For all $n \in \omega, s \in Y_n, t \in X \setminus Y_n$, if $d(s, t) \leq 1$ then $t = (a'_i, n)$ for some $i \in \mathbb{Z}^+$.

Lemma 3.5. The set $D = \bigcup_{n \in \omega} (A \times \{n\}) = A \times \omega$ is a countable dense subset of (X, d) .

Proof. Since A is countable, $D = A \times \omega$ is countable. Furthermore, since $X = D \cup (\bigcup_{n \in \omega} Y_n)$, to show that D is dense in (X, d) it suffices to prove that every $t \in \bigcup_{n \in \omega} Y_n$ is a limit point of D . By definition $\{(a'_i, n) : i \in \mathbb{Z}^+ \wedge n \in \omega\} \subseteq D$. Therefore part (1) of Lemma 3.4 implies t is a limit point of D . $\square$

Let $S = \bigcup_{n \in \omega} (\{a_i : i \in \mathbb{Z}^+\} \times \{n\}) = \{a_i : i \in \mathbb{Z}^+\} \times \omega$.

Lemma 3.6. The set S is strictly 1-separated.

Proof. Assume (a_i, n) and (a_j, m) are distinct elements in S . If $n \neq m$ then by the first clause of the definition of d , $d((a_i, n), (a_j, m)) = 3 > 1$. On the other hand, if $n = m$ then $i \neq j$, so by part (2) of Lemma 3.4, $d((a_i, n), (a_j, m)) > 1$. $\square$

Lemma 3.7. If S' is a maximal strictly 1-separated set containing S then for all $n \in \omega, |S' \cap Y_n| = 1$.

Proof. Assume S' satisfies the hypotheses of the Lemma and assume $n \in \omega$. Clearly, $|S' \cap Y_n| \leq 1$. We prove $|S' \cap Y_n| \geq 1$ by contradiction. Suppose $S' \cap Y_n = \emptyset$. Since S' is maximal, for all $y \in Y_n$ there is a $t \in S' \setminus Y_n$ such that $d(t, y) \leq 1$. By part (4) of Lemma 3.4, $t = (a'_i, n)$ for some $i \in \mathbb{Z}^+$. But by part (3) in Lemma 3.4, $d((a'_i, n), (a_i, n)) = 1$. This is a contradiction since $(a_i, n) \in S \subseteq S'$. $\square$

By Lemmas 3.5 and 3.6 and **PSE**(1), there is a maximal strictly 1-separated set S' containing S . Therefore, by Lemma 3.7, $\mathcal{Y}$ has a choice function. $\square$

4. NON-STRICT δ -SEPARATION IN SEPARABLE PSEUDOMETRIC SPACES

Theorem 4.1 (ZFA). **PN** $\implies$ **CC**.

Proof. As in the proof of Theorem 3.3, it suffices to prove **PN**(1) $\implies$ **CC**. Assume **PN**(1) and let X be a countable family of non-empty, pairwise disjoint sets where the function $n \mapsto X_n$ is one to one from ω onto X . Also assume without loss of generality that $(\bigcup X) \cap \mathbb{R} = \emptyset$. We will construct a choice function for X .

Lemma 4.2. There is a separable pseudometric space (Z, p) such that


- (1) $\bigcup_{n \in \omega} (X_n \times \{0, 1\}) \subseteq Z$
- (2) Every maximal non-strictly 1-separated set S in (Z, p) satisfies for all $n \in \omega$,
 - (a) $0 < |S \cap (X_n \times \{0, 1\})|$
 - (b) $|S \cap (X_n \times \{0\})| \leq 1$ and $|S \cap (X_n \times \{1\})| \leq 1$

Before giving the proof of the Lemma we argue that Lemma 4.2 is sufficient to prove Theorem 4.1 in the case $\delta = 1$.

To construct a choice function f for X let (Z, p) be the pseudometric given by Lemma 4.2. By **PN**(1) there is a maximal non-strictly 1-separated set S in (Z, p) and by Lemma 4.2, for each $n \in \omega$, each of $S \cap (X_n \times \{0\})$ and $S \cap (X_n \times \{1\})$ contains at most one element and one of these two sets is non-empty. We can therefore define $f(X_n)$ to be the element of $x \in X_n$ for which $(x, 0) \in S$ if such an element exists. Otherwise $f(X_n)$ is the element $x \in X_n$ for which $(x, 1) \in S$.

Proof of Lemma 4.2. We begin by defining a certain subset Y_0 of the Euclidean plane $(\mathbb{R}^2, d)$ as follows. Let

$$\begin{aligned} A_0^0 &= \left\{ (x, y) \in \mathbb{R}^2 : x^2 + (y-1)^2 = 1 \text{ and } 0 < x < \sqrt{3}/2 \text{ and } x \text{ is rational and } 0 < y < 1/2 \right\}, \\ A_0^1 &= \left\{ (x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1 \text{ and } 0 < x < \sqrt{3}/2 \text{ and } x \text{ is rational and } 1/2 < y < 1 \right\}, \\ Y_0 &= A_0^0 \cup A_0^1. \end{aligned}$$



Note that both A_0^0 and A_0^1 are countable. For every positive integer n we let $A_n^0 = \{(a, b+2n) : (a, b) \in A_0^0\}$, $A_n^1 = \{(x, y+2n) : (x, y) \in A_0^1\}$ and $Y_n = A_n^0 \cup A_n^1$. Since for every $(x, y) \in Y_0$, we have $0 < y < 1$ we conclude that for every pair $(x, y) \in Y_n$ we have $2n < y < 2n+1$.

Sublemma 4.2.1. Let $n \in \omega$, then for all (a_1, b_1) and (a_2, b_2) in A_n^0 and for all (x_1, y_1) and (x_2, y_2) in A_n^1 we have

$$d((a_1, b_1), (x_1, y_1)) < 1, \quad (5)$$

$$d((a_1, b_1), (a_2, b_2)) < 1 \text{ and } d((x_1, y_1), (x_2, y_2)) < 1, \quad (6)$$

$$d((a_1, b_1), (0, 2n)) < 1 \text{ and } d((x_1, y_1), (0, 2n+1)) < 1, \quad (7)$$

$$d(a_1, b_1), (0, 2n+1) = 1 \text{ and } d((x_1, y_1), (0, 2n)) = 1. \quad (8)$$

Proof. It is enough to prove the Sublemma for $n = 0$. Parts (6), (7) and (8) are clear from the picture. For part (5) an algebraic proof can be given by assuming that $(a_1, b_1) \in A_0^0$ and $(x_1, y_1) \in A_0^1$ and calculating the square of the distance from (a_1, b_1) to (x_1, y_1) . $\square$

For $n \in \omega$, let $Z_n = Y_n \cup (X_n \times \{0, 1\})$ and let $Z = \bigcup_{n \in \omega} Z_n$. We shall define a pseudometric p on Z using the usual metric on $\mathbb{R}^2$ and for each $n \in \omega$, placing one copy of X_n (namely $X_n \times \{0\}$) at the point $(0, 2n)$ and one copy of X_n (namely $X_n \times \{1\}$) at $(0, 2n+1)$. Here are the details. Define a function $c : Z \rightarrow \mathbb{R}^2$ by

$$c(t, s) = \begin{cases} (t, s) & \text{if } (t, s) \in \bigcup_{n \in \omega} Y_n \\ (0, 2n) & \text{if } t \in X_n \text{ and } s = 0 \\ (0, 2n+1) & \text{if } t \in X_n \text{ and } s = 1 \end{cases}$$

Function c assigns each point of Z to a point in the plane and the distance between two elements of Z will be the Euclidean distance between their assignments. That is, define a pseudometric p on Z by

$$p((t_1, s_1), (t_2, s_2)) = d(c(t_1, s_1), c(t_2, s_2)). \quad (9)$$

Sublemma 4.2.2. Assume $z_1, z_2 \in A_n^0$, $w_1, w_2 \in A_n^1$, $u_1, u_2 \in X_n \times \{0\}$ and $v_1, v_2 \in X_n \times \{1\}$, then

- (1) $p(z_1, w_1) < 1$,
- (2) $p(z_1, z_2) < 1$ and $p(w_1, w_2) < 1$,
- (3) $p(z_1, u_1) < 1$ and $p(w_1, v_1) < 1$,
- (4) $p(z_1, v_1) = 1$ and $p(w_1, u_1) = 1$,
- (5) $p(u_1, v_1) = 1$,
- (6) $p(u_1, u_2) = 0$ and $p(v_1, v_2) = 0$.

Proof. Using the definitions of c and p , items (1) through (4) follow from equations (5) through (8) respectively and items (5) and (6) are clear from the definitions of c and p . $\square$

It is clear that (Z, p) is a pseudometric space. It is also the case that $\bigcup_{n \in \omega} Y_n$ is a countable dense subset of Z since every element of $X_n \times \{0\}$ is a limit point of A_n^0 and every element of $X_n \times \{1\}$ is a limit point of A_n^1 . Therefore (Z, p) is a separable pseudometric space satisfying condition (1) of Lemma 4.2.

To argue that condition (2) of Lemma 4.2 holds assume that S is a maximal non-strictly 1-separated set in (Z, p) and that $n \in \omega$. Part (2b) follows from Sublemma 4.2.2 part (6). To argue for part (2a), we begin by showing that for $i \in \{0, 1\}$

$$\forall u \in X_n \times \{i\}, \forall z \in Z \setminus (X_n \times \{i\}) (p(u, z) < 1 \Rightarrow z \in A_n^i). \quad (10)$$

We will argue for $i = 0$ and the case $i = 1$ we leave to the reader. Assume $u \in X_n \times \{0\}$, $z \in Z \setminus (X_n \times \{0\})$ and $p(u, z) < 1$. Under these assumptions $z \notin Z_m$ where $m \in \omega \setminus \{n\}$ because for every element z of Z_m , if r_1 is the second component of $c(z)$ then $2m \leq r_1 \leq 2m + 1$. Whereas, if r_2 is the second component $c(u)$ then $r_2 = 2n$. This implies that $p(u, z) = d(c(u), c(z)) \geq 1$. Since $z \notin \bigcup_{m \in \omega \setminus \{n\}} Z_m$ we have $z \in Z_n \setminus (X_n \times \{0\}) = A_n^0 \cup A_n^1 \cup (X_n \times \{1\})$. By Sublemma 4.2.2, item (4), $z \notin A_n^1$. By Sublemma 4.2.2, item (5), $z \notin X_n \times \{1\}$. Therefore $z \in A_n^0$ completing the proof of (10).

We complete the proof of Lemma 4.2, part (2) by contradiction. Assume that both $S \cap (X_n \times \{0\})$ and $S \cap (X_n \times \{1\})$ are empty. By the first of these assumptions, since S is maximal, for every $z \in X_n \times \{0\}$ there is an element $s \in S$ such that $p(z, s) < 1$. Hence, by equation (10) this means that there is an element $e_0 \in S \cap A_n^0$. Similarly, there is an element $e_1 \in S \cap A_n^1$. But this is a contradiction, since by Sublemma 4.2.2, part (1), $p(e_0, e_1) < 1$. $\square$

Corollary 4.3.

- (i) **CC** $\iff$ **PSE** $\implies$ **MSE**;
- (ii) **PN** $\iff$ **PNE** $\implies$ **CC**.

Proof. With the use of Proposition 2.3 statement (i) follows from Theorems 3.1 and 3.3 and statement (ii) follows from Theorem 4.1. $\square$

5. STRICT AND NON-STRICT δ -SEPARATION IN SEPARABLE METRIC SPACES

We first make several remarks about the consequences of Proposition 2.1 for Fraenkel-Mostowski models of **ZFA**.

Remark 5.1.

- (1) In any Fraenkel-Mostowski model of **ZFA** the real numbers are well orderable and therefore, by Proposition 2.1, every separable metric space is well orderable. It follows from the discussion in [7, Section 4.2] that all of the statements **MSE**, **MNE**, **MS** and **MN** are true in all Fraenkel-Mostowski models.
- (2) The statements **MSE**, **MNE**, **MS** and **MN** are injectively boundable. See [5, Note 103] or [9] for a definition of *injectively boundable*.

- (3) Therefore using Theorem 2.5, if Φ is any consequence of **AC** which is false in some Fraenkel-Mostowski model and whose negation is injectively boundable then none of the statements **MSE**, **MS**, **MNE** or **MN** implies Φ in **ZF**. For most of the sentences Φ which are consequences of **AC** and mentioned in [5], the negation of Φ is injectively boundable. In particular the negations of **CC** and **DC** are injectively boundable. So none of these four statements implies **CC** nor does any one of them imply **DC**. See [10, Theorem 4] or Note 103 of [5] for a more complete discussion of the transfer theorem of Pincus used here.

Theorem 5.2 (ZFA). $\mathbf{CC}(\mathbb{R}) \iff \mathbf{MSE}$.

Proof. $\mathbf{MSE} \implies \mathbf{CC}(\mathbb{R})$: Assume that $\mathcal{Y} = \{Y_n : n \in \omega\}$ is a countable family of pairwise disjoint non-empty subsets of $\mathbb{R}$ and that the mapping $n \mapsto Y_n$ is one-to-one. Since $\mathbb{R}$ is bijective with $(3n, 3n+1)$ for all $n \in \omega$, we can assume that $Y_n \subset (3n, 3n+1)$ for all $n \in \omega$. Let $Y = \bigcup \mathcal{Y}$. Let D_0 be the set of dyadic numbers i.e. $D_0 = \{m/2^n : m \in \mathbb{Z}, n \in \mathbb{Z}^+\}$ and $D_1 = D_0 + 1/3$. Sets D_0 and D_1 are countable and dense in $\mathbb{R}$ and it is easy to check that $D_0 \cap D_1 = \emptyset$. We can assume⁴ that $Y \cap (D_0 \cup D_1) = \emptyset$. Finally, we put $X = Y \cup D_0 \cup D_1$ and define metric d on X in the following manner:

$$d(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 + |x - y| & \text{if } x \neq y \text{ and } x, y \in D_1 \\ 1 + |y - x + 1/3| & \text{if } x \in D_1 \text{ and } y \notin D_1 \\ 1 + |x - y + 1/3| & \text{if } x \notin D_1 \text{ and } y \in D_1 \\ |x - y| & \text{otherwise} \end{cases}.$$

It is easy to check that d is indeed metric and (X, d) is separable metric space with $D_0 \cup D_1$ as the countable and dense set in X . Since D_1 is strictly 1-separated set in X and $X \subset \mathbb{R}$ is separable, we can use the assumed statement so there exists a maximal strictly 1-separated set S' in X such that $D_1 \subset S'$. We shall show that $|Y_n \cap S'| = 1$ for all $n \in \omega$.

Let us notice that for every $x \in D_0$ there exists $y \in D_1$ such that $d(x, y) = 1$. Indeed, it is enough to take $y = x + 1/3 \in D_1$ and then $d(x, y) = 1 + |x - y + 1/3| = 1$. Since S' is strictly 1-separated set and $D_1 \subset S'$, it means that $S' \cap D_0 = \emptyset$.

Let us fix $n \in \omega$. If $x, y \in Y_n \cap S' \subset Y_n \subset (3n, 3n+1)$, then $d(x, y) = |x - y| < 1$. Thus $x = y$ since $x, y \in S'$. Hence $|Y_n \cap S'| \leq 1$.

Let us suppose that $Y_n \cap S' = \emptyset$ for some $n \in \omega$. Let us take $x \in Y_n$ and fix $y \in S'$. Since $S' \cap D_0 = \emptyset$, either $y \in D_1$ or $y \in Y$. If $y \in D_1$, then $y - 1/3 \in D_0$ and $Y_n \cap D_0 = \emptyset$ so $x \neq y - 1/3$. Thus $d(x, y) = 1 + |x - y + 1/3| > 1$. If $y \in Y$, then $y \notin Y_n$ since $Y_n \cap S' = \emptyset$ so $d(x, y) > 1$. We have $d(x, y) > 1$ for all $y \in S'$ so it contradicts the maximality of S' . Hence $Y_n \cap S' \neq \emptyset$ for all $n \in \omega$.

$\mathbf{CC}(\mathbb{R}) \implies \mathbf{MSE}$: By Proposition 2.3 we have that $\mathbf{CC}(\mathbb{R}) + \mathbf{MS} \implies \mathbf{MSE}$ but **MS** is true in **ZFA** by Theorem 3.1. $\square$

Corollary 5.3. **MSE** is not provable in **ZF**.

Proof. It follows immediately from Theorem 5.2. $\square$

Proposition 5.4 (ZFA). $\mathbf{MNE} \implies \mathbf{CC}(\mathbb{R})$.

Proof. Assume that $\mathcal{Y} = \{Y_n : n \in \omega\}$ is a countable family of pairwise disjoint non-empty subsets of $\mathbb{R}$ and that the mapping $n \mapsto Y_n$ is one-to-one. Let S^1 be unit circle in $\mathbb{R}^2$, namely $S^1 = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$. There exists a bijection $f : \mathbb{R} \rightarrow S^1$. Let $\hat{Y}_n = f(Y_n) + (4n, 0)$ for all $n \in \omega$. Since f is bijective and $\mathbb{R}$ is linearly ordered, it is enough to prove that there exists set S' such that $S' \cap \hat{Y}_n$ is non-empty and finite for all $n \in \omega$ to complete the proof.

⁴If $Y_n \cap D_i \neq \emptyset$ for some $n \in \omega$ and $i = 0, 1$, then we can take $j_0 = \min \{j \in \mathbb{Z}^+ : d_j^i \in Y_n \cap D_i\}$ where $D_i = \{d_j^i\}_{j=1}^\infty$. Thus we can put $y_n = d_{j_0}^i \in Y_n$. Hence from those sets Y_n such that $Y_n \cap (D_0 \cup D_1) \neq \emptyset$ we can select elements directly, without any choice.

Let $D^1 = \{(x, y) \in \mathbb{Q}^2: x^2 + y^2 < 1\}$ and $D_n^1 = D^1 + (4n, 0)$ for all $n \in \omega$. We consider metric space (X, d) where $X = \bigcup_{n \in \omega} (\hat{Y}_n \cup D_n^1)$ and d is the usual Euclidean metric on $\mathbb{R}^2$. Obviously, X is separable with $\bigcup_{n \in \omega} D_n^1$ as a countable and dense set. Let $S = \{(4n, 0): n \in \omega\}$. It is non-strictly 1-separated set in X so by **MNE** there exists a maximal non-strictly 1-separated set S' such that $S \subset S'$.

Let us notice that for all $n \in \omega, x \in D_n^1$ and $y \in \hat{Y}_n$ we have $d((4n, 0), x) < 1$ and $d((4n, 0), y) = 1$. Thus $S' \cap \bigcup_{n \in \omega} D_n^1 \setminus \{(4n, 0)\} = \emptyset$. Since S' is maximal and $d(x, y) \geq 1$ for all $x \in D_n^1 \cup \hat{Y}_n$ and $y \in D_m^1 \cup \hat{Y}_m$ where $n \neq m$, it is easy to see that $S' \cap \hat{Y}_n \neq \emptyset$ for all $n \in \omega$. Moreover for all $n \in \omega$ set $S' \cap \hat{Y}_n$ is finite since every 1-separated set contained in $S^1 + (4n, 0)$ is finite. $\square$

Corollary 5.5 (ZFA). $\mathbf{CC}(\mathbb{R}) + \mathbf{MN} \iff \mathbf{MNE}$.

Proof. It follows from the Proposition 5.4 and Proposition 2.3. $\square$

Theorem 5.6 (ZFA). $\mathbf{MN} + \mathbf{CC} \iff \mathbf{PN}$.

Proof. For the implication " $\Leftarrow$ " we note that $\mathbf{PN} \implies \mathbf{MN}$ is clear and from Theorem 4.1 we have $\mathbf{PN} \implies \mathbf{MN}$.

For the other implication let (X, p) be a separable pseudometric space and D be a countable and dense set. We convert this space into metric space in the standard way. We define an equivalence relation $\sim$ on X as follows: $x \sim y \iff p(x, y) = 0$ for all $x, y \in X$. Then $(X/\sim, d)$ is metric space where $d([x], [y]) = p(x, y)$ for all $x, y \in X$ and $[x]$ denotes the equivalence class of x . Obviously $D/\sim$ is a countable and dense set in $X/\sim$ so $(X/\sim, d)$ is separable.

Since we assumed **MN**, there exists a maximal non-strictly 1-separated set S in $(X/\sim, d)$. Every non-strictly 1-separated in every separable metric space is at most countable so S is at most countable. Let $S = \{S_i\}_{i=1}^\infty$. From **CC** there exists set $\hat{S} = \{s_i\}_{i=1}^\infty \subset X$ such that $s_i \in S_i$ for every $i \in \mathbb{Z}^+$. Then $[s_i] = S_i$ for every $i \in \mathbb{Z}^+$ and it is easy to see that $\hat{S}$ is a maximal non-strictly 1-separated set in (X, p) . $\square$

Corollary 5.7. In the theory **ZF**, the sentence **PN** (and therefore the sentence **PNE**) does not imply **DC**.

Proof. In [6] a Fraenkel-Mostowski model is constructed in which **CC** is true and **DC** is false. By Remark 5.1 Part (1), **MN** is also true in this model. By Remark 5.1 Part (2), **MN** is injectively boundable. It follows from Theorem 2.5 that there is a model of **ZF** in which both **CC** and **MN** are true and **DC** is false. Therefore, by Theorem 5.6, in this model **PN** is true and **DC** is false. $\square$

We close this section with the following observation.

Theorem 5.8 (ZFA). Let (X, d) be a separable, complete metric space and let $\delta > 0$. Then, for every non-strictly δ -separated subset S of X there exists a maximal non-strictly δ -separated subset S' of X such that $S \subset S'$.

Proof. First of all we shall prove the following lemma.

Lemma 5.9 (ZFA). Let (X, d) be a separable, complete metric space and $\mathcal{F}$ be the family of all non-empty closed subsets of X . Then, there exists choice function for $\mathcal{F}$ i.e. set $\{a_F\}_{F \in \mathcal{F}} \subset X$ such that $a_F \in F$ for all $F \in \mathcal{F}$.

Proof. Let $\{x_i\}_{i=1}^\infty \subset X$ be a countable and dense subset of X . Let $F \in \mathcal{F}$. We shall give the construction of $a_F \in F$. We define sequence $\{n_k\}_{k=1}^\infty \subset \omega$ in the following manner:

- $n_1 = \min \{i \in \mathbb{Z}^+: x_i \in \bigcup_{x \in F} B(x, 1/2)\},$
- $n_{k+1} = \min \{i \in \mathbb{Z}^+: x_i \in B(x_{n_k}, 1/2^k) \cap \bigcup_{x \in F} B(x, 1/2^{k+1})\}$ for all $k \geq 1$.

Obviously, set $\bigcup_{x \in F} B(x, 1/2)$ is open and non-empty so n_1 is well-defined. Let us assume that n_k are defined for $k \leq m$. We will show that n_{m+1} is well-defined. Since set $B(x_{n_m}, 1/2^m) \cap \bigcup_{x \in F} B(x, 1/2^{m+1})$ is open, it is enough to show that it is non-empty. Let us notice that $x_{n_m} \in \bigcup_{x \in F} B(x, 1/2^m)$ so $B(x_{n_m}, 1/2^m) \cap F \neq \emptyset$. As a consequence $B(x_{n_m}, 1/2^m) \cap \bigcup_{x \in F} B(x, 1/2^{m+1}) \neq \emptyset$ and it shows that n_{m+1} is well-defined. The sequence $\{n_k\}_{k=1}^\infty$ has the following properties:

- (1) $x_{n_{k+1}} \in B(x_{n_k}, 1/2^k)$ for all $k \geq 1$,
- (2) $\text{dist}(x_{n_k}, F) < 1/2^k$ for all $k \geq 1$.

From (1) we conclude that sequence $\{x_{n_k}\}_{k=1}^\infty$ is Cauchy sequence so it converges since X is complete. Let $\lim_{k \rightarrow \infty} x_{n_k} = x \in X$. From (2), since $\text{dist}(x, F) \leq \text{dist}(x_{n_k}, F) + d(x, x_{n_k})$ for every $k \geq 1$, then $\text{dist}(x, F) = 0$. But F is closed so $x \in F$. We put $a_F := x$. $\square$

Let $\{x_i\}_{i=1}^\infty \subset X$ be a countable and dense subset of X and let S be non-strictly δ -separated subset of X . By virtue of Lemma 5.9 let $\mathcal{F}$ be the family of all non-empty closed subset of X and $\{a_F\}_{F \in \mathcal{F}} \subset X$ be a set such that $a_F \in F$ for all $F \in \mathcal{F}$. Let us denote $N_\delta(S) = \bigcup_{x \in S} B(x, \delta)$. If $N_\delta(S) = X$, then S is maximal non-strictly δ -separated set so we assume $N_\delta(S) \neq X$. Let $y \in X \setminus N_\delta(S)$. We define sequences $\{y_n\}_{n=1}^\infty \subset X$ and $\{k_n\}_{n=1}^\infty \subset \omega$ as follows⁵:

- $k_1 = 0, y_1 = y$,
- $k_n = \min \left\{ i > k_{n-1} : \overline{B}(x_i, \delta/3) \setminus \left(\bigcup_{j=1}^{n-1} B(y_j, \delta) \cup N_\delta(S) \right) \neq \emptyset \right\}$ for all $n \geq 2$,
- $y_n = a_F$ where $F = \overline{B}(x_{k_n}, \delta/3) \setminus \left(\bigcup_{j=1}^{n-1} B(y_j, \delta) \cup N_\delta(S) \right)$ for all $n \geq 2$.

We shall show that $S' := S \cup \{y_n\}_{n=1}^\infty$ is a maximal non-strictly δ -separated set in X containing S . Let us fix $n \in \mathbb{Z}^+$. Since $y_n \notin \left(\bigcup_{j=1}^{n-1} B(y_j, \delta) \cup N_\delta(S) \right)$, then $d(y_n, y_j) \geq \delta$ for all $j < n$ and $d(y_n, s) \geq \delta$ for all $s \in S$. But $n \in \mathbb{Z}^+$ was arbitrary so $d(y_n, y_j) \geq \delta$ for all $n, j \in \mathbb{Z}^+, n \neq j$. It proves that S' is non-strictly δ -separated set.

Let us suppose that S' is not maximal i.e. there exists $x \in X$ such that $d(x, y_n) \geq \delta$ for all $n \in \mathbb{Z}^+$ and $d(x, s) \geq \delta$ for all $s \in S$. There exists $i \in \mathbb{Z}^+$ such that $x \in \overline{B}(x_i, \delta/3)$. Thus $x \in \overline{B}(x_i, \delta/3) \setminus \left(\bigcup_{j=1}^\infty B(y_j, \delta) \cup N_\delta(S) \right)$. Let us suppose that $i \neq k_n$ for any $n \in \mathbb{Z}^+$. Sequence $\{k_n\}_{n=1}^\infty$ is increasing so there exists $n \in \mathbb{Z}^+$ such that $k_n < i < k_{n+1}$. By the definition of k_{n+1} we have $\overline{B}(x_i, \delta/3) \setminus \left(\bigcup_{j=1}^n B(y_j, \delta) \cup N_\delta(S) \right) = \emptyset$ so $\overline{B}(x_i, \delta/3) \setminus \left(\bigcup_{j=1}^\infty B(y_j, \delta) \cup N_\delta(S) \right) = \emptyset$ which contradicts the fact that x is an element of this set. Hence $i = k_n$ for some $n \in \mathbb{Z}^+$. Obviously $n \geq 2$ since $i \in \mathbb{Z}^+$ and $k_1 = 0 \notin \mathbb{Z}^+$. It means that $x \in \overline{B}(x_{k_n}, \delta/3) \setminus \left(\bigcup_{j=1}^{n-1} B(y_j, \delta) \cup N_\delta(S) \right)$. Since $y_n \in \overline{B}(x_{k_n}, \delta/3) \setminus \left(\bigcup_{j=1}^{n-1} B(y_j, \delta) \cup N_\delta(S) \right)$ either, we obtain $x, y_n \in \overline{B}(x_{k_n}, \delta/3)$ so $d(x, y_n) \leq 2\delta/3$. It gives a contradiction with the fact that $d(x, y_n) \geq \delta$. Hence S' is maximal non-strictly δ -separated set. $\square$

REFERENCES

- [1] H. L. Bentley and H. Herrlich, *Countable choice and pseudometric spaces*, Topology and its Applications, **85**, 153-164, 1998.
- [2] M. Dybowski and P. Górka, *The axiom of choice in metric measure spaces and maximal δ -separated sets*, Archive for Mathematical Logic **62**, 735-749, 2023.
- [3] P. Górka, *Separability of a metric space is equivalent to the existence of a Borel measure*, The American Mathematical Monthly, **128**, 84-86, 2020.
- [4] H. Herrlich, *Axiom of choice*, Springer, 2006.

⁵It may happen that construction of these sequences stops at some point i.e. we constructed k_n for $n \leq m$ and $\overline{B}(x_i, \delta/3) \setminus \left(\bigcup_{j=1}^m B(y_j, \delta) \cup N_\delta(S) \right) = \emptyset$ for all $i > k_m$. The proof in that case is similar as in the case where construction of k_n goes to infinity.

- [5] P. Howard and J. E. Rubin, *Consequences of the Axiom of Choice*, Mathematical Surveys and Monographs (59), American Mathematical Society, Providence, RI, 1998.
- [6] P. Howard and J. E. Rubin, *The Boolean prime ideal theorem plus countable choice do not imply dependent choice*, MathLogic Quart., **42**, 410-420, 1996.
- [7] T. J. Jech, *The Axiom of Choice*, Studies in Logic and the Foundations of Mathematics **75**, North-Holland Publishing Co., Amsterdam, 1973.
- [8] R. B. Jensen, *Consistency results for **ZF***, Notices Am. Math. Soc. **14**, 137, 1967.
- [9] D. Pincus, *Zermelo-Fraenkel consistency results by Fraenkel-Mostowski methods*, J. Symbolic Logic **37**, 721-743, 1972.
- [10] D. Pincus, *Adding dependent choice*, Ann. Math. Logic **11**, 105-145, 1977.
- [11] E. Tachtsis, *On the existence of almost disjoint and MAD families without AC*, Bulletin Polish Acad. Sci. Math. **67**, 101-124, 2019.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

FACULTY OF MATHEMATICS AND INFORMATION SCIENCES, WARSAW UNIVERSITY OF TECHNOLOGY, PL. POLITECHNIKI
1, 00-661 WARSAW, POLAND
Email address: `michal.dybowski.dokt@pw.edu.pl`

FACULTY OF MATHEMATICS AND INFORMATION SCIENCES, WARSAW UNIVERSITY OF TECHNOLOGY, PL. POLITECHNIKI
1, 00-661 WARSAW, POLAND
Email address: `przemyslaw.gorka@pw.edu.pl`

2770 EMBER WAY, ANN ARBOR, MI 48104, USA
Email address: `phoward@umich.edu`