

SPECTRAL ENERGY OF DIRAC OPERATORS FOR FERMION SCATTERING ON TOPOLOGICAL SOLITONS IN THE NONLINEAR $O(3)$ σ -MODEL

DAIJU FUNAKAWA, SATOSHI OKUMURA AND YUKI UEDA

ABSTRACT. We present a criteria affirming the existence of discrete ground states of the Dirac operators which describe the fermion scattering on topological solitons in the nonlinear $O(3)$ σ -model. Additionally, we investigate a sufficient condition that elucidates the phenomenon of supersymmetry breaking within the supersymmetric quantum mechanics associated with this Dirac operator.

1. INTRODUCTION

Topological solitons of nonlinear field models play an important role in field theory, condensed matter physics and hydrodynamics etc (see e.g. [7, 8]). In this paper, we study the Dirac operators which describe fermion scattering on topological solitons in the nonlinear $O(3)$ σ -model (see [6] for details). Moreover, we investigate whether the supersymmetric quantum mechanics constructed from this model has supersymmetry breaking or not. As below, we introduce the Dirac operator for fermion scattering on topological solitons in the nonlinear $O(3)$ σ -model.

Let $\sigma_j \in M_2(\mathbb{C})$ be the *Pauli matrices* for $j = 1, 2, 3$, that is,

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

The *spin matrices* are defined by $\gamma_j = (i\sigma_j) \otimes 1_2$ for $j = 1, 2$ and $\gamma_3 = (-\sigma_3) \otimes 1_2$, where 1_d is the $d \times d$ identity matrix. It is easy to verify that

- $\sigma_j^2 = 1_2$ for $j = 1, 2, 3$ and also $\sigma_1\sigma_2 = i\sigma_3$, $\sigma_2\sigma_3 = i\sigma_1$ and $\sigma_3\sigma_1 = i\sigma_2$;
- $\gamma_j^2 = 1_4$ for $j = 1, 2, 3$ and also $\gamma_1\gamma_2 = i\gamma_3$, $\gamma_2\gamma_3 = -i\gamma_1$ and $\gamma_3\gamma_1 = -i\gamma_2$;
- $\{\sigma_j, \sigma_k\} = 0_2$ and $\{\gamma_j, \gamma_k\} = 0_4$ for all $1 \leq j \neq k \leq 3$,

where $\{a, b\} = ab + ba$ and 0_d is the $d \times d$ zero matrix.

Denote by \mathcal{N} the set of all vector fields $\mathbf{n} = (n_1, n_2, n_3)$ such that

- (n-1) n_j is continuously differentiable a.e. \mathbb{R}^2 for $j = 1, 2, 3$;
- (n-2) $\|\mathbf{n}(\mathbf{x})\| = 1$ a.e. $\mathbf{x} \in \mathbb{R}^2$;
- (n-3) $D_k n_j$ is bounded a.e. \mathbb{R}^2 for all $j = 1, 2, 3$ and $k = 1, 2$;
- (n-4) $\lim_{\|\mathbf{x}\| \rightarrow \infty} \|\nabla n_j(\mathbf{x})\| = 0$ for $j = 1, 2, 3$,

where $\|\cdot\|$ is the standard Euclidian distance and D_j is the generalized partial differential operator in the variable x_j , and also $\nabla = (D_1, D_2)$ is a (generalized) gradient. Due to condition (n-2), which asserts that $\sup_{\mathbf{x} \in \mathbb{R}^2} |n_j(\mathbf{x})| \leq 1$, the function n_j can be considered a bounded multiplication operator

2010 *Mathematics Subject Classification*. Primary: 81Q10; Secondary: 47A75, 47B25, 81Q60,

Key words and phrases. Dirac operators, fermion scattering on topological solitons, nonlinear $O(3)$ σ -model, ground states, supersymmetry breaking.

on $L^2(\mathbb{R}^2)$. Additionally, in accordance with condition (n-3), the operator $D_k n_j$ can also be viewed as a bounded multiplication operator on $L^2(\mathbb{R}^2)$ for $j = 1, 2, 3$ and $k = 1, 2$.

Definition 1.1 ([6]). *A Dirac operator for fermion scattering on topological solitons in the nonlinear $O(3)$ σ -model (for short, fermionic Dirac operator) is a linear operator on $L^2(\mathbb{R}^2; \mathbb{C}^4) \otimes \mathbb{C}^2$ defined by*

$$H_{\text{ferm}} \equiv H_{\text{ferm}}(\mathbf{n}) := -i \sum_{k=1}^2 \gamma_3 \gamma_k D_k \otimes 1_2 + m \sum_{j=1}^3 \gamma_3 n_j \otimes \tau_j$$

for Yukawa coupling constant $m > 0$, $\mathbf{n} = (n_1, n_2, n_3) \in \mathcal{N}$ and $\boldsymbol{\tau} \equiv (\tau_1, \tau_2, \tau_3) := (\sigma_1, \sigma_2, \sigma_3)$.

By the natural identification $L^2(\mathbb{R}^2; \mathbb{C}^4) \otimes \mathbb{C}^2 \cong \oplus_{i=1}^2 L^2(\mathbb{R}^2; \mathbb{C}^2) \otimes \mathbb{C}^2$, one can see that H_{ferm} has the matrix representation $H_{\text{ferm}} = H \oplus H$, where H is a linear operator on $L^2(\mathbb{R}^2; \mathbb{C}^2) \otimes \mathbb{C}^2$ as

$$(1.1) \quad H \equiv H(\mathbf{n}) := (-i\sigma_2 D_1 + i\sigma_1 D_2) \otimes 1_2 - m \sum_{j=1}^3 \sigma_3 n_j \otimes \tau_j.$$

Since H_{ferm} is diagonal, we have $\sigma(H_{\text{ferm}}) = \sigma(H)$. In the following, we use H as the fermionic Dirac operator instead of H_{ferm} .

Denote by $\mathcal{C} = \{(r, \varphi) : r > 0, 0 \leq \varphi < 2\pi\}$ the polar coordinate system and $\Phi : \mathcal{C} \rightarrow \mathbb{R}^2$ the polar coordinate conversion. A unitary operator $U : L^2(\mathbb{R}^2) \rightarrow L^2(\mathcal{C}, r dr d\varphi)$ defined by $Uf = f \circ \Phi$ plays a role of the polar coordinate conversion for operators. Then we get the equivalence of operators:

$$H(\mathbf{n}) \simeq \tilde{H}(\mathbf{n}) := (-i\sigma_2 \tilde{D}_1 + i\sigma_1 \tilde{D}_2) \otimes 1_2 - m \sum_{j=1}^3 \sigma_3 \tilde{n}_j \otimes \tau_j$$

with respect to the unitary conjugation by $U \otimes 1_2$, where $\tilde{n}_j = U n_j U^{-1}$ for $j = 1, 2, 3$ and

$$\tilde{D}_1 = \cos \varphi \cdot \partial_r - \frac{\sin \varphi}{r} \cdot \partial_\varphi, \quad \text{and} \quad \tilde{D}_2 = \sin \varphi \cdot \partial_r + \frac{\cos \varphi}{r} \cdot \partial_\varphi,$$

see Sections 3.1 and 3.2 for more details. By the unitary invariance of spectrums, we have $\sigma(H(\mathbf{n})) = \sigma(\tilde{H}(\mathbf{n}))$, $\sigma_{\text{d}}(H(\mathbf{n})) = \sigma_{\text{d}}(\tilde{H}(\mathbf{n}))$ and $\sigma_{\text{ess}}(H(\mathbf{n})) = \sigma_{\text{ess}}(\tilde{H}(\mathbf{n}))$, where $\sigma_{\text{d}}(A)$ and $\sigma_{\text{ess}}(A) := \sigma(A) \setminus \sigma_{\text{d}}(A)$ are *discrete* and *essential spectrums* of an operator A , respectively. Especially, we are interested in the Dirac operator $H(\mathbf{n}) \simeq \tilde{H}(\mathbf{n})$ when a vector field \mathbf{n} is the *hedgehog-ansatz* $\mathbf{n}(F)$, that is, for $\omega = (r, \varphi) \in \mathcal{C}$,

$$(1.2) \quad \tilde{n}_1(\omega) = \sin F(r) \cos N\varphi, \quad \tilde{n}_2(\omega) = \sin F(r) \sin N\varphi, \quad \tilde{n}_3(\omega) = \cos F(r),$$

where $N \in \mathbb{Z}$ and $F : (0, \infty) \rightarrow [0, \pi]$ is a function that is continuously differentiable and subjects to specific boundary conditions. The function F is called a *profile function*.

In nuclear physics and quantum field theory, the notion of *ground states* is significantly important in elucidating the lowest energy of the Hamiltonian (Dirac operator) in a quantum mechanical system. Rigorously, a ground state of a self-adjoint operator T bounded from below is defined as a nonzero eigenvector in $\ker(T - E_0(T))$, where $E_0(T) := \inf \sigma(T)$ denotes the lowest energy of T . Notably, as $H(\mathbf{n})^2$ is positive, its lowest energy $E_0(H(\mathbf{n})^2)$ is grater than or equal to zero. In Proposition 2.4, we prove that $\sigma_{\text{ess}}(H(\mathbf{n})) = (-\infty, -m] \cup [m, \infty)$ holds true for all vector fields $\mathbf{n} \in \mathcal{N}$, consequently implying $\sigma_{\text{ess}}(H(\mathbf{n})^2) = [m^2, \infty)$ and $\sigma_{\text{d}}(H(\mathbf{n})^2) \subset [0, m^2)$. From diverse perspectives such as nuclear physics, quantum mechanics and condensed matter physics, it is concerned whether $E_0(H(\mathbf{n})^2)$ belongs to the discrete spectrum $\sigma_{\text{d}}(H(\mathbf{n})^2)$. As an interest of purely mathematics, the similar type

problem has already studied (see [3, 4]; the existence of discrete ground states of Dirac operators in the chiral quark soliton model). Here, we introduce the notion of discrete ground states.

Definition 1.2. Let T be a self-adjoint operator bounded from below. If $E_0(T) \in \sigma_d(T)$, then a non-zero vector in $\ker(T - E_0(T))$ is called a *discrete ground state of T* . In addition, we denote by $E_0^+(T) = \inf(\sigma(T) \cap [0, \infty))$ and $E_0^-(T) = \sup(\sigma(T) \cap (-\infty, 0])$. If $E_0^+(T) \in \sigma_d(T)$ (resp. $E_0^-(T) \in \sigma_d(T)$), then a non-zero vector in $\ker(T - E_0^+(T))$ (resp. $\ker(T - E_0^-(T))$) is called a *discrete positive (resp. negative) energy ground state of T* .

In this paper, we firstly establish sufficient conditions for the existence of discrete energy ground states for the fermionic Dirac operator $H(\mathbf{n}(F))$ with a hedgehog-ansatz. These conditions provide insights into the stability and nature of the solutions associated with the model. To derive sufficient conditions, our attention is directed towards the reduced component $H(\ell, s, t)$ of the Dirac operator $H(\mathbf{n}(F))$. Specifically, we focus on its action within an eigenspace of the grand spin operator K_3 , characterized by an eigenvalue of the form $\ell + \frac{s}{2} + \frac{Nt}{2}$, where ℓ is an integer, and s and t take values of ± 1 . By scrutinizing the operator $H(\ell, s, t)^2$, our first main theorem is established.

Theorem 1.3. Let us consider a profile function F . If there exist $\ell \in \mathbb{Z}$ and $t \in \{1, -1\}$ such that

$$\inf_{\substack{f \in C_0^\infty((0, \infty)) \\ \|f\|_2 = 1}} \left\langle f, \left(-\frac{d^2}{dr^2} - \frac{1}{r} \frac{d}{dr} + \frac{\ell^2 + (\ell + 1)^2}{2r^2} + mt \frac{dF}{dr} \sin F \right) f \right\rangle_2 < 0,$$

then $H(\mathbf{n}(F))$ has a discrete positive or negative energy ground state, where $C_0^\infty((0, \infty))$ is the set of all compactly supported C^∞ -functions on $(0, \infty)$ and

$$\langle u, v \rangle_2 := \int_{(0, \infty)} \overline{u(r)} v(r) r dr \quad \text{and} \quad \|u\|_2 := \sqrt{\langle u, u \rangle_2}$$

for all $u, v \in L^2((0, \infty), r dr)$.

A strategy of proof is to use the min-max principle as well as [3] and [4]. However, the choice of vectors in the eigenspace of the grand spin operator K_3 is quite different from the previous works, see Lemma 4.1 for details.

The Dirac operator $H = H(\mathbf{n})$ is employed to construct a *supersymmetric quantum mechanics* denoted as a quadruple $\{L^2(\mathbb{R}^2; \mathbb{C}^2) \otimes \mathbb{C}^2, H^2, H, \Gamma\}$, in which Γ is a unitary and selfadjoint operator on $L^2(\mathbb{R}^2; \mathbb{C}^2) \otimes \mathbb{C}^2$ satisfying the condition $\{H, \Gamma\} = 0$. The supersymmetry is a theoretical concept in physics denoting a symmetry between bosons and fermions. The subsequent goal in the paper is to establish a sufficient condition for the supersymmetric quantum mechanics associated with the fermionic Dirac operator H , to exhibit supersymmetry (SUSY) breaking. To elaborate further, the SUSY breaking means the absence of zero-energy states in the spectrum of the Dirac operator H , i.e. $\text{Ker} H = \{0\}$. Actually, we will find a sufficient condition to be $E_0(H^2) > 0$, which implies that $\text{Ker} H = \{0\}$.

Theorem 1.4. Let $\mathbf{n} = \mathbf{n}(F)$ be the hedgehog-ansatz with $\limsup_{r \rightarrow 0^+} \left| \frac{\sin F(r)}{r} \right| < \infty$. If

$$m > \sup_{r > 0} \sqrt{\left| F'(r)^2 - \left(\frac{N}{r} \sin F(r) \right)^2 \right|},$$

then the supersymmetric quantum mechanics $\{L^2(\mathbb{R}^2; \mathbb{C}^2) \otimes \mathbb{C}^2, H^2, H, \Gamma\}$ is SUSY breaking, where Γ is a unitary self-adjoint operator defined in Lemma 5.1.

Note that, if H has no discrete energy ground states, then $\{L^2(\mathbb{R}^2; \mathbb{C}^2) \otimes \mathbb{C}^2, H^2, H, \Gamma\}$ is automatically SUSY breaking. From the reason, we are interested in the supersymmetric quantum mechanics $\{L^2(\mathbb{R}^2; \mathbb{C}^2) \otimes \mathbb{C}^2, H^2, H, \Gamma\}$ in which H has a discrete energy ground state and this mechanics is SUSY breaking. Finally, we give the expected example as above (see Example 5.5).

The present paper is organized as follows: In Section 2, we derive into the general spectral properties of the fermionic Dirac operators. Specifically, our focus is on investigating the essential and discrete spectrums of $H(\mathbf{n})$. In Section 3, we explain a basic theory for differentiable operators via the polar coordinate conversion. Furthermore, we introduce and study the fermionic Dirac operators with the hedgehog-ansatz. In Section 4, we rigorously prove the first main theorem (i.e. Theorem 1.3). In section 5, we establish the class of unitary and selfadjoint operators ξ on $L^2(\mathbb{R}^2; \mathbb{C}^2)$ such that a quadruple $\{L^2(\mathbb{R}^2; \mathbb{C}^2) \otimes \mathbb{C}^2, H^2, H, \sigma_3 \otimes \xi\}$ embodies the supersymmetric quantum mechanics (see Lemma 5.1). Finally, we provide a sufficient condition for the SUSY breaking of this quantum mechanics (i.e. Theorem 1.4) and we give these examples (see Example 5.5).

2. CONSTRUCTION OF SPECTRUM

In this section, we denote H as the operator $H(\mathbf{n})$ defined in (1.1) as the specific vector fields \mathbf{n} under consideration are unimportant. We explore the spectral properties of the fermionic Dirac operator H .

We denote by $H_0 := (-i\sigma_2 D_1 + i\sigma_1 D_2) \otimes 1_2$ the *spin-Dirac operator*. It is known that H_0 is a self-adjoint operator on $L^2(\mathbb{R}^2; \mathbb{C}^2) \otimes \mathbb{C}^2$. Since the operator $m \sum_{j=1}^3 \sigma_3 n_j \otimes \tau_j$ is bounded and self-adjoint, it follows from the Kato-Rellich theorem (see e.g. [10, Theorem X.12]) that $\mathcal{D}(H) = \mathcal{D}(H_0)$ and $H = H_0 - m \sum_{j=1}^3 \sigma_3 n_j \otimes \tau_j$ is self-adjoint. Therefore $\sigma(H)$ is contained in \mathbb{R} . Furthermore, the symmetry of $\sigma(H)$ is established.

Lemma 2.1. The spectrum $\sigma(H)$ is symmetric with respect to zero, that is, $\lambda \in \sigma(H)$ implies that $-\lambda \in \sigma(H)$. Moreover, so are $\sigma_d(H)$ and $\sigma_{\text{ess}}(H)$.

Proof. Define an operator $\Gamma := \sigma_3 \otimes \xi$ on $L^2(\mathbb{R}^2; \mathbb{C}^2) \otimes \mathbb{C}^2$, where ξ is a unitary self-adjoint operator on $L^2(\mathbb{R}^2; \mathbb{C}^2)$ such that $\{\xi, \mathbf{n} \cdot \boldsymbol{\tau}\} = 0$ (see Example 5.2 for the existence of ξ). It is easy to see that Γ is also a self-adjoint and unitary operator. Moreover, we obtain $\Gamma \mathcal{D}(H) = \mathcal{D}(H)$. A direct calculation shows that $\{\Gamma, H\} = 0$ on $\mathcal{D}(H)$ (or see Lemma 5.1). By the relation between Γ and H , and also unitarity of Γ , we get $\sigma(H) = \sigma(\Gamma(-H)\Gamma^{-1}) = \sigma(-H)$, as desired. A similar argument to the above proof shows that $\sigma_{\text{ess}}(H)$ and $\sigma_d(H)$ are also symmetric with respect to zero. \square

Lemma 2.2. For any $\mathbf{n} = (n_1, n_2, n_3) \in \mathcal{N}$, we have

$$H^2 = (-\Delta + m^2)1_2 \otimes 1_2 - m \sum_{j=1}^3 \left(\sum_{k=1}^2 \sigma_k(D_k n_j) \right) \otimes \tau_j$$

where $\Delta := D_1^2 + D_2^2$ denotes the 2-dimensional Laplacian.

Proof. A direct computation shows

$$\begin{aligned}
H^2 &= (i\sigma_2 D_1 \otimes 1_2)^2 + (i\sigma_1 D_2 \otimes 1_2)^2 + m^2 \left(\sum_{j=1}^3 \sigma_3 n_j \otimes \tau_j \right)^2 \\
&\quad + \underbrace{\{\sigma_2 D_1, \sigma_1 D_2\}}_{=0} \otimes 1_2 + im \sum_{j=1}^3 \{\sigma_2 D_1, \sigma_3 n_j\} \otimes \tau_j - im \sum_{j=1}^3 \{\sigma_1 D_2, \sigma_3 n_j\} \otimes \tau_j \\
&= (-\Delta + m^2) 1_2 \otimes 1_2 + im \sum_{j=1}^3 \underbrace{\sigma_2 \sigma_3}_{=i\sigma_1} (D_1 n_j) \otimes \tau_j - im \sum_{j=1}^3 \underbrace{\sigma_1 \sigma_3}_{=-i\sigma_2} (D_2 n_j) \otimes \tau_j \\
&= (-\Delta + m^2) 1_2 \otimes 1_2 - m \sum_{j=1}^3 \left(\sum_{k=1}^2 \sigma_k (D_k n_j) \right) \otimes \tau_j.
\end{aligned}$$

□

For simply, we write $-\Delta + m^2$ as $(-\Delta + m^2) 1_2 \otimes 1_2$. Here, we define

$$V \equiv V(\mathbf{n}) := -m \sum_{j=1}^3 \left(\sum_{k=1}^2 \sigma_k (D_k n_j) \right) \otimes \tau_j$$

By the definition of τ_j 's, one can see the following matrix decomposition of V .

Lemma 2.3. The operator V has

$$V = \begin{pmatrix} 0 & W^* \\ W & 0 \end{pmatrix},$$

where $W \equiv W(\mathbf{n}) := -m \sum_{j=1}^3 ((D_1 n_j) + i(D_2 n_j)) \otimes \tau_j$.

Further, we define

$$(2.1) \quad L \equiv L(\mathbf{n}) := -\Delta + V$$

Then H^2 is written by $H^2 = L + m^2$. Since V is a bounded and self-adjoint operator, L is self-adjoint and $\mathcal{D}(L) = \mathcal{D}(-\Delta)$ by Kato-Rellich theorem. Finally, we derive a construction of $\sigma_{\text{ess}}(H)$ and $\sigma_{\text{d}}(H)$.

Proposition 2.4. We have $\sigma_{\text{ess}}(H) = (-\infty, -m] \cup [m, \infty)$ and $\sigma_{\text{d}}(H) \subset (-m, m)$.

Proof. By [5, Theorem 6.7] and two assumptions (n-3) and (n-4), the operator V is compact. Since $\sigma(-\Delta) = \sigma_{\text{ess}}(-\Delta) = [0, \infty)$, we get $\sigma_{\text{ess}}(L) = \sigma_{\text{ess}}(-\Delta) = [0, \infty)$ by the Weyl theorem (see e.g. [9, Example 3, Page 117]). Finally, $\sigma_{\text{ess}}(H^2) = [m^2, \infty)$, and hence it follows that $\sigma_{\text{ess}}(H) = (-\infty, -m] \cup [m, \infty)$. □

3. HEDGEHOG-ANSATZ

3.1. Differential operators under the polar coordinate conversion. Suppose the polar coordinate conversion $\Phi : \mathcal{C} \rightarrow \mathbb{R}^2$, that is, $\Phi(\omega) := (r \cos \varphi, r \sin \varphi)$ for $\omega = (r, \varphi) \in \mathcal{C}$. The map Φ is injective and $\Phi(\mathcal{C}) = \mathbb{R}^2 \setminus \{(0, 0)\}$. For a complex valued function f on domain which contains $\Phi(\mathcal{C})$ and $\omega \in \mathcal{C}$, we define $\tilde{f}(\omega) := f(\Phi(\omega))$. Then for any $\psi \in L^2(\mathbb{R}^2)$, we have

$$\int_{\mathbb{R}^2} |\psi(\mathbf{x})|^2 d\mathbf{x} = \int_{\Phi(\mathcal{C})} |\psi(\mathbf{x})|^2 d\mathbf{x} = \int_{\mathcal{C}} |\tilde{\psi}(\omega)|^2 r dr d\varphi,$$

and therefore the isometry $U : L^2(\mathbb{R}^2) \rightarrow L^2(\mathcal{C}, r dr d\varphi)$ can be defined by $U\psi = \tilde{\psi}$ for all $\psi \in L^2(\mathbb{R}^2)$. The operator U is unitary. For $(x_1, x_2) \in \mathbb{R}^2 \setminus \{(0, 0)\}$, it is easy to see that $Ux_1U^{-1} = r \cos \varphi$ and $Ux_2U^{-1} = r \sin \varphi$, where x_j is regarded as a multiplication operator on $L^2(\mathbb{R}^2)$ for $j = 1, 2$. By the definition of cylindrical coordinate conversion and the chain rule of partial differential, we see that

$$UD_1U^{-1} = \cos \varphi \cdot \partial_r - \frac{\sin \varphi}{r} \cdot \partial_\varphi \quad \text{and} \quad UD_2U^{-1} = \sin \varphi \cdot \partial_r + \frac{\cos \varphi}{r} \cdot \partial_\varphi,$$

on $U \left(\bigcap_{j=1}^2 \mathcal{D}(D_j) \right)$, where $\partial_\#$ is a partial differential operator in the variable $\# = r, \varphi$.

By the above relation, we have $\tilde{\Delta} := U\Delta U^{-1} = \partial_r^2 + \frac{1}{r}\partial_r + \frac{1}{r^2}\partial_\varphi^2$ on $U[\mathcal{D}(\Delta) \cap C^2(\Phi(\mathcal{C}))]$. The unitary invariance implies $\sigma(-\tilde{\Delta}) = \sigma_{\text{ess}}(-\tilde{\Delta}) = [0, \infty)$.

3.2. Hedgehog-ansatz. Consider $N \in \mathbb{Z}$. Define a vector field $\mathbf{n}(F) = (n_1, n_2, n_3)$, where n_j are defined in (1.2) and $F : (0, \infty) \rightarrow [0, \pi]$ is a function such that

- (F-1) F is continuously differentiable on $(0, \infty)$;
- (F-2) $\sup_{r>0} |F'(r)| < \infty$ and $F' \not\equiv 0$ on $(0, \infty)$;
- (F-3) $\lim_{r \rightarrow \infty} F'(r) = 0$.

Define $\mathcal{F} = \{F : (0, \infty) \rightarrow [0, \pi] : F \text{ satisfies the conditions (F-1) - (F-3)}\}$. For $F \in \mathcal{F}$ and $N \in \mathbb{Z}$, the vector field $\mathbf{n}(F)$ is called the *hedgehog-ansatz*¹.

Lemma 3.1. For any $F \in \mathcal{F}$, we have $\mathbf{n}(F) \in \mathcal{N}$.

Proof. Since $n_j = U^{-1}\tilde{n}_j U$ on $\Phi(\mathcal{C})$ for $j = 1, 2, 3$, we get $n_1(\mathbf{x}) = \sin F(\sqrt{x_1^2 + x_2^2}) \cdot \cos(N \text{Tan}^{-1}(x_2/x_1))$, $n_2(\mathbf{x}) = \sin F(\sqrt{x_1^2 + x_2^2}) \cdot \sin(N \text{Tan}^{-1}(x_2/x_1))$ and $n_3(\mathbf{x}) = \cos F(\sqrt{x_1^2 + x_2^2})$ for all $\mathbf{x} \in \Phi(\mathcal{C})$. By the assumptions (F-1) and (F-2), n_j is continuously differentiable and $D_k n_j$ is bounded on $\Phi(\mathcal{C})$ for $j = 1, 2, 3$ and $k = 1, 2$. It is easy to see that $\|\mathbf{n}(F)(\mathbf{x})\| = 1$ on $\Phi(\mathcal{C})$. By the chain rule of the partial differential, for $\omega = (r, \varphi) \in \mathcal{C}$, we have

$$\frac{\partial \tilde{n}_1}{\partial r}(\omega) = \frac{\partial n_1}{\partial r}(r \cos \varphi, r \sin \varphi) = \frac{\partial n_1}{\partial x_1} \frac{\partial x_1}{\partial r} + \frac{\partial n_1}{\partial x_2} \frac{\partial x_2}{\partial r} = \frac{\partial n_1}{\partial x_1} \cos \varphi + \frac{\partial n_1}{\partial x_2} \sin \varphi.$$

Since $\tilde{n}_1(r, \varphi) = \sin F(r) \cos N\varphi$, it follows from the assumption (F-3) that

$$\frac{\partial \tilde{n}_1}{\partial r}(\omega) = F'(r) \cos F(r) \cos N\varphi \rightarrow 0, \quad r \rightarrow \infty.$$

Therefore we obtain $D_1 n_1(\mathbf{x}) \rightarrow 0$ and $D_2 n_1(\mathbf{x}) \rightarrow 0$ as $\|\mathbf{x}\| \rightarrow \infty$. Finally, we get $\|\nabla n_1(\mathbf{x})\| \rightarrow 0$ as $\|\mathbf{x}\| \rightarrow \infty$. Similarly, we obtain $\|\nabla n_2(\mathbf{x})\| \rightarrow 0$ as $\|\mathbf{x}\| \rightarrow \infty$. Therefore $\mathbf{n}(F) \in \mathcal{N}$. \square

For an operator A on $L^2(\mathbb{R}^2; \mathbb{C}^2) \otimes \mathbb{C}^2$, we define the operator \tilde{A} on $L^2(\mathcal{C}; \mathbb{C}^2) \otimes \mathbb{C}^2$ by

$$\tilde{A} := (U \otimes 1_2) A (U \otimes 1_2)^{-1},$$

where we use the same symbol \tilde{A} as in the unitary conjugation by $U : L^2(\mathbb{R}^2) \rightarrow L^2(\mathcal{C}, r dr d\varphi)$. Consider $H \equiv H(\mathbf{n}(F))$, $V \equiv V(\mathbf{n}(F))$ and $L \equiv L(\mathbf{n}(F))$ for $F \in \mathcal{F}$ as below. By Lemma 3.1 and Theorem 2.4, we have $\sigma_d(\tilde{H}) = \sigma_d(H) \subset (-m, m)$. By Lemma 2.2, we have $\tilde{H}^2 = \tilde{L} + m^2 = -\tilde{\Delta} + \tilde{V} + m^2$, where

$$\tilde{V} = \begin{pmatrix} 0 & m \sum_{j=1}^3 e^{-i\varphi} \left(\frac{\partial \tilde{n}_j}{\partial r} - \frac{i}{r} \frac{\partial \tilde{n}_j}{\partial \varphi} \right) \otimes \tau_j \\ -m \sum_{j=1}^3 e^{i\varphi} \left(\frac{\partial \tilde{n}_j}{\partial r} + \frac{i}{r} \frac{\partial \tilde{n}_j}{\partial \varphi} \right) \otimes \tau_j & 0 \end{pmatrix}.$$

¹The number N is known as the topological invariant, see [1]

3.3. Reduced part with respect to the grand spin. Let $L_3 = \overline{-ix_1 D_2 + ix_2 D_1}$ be the third component of the angular momentum. Define the *grand spin* $K_3 = L_3 \otimes 1_2 + \frac{1}{2}\sigma_3 \otimes 1_2 + \frac{N}{2} \otimes \tau_3$. Note that K_3 is a self-adjoint operator on $L^2(\mathbb{R}^2; \mathbb{C}^2) \otimes \mathbb{C}^2$. By [4, (5.16)],

$$\sigma(K_3) = \sigma_p(K_3) = \left\{ \ell + \frac{s}{2} + \frac{Nt}{2} : \ell \in \mathbb{Z}, s, t \in \{1, -1\} \right\},$$

where $\sigma_p(A)$ is the point spectrum of A . By [4, (5.17)], the eigenspace of K_3 with an eigenvalue $\ell + (s/2) + (Nt/2)$ is given by $\mathcal{M}_{\ell, s, t} := \ker(L_3 - \ell) \otimes \ker(\sigma_3 - s) \otimes \ker(\tau_3 - t)$. Therefore $L^2(\mathbb{R}^2; \mathbb{C}^2) \otimes \mathbb{C}^2$ has the following orthogonal decomposition $L^2(\mathbb{R}^2; \mathbb{C}^2) \otimes \mathbb{C}^2 \cong \bigoplus_{\ell \in \mathbb{Z}, s, t \in \{\pm 1\}} \mathcal{M}_{\ell, s, t}$, under the natural identification $L^2(\mathbb{R}^2; \mathbb{C}^2) \otimes \mathbb{C}^2 \cong L^2(\mathbb{R}^2) \otimes \mathbb{C}^2 \otimes \mathbb{C}^2$.

Lemma 3.2 ([6]). Consider $H \equiv H(\mathbf{n}(F))$ for $F \in \mathcal{F}$.

- (1) The operators H and K_3 strongly commute, that is, their spectral projections commute.
- (2) For any $\Psi \in \mathcal{D}(H)$, we obtain $P_{\mathcal{M}_{\ell, s, t}} \Psi \in \mathcal{D}(H)$ and $HP_{\mathcal{M}_{\ell, s, t}} \Psi = P_{\mathcal{M}_{\ell, s, t}} H \Psi$, where $P_{\mathcal{M}_{\ell, s, t}}$ is a projection onto $\mathcal{M}_{\ell, s, t}$.

Proof. The first statement can be proven using a similar approach as outlined in [4, Lemmas 5.2 and 5.3]. The second statement follows readily from the first one. \square

By Lemma 3.2, one can define the reduced part $H(\ell, s, t)$ of H to $\mathcal{M}_{\ell, s, t}$, that is, $H(\ell, s, t) := H \upharpoonright_{\mathcal{D}(H(\ell, s, t))}$, where $\mathcal{D}(H(\ell, s, t)) = \mathcal{D}(H) \cap \mathcal{M}_{\ell, s, t}$. Note that $\mathcal{D}(\tilde{H}(\ell, s, t)) = \mathcal{D}(\tilde{H}) \cap \tilde{\mathcal{M}}_{\ell, s, t}$, where $\tilde{\mathcal{M}}_{\ell, s, t} := \ker(-i\partial_\varphi - \ell) \otimes \ker(\sigma_3 - s) \otimes \ker(\tau_3 - t)$. Since $UL_3U^{-1} = -i\partial_\varphi$, we get $-\frac{\partial^2}{\partial\varphi^2} \upharpoonright_{\mathcal{D}(\tilde{H}(\ell, s, t))} = \ell^2$, and therefore $\tilde{\Delta} \upharpoonright_{\mathcal{D}(\tilde{H}(\ell, s, t))} = \frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{\ell^2}{r^2}$. In the following, we denote by

$$\Delta_\ell := \frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{\ell^2}{r^2}, \quad \ell \in \mathbb{R}.$$

Moreover, for any $F \in \mathcal{F}$, we get

$$\tilde{V} \upharpoonright_{\mathcal{D}(\tilde{H}(\ell, s, t))} \equiv \tilde{V}(\mathbf{n}(F)) \upharpoonright_{\mathcal{D}(\tilde{H}(\ell, s, t))} = \begin{pmatrix} 0 & W(t)^* \\ W(t) & 0 \end{pmatrix},$$

where

$$(W(t)\Psi)(\omega) := -me^{i\varphi} \left\{ \sum_{j=1}^2 \left(\frac{\partial \tilde{n}_j}{\partial r} + \frac{i}{r} \frac{\partial \tilde{n}_j}{\partial \varphi} \right) \otimes \tau_j + t \frac{\partial \tilde{n}_3}{\partial r} \otimes 1_2 \right\} \Psi(\omega)$$

for any $\Psi \in \mathcal{D}(W(t))$ and $\omega = (r, \varphi) \in \mathcal{C}$. Finally, we get

$$\tilde{H}(\ell, s, t)^2 = -\Delta_\ell + \underbrace{\begin{pmatrix} 0 & W(t)^* \\ W(t) & 0 \end{pmatrix}}_{=: L(\ell, t)} + m^2$$

on $C_0^\infty((0, \infty))$. Note that $L(\ell, t) = \tilde{L} \upharpoonright_{\mathcal{D}(\tilde{H}(\ell, s, t))}$ and $\sigma_{\text{ess}}(L(\ell, t)) \subset [0, \infty)$.

4. EXISTENCE OF DISCRETE GROUND STATES

In this section, we establish sufficient conditions for that $H \equiv H(\mathbf{n}(F))$ has a discrete positive or negative energy ground state (see Theorems 1.3 and 1.4). For after discussions, we denote by

$$v_\ell(f)(\omega) = f(r)e^{i\ell\varphi}$$

for all $\omega = (r, \varphi) \in \mathcal{C}$ and complex-valued functions f on $(0, \infty)$ and $\ell \in \mathbb{Z}$. Let $u_t \in \mathbb{C}^2$ be an eigenvector of τ_3 for each $t \in \{1, -1\}$, that is, $\tau_3 u_t = t u_t$ for each $t \in \{1, -1\}$. Further, we consider $\ell \in \mathbb{Z}$ and unit vectors $f \in C_0^\infty((0, \infty))$. We then define the following vectors:

$$\Psi_{f,+}^{(\ell,t)} = (v_\ell(f) \otimes u_t, 0)^\top \in \widetilde{\mathcal{M}}_{\ell,1,t} \quad \text{and} \quad \Psi_{f,-}^{(\ell,t)} = (0, v_\ell(f) \otimes u_t)^\top \in \widetilde{\mathcal{M}}_{\ell,-1,t}.$$

Moreover, for $\ell_1, \ell_2 \in \mathbb{Z}$ and $t \in \{1, -1\}$, we define $\Psi_f^{(\ell_1, \ell_2, t)} := \frac{1}{\sqrt{2}} \left(\Psi_{f,+}^{(\ell_1, t)} + \Psi_{f,-}^{(\ell_2, t)} \right)$.

Lemma 4.1. Let us consider $\ell_1, \ell_2 \in \mathbb{Z}$, $t \in \{1, -1\}$ and a unit vector $f \in C_0^\infty((0, \infty))$. Moreover, $\widetilde{V} \equiv \widetilde{V}(\mathbf{n}(F))$ for $F \in \mathcal{F}$. Then we obtain the following formula.

$$(1) \quad \langle \Psi_f^{(\ell_1, \ell_2, t)}, \widetilde{\Delta} \Psi_f^{(\ell_1, \ell_2, t)} \rangle = \frac{1}{2} \sum_{k=1}^2 \langle f, \Delta_{\ell_k} f \rangle_2.$$

$$(2) \quad \langle \Psi_f^{(\ell_1, \ell_2, t)}, \widetilde{V} \Psi_f^{(\ell_1, \ell_2, t)} \rangle = mt \langle f, (F' \sin F) f \rangle_2 \delta_{\ell_1+1, \ell_2}$$

Proof. (1) Since $\langle \Psi_{f,+}^{(\ell_1, t)}, \widetilde{\Delta} \Psi_{f,-}^{(\ell_2, t)} \rangle = \langle \Psi_{f,-}^{(\ell_2, t)}, \widetilde{\Delta} \Psi_{f,+}^{(\ell_1, t)} \rangle = 0$, we obtain

$$\begin{aligned} \langle \Psi_f^{(\ell_1, \ell_2, t)}, \widetilde{\Delta} \Psi_f^{(\ell_1, \ell_2, t)} \rangle &= \frac{1}{2} \left(\langle \Psi_{f,+}^{(\ell_1, t)}, \widetilde{\Delta} \Psi_{f,+}^{(\ell_1, t)} \rangle + \langle \Psi_{f,-}^{(\ell_2, t)}, \widetilde{\Delta} \Psi_{f,-}^{(\ell_2, t)} \rangle \right) \\ &= \frac{1}{2} \left(\langle \Psi_{f,+}^{(\ell_1, t)}, \Delta_{\ell_1} \Psi_{f,+}^{(\ell_1, t)} \rangle + \langle \Psi_{f,-}^{(\ell_2, t)}, \Delta_{\ell_2} \Psi_{f,-}^{(\ell_2, t)} \rangle \right) \\ &= \frac{1}{2} (\langle f, \Delta_{\ell_1} f \rangle_2 + \langle f, \Delta_{\ell_2} f \rangle_2). \end{aligned}$$

(2) We have

$$\begin{aligned} \langle \Psi_{f,+}^{(\ell_1, t)}, \widetilde{V} \Psi_{f,+}^{(\ell_1, t)} \rangle &= \left\langle \begin{pmatrix} v_{\ell_1}(f) \otimes u_t \\ 0 \end{pmatrix}, \begin{pmatrix} 0 & W(t)^* \\ W(t) & 0 \end{pmatrix} \begin{pmatrix} v_{\ell_1}(f) \otimes u_t \\ 0 \end{pmatrix} \right\rangle \\ &= \left\langle \begin{pmatrix} v_{\ell_1}(f) \otimes u_t \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ W(t)(v_{\ell_1}(f) \otimes u_t) \end{pmatrix} \right\rangle \\ &= 0. \end{aligned}$$

Similarly, we obtain $\langle \Psi_{f,-}^{(\ell_2, t)}, \widetilde{V} \Psi_{f,-}^{(\ell_2, t)} \rangle = 0$. Hence,

$$\begin{aligned} \langle \Psi_f^{(\ell_1, \ell_2, t)}, \widetilde{V} \Psi_f^{(\ell_1, \ell_2, t)} \rangle &= \frac{1}{2} \left(\langle \Psi_{f,+}^{(\ell_1, t)}, \widetilde{V} \Psi_{f,-}^{(\ell_2, t)} \rangle + \langle \Psi_{f,-}^{(\ell_2, t)}, \widetilde{V} \Psi_{f,+}^{(\ell_1, t)} \rangle \right) \\ &= \frac{1}{2} (\langle v_{\ell_1}(f) \otimes u_t, W(t)^*(v_{\ell_2}(f) \otimes u_t) \rangle + \langle v_{\ell_2}(f) \otimes u_t, W(t)(v_{\ell_1}(f) \otimes u_t) \rangle) \\ &= \text{Re} \langle W(t)(v_{\ell_1}(f) \otimes u_t), v_{\ell_2}(f) \otimes u_t \rangle \\ &= -mt \text{Re} \left(\int_0^{2\pi} e^{-i(1+\ell_1)\varphi} e^{i\ell_2\varphi} \cdot \frac{1}{2\pi} d\varphi \right) \left(\int_{(0, \infty)} \frac{d \cos F(r)}{dr} \overline{f(r)} f(r) r dr \right) \\ &= mt \langle f, (F' \sin F) f \rangle_2 \delta_{\ell_1+1, \ell_2}. \end{aligned}$$

□

For $\ell \in \mathbb{Z}$, we define $c(\ell) := \sqrt{\frac{\ell^2 + (\ell+1)^2}{2}}$ and

$$R(\ell, t) := -\Delta_{c(\ell)} + mt \frac{dF}{dr} \sin F,$$

acting on $L^2((0, \infty), r dr)$ with domain $\mathcal{D}(R(\ell, t)) := C_0^\infty((0, \infty))$.

Theorem 4.2 (Theorem 1.3). Let us consider $F \in \mathcal{F}$. If there exist $\ell \in \mathbb{Z}$ and $t \in \{1, -1\}$ such that

$$(4.1) \quad \mathcal{E}_0(R(\ell, t)) := \inf_{\substack{f \in C_0^\infty((0, \infty)) \\ \|f\|_2=1}} \langle f, R(\ell, t)f \rangle_2 < 0,$$

then H has a discrete positive or negative energy ground state.

Proof. For any $\ell \in \mathbb{Z}$, $t \in \{1, -1\}$ and unit vectors $f \in C_0^\infty((0, \infty))$, we get

$$\begin{aligned} \langle \Psi_f^{(\ell, \ell+1, t)}, \tilde{L}\Psi_f^{(\ell, \ell+1, t)} \rangle &= \langle \Psi_f^{(\ell, \ell+1, t)}, -\tilde{\Delta}\Psi_f^{(\ell, \ell+1, t)} \rangle + \langle \Psi_f^{(\ell, \ell+1, t)}, \tilde{V}\Psi_f^{(\ell, \ell+1, t)} \rangle \\ &= \langle f, (-\Delta_{c(\ell)} + mt(F' \sin F))f \rangle_2 = \langle f, R(\ell, t)f \rangle_2 \end{aligned}$$

by Lemma 4.1. If $\mathcal{E}_0(R(\ell, t)) < 0$ for some $\ell \in \mathbb{Z}$ and $t \in \{1, -1\}$, then there exists a unit vector $f \in C_0^\infty((0, \infty))$ such that $\langle f, R(\ell, t)f \rangle_2 < 0$. The above equality shows that $\langle \Psi_f^{(\ell, \ell+1, t)}, \tilde{L}\Psi_f^{(\ell, \ell+1, t)} \rangle < 0$. Since $\sigma_{\text{ess}}(\tilde{L}) = [0, \infty)$, it follows from the min-max principal (see [9, Theorem XIII.2]) that the operator \tilde{L} has a discrete ground state. Therefore so does \tilde{H}^2 . It follows the existence of a discrete positive or negative energy ground state of \tilde{H} (and so is H). \square

5. SUPERSYMMETRY BREAKING

A quadruple $\{\mathcal{H}, Q^2, Q, \Gamma\}$ is called a *supersymmetric quantum mechanics* if \mathcal{H} is a Hilbert space, Q is a self-adjoint operator on \mathcal{H} and Γ is a unitary self-adjoint operator on \mathcal{H} such that $\{\Gamma, Q\} = 0$. In the physical literature, the operators Q and Γ are called a *supercharge* and a *grading operator*, respectively. Initially, we build a supersymmetric quantum mechanics by the fermionic Dirac operators as follows.

Lemma 5.1. Let $H \equiv H(\mathbf{n})$ be the fermionic Dirac operator and ξ a unitary self-adjoint operator on $L^2(\mathbb{R}^2; \mathbb{C}^2)$. A quadruple $\mathcal{Q}(\mathbf{n}, \xi) := \{L^2(\mathbb{R}^2; \mathbb{C}^2) \otimes \mathbb{C}^2, H^2, H, \sigma_3 \otimes \xi\}$ consists a supersymmetric quantum mechanics if and only if $\{\xi, \mathbf{n} \cdot \boldsymbol{\tau}\} = 0$.

Proof. Let us set $\Gamma = \sigma_3 \otimes \xi$. We can regard $m \sum_{j=1}^3 \sigma_3 n_j \otimes \tau_j$ as the operator $m\sigma_3 \otimes (\mathbf{n} \cdot \boldsymbol{\tau})$. Since $\{\sigma_3, \sigma_1\} = \{\sigma_3, \sigma_2\} = 0$, we get $\{\sigma_3, (-i\sigma_2 D_1 + i\sigma_1 D_2)\} = 0$, and therefore $\{\Gamma, H_0\} = 0$ without a condition of ξ . Thus, the desired result holds thanks to the following equality

$$\{\Gamma, \sigma_3 \otimes (\mathbf{n} \cdot \boldsymbol{\tau})\} = [\sigma_3, \sigma_3] \otimes \xi(\mathbf{n} \cdot \boldsymbol{\tau}) + \sigma_3^2 \otimes \{\xi, \mathbf{n} \cdot \boldsymbol{\tau}\} = 1_2 \otimes \{\xi, \mathbf{n} \cdot \boldsymbol{\tau}\}.$$

\square

By using Lemma 5.1, we present a characterization of unitary self-adjoint operators ξ on $L^2(\mathbb{R}^2; \mathbb{C}^2)$ such that $\mathcal{Q}(\mathbf{n}, \xi)$ constitutes a supersymmetric quantum mechanics.

Example 5.2. If $\xi = f_0\tau_0 + f_1\tau_1 + f_2\tau_2 + f_3\tau_3$ for some $f_j \in L^2(\mathbb{R}^2)$ ($j = 0, 1, 2, 3$) and $\tau_0 = 1_2$, then

- (i) For any $\mathbf{x} \in \mathbb{R}^2$, we have $f_j(\mathbf{x}) \in \mathbb{R}$ for all $j = 0, 1, 2, 3$ (since ξ is self-adjoint).
- (ii) For any $\mathbf{x} \in \mathbb{R}^2$, $\sum_{j=0}^3 |f_j(\mathbf{x})|^2 = 1$ (since ξ is unitary).

Note that $\{\xi, \mathbf{n} \cdot \boldsymbol{\tau}\} = 2(\sum_{j=1}^3 f_j n_j)\tau_0 + 2f_0 \sum_{j=1}^3 n_j \tau_j$. If $\{\xi, \mathbf{n} \cdot \boldsymbol{\tau}\} = 0$, then $\sum_{j=1}^3 f_j n_j = 0$ and $f_0 n_j = 0$ for all $j = 1, 2, 3$ since $\{\tau_j\}_{j=0}^3$ is an orthonormal basis in $M_2(\mathbb{C})$. Because $\|\mathbf{n}(\mathbf{x})\| = 1$ a.e.

$\mathbf{x} \in \mathbb{R}^2$, we have $(n_1, n_2, n_3) \neq (0, 0, 0)$, and therefore it must be $f_0 = 0$. We particularly consider $n_3 \neq 0$. Then $f_3 = -\frac{n_1}{n_3}f_1 - \frac{n_2}{n_3}f_2$. Finally, the above condition (ii) implies

$$1 = \left(1 + \frac{n_1(\mathbf{x})^2}{n_3(\mathbf{x})^2}\right) f_1(\mathbf{x})^2 + \left(1 + \frac{n_2(\mathbf{x})^2}{n_3(\mathbf{x})^2}\right) f_2(\mathbf{x})^2 - \frac{2n_1(\mathbf{x})n_2(\mathbf{x})}{n_3(\mathbf{x})^2} f_1(\mathbf{x})f_2(\mathbf{x}),$$

for $\mathbf{x} \in \mathbb{R}^2$. For example,

- If $f_2 = 0$, then $f_1 = \frac{|n_3|}{\sqrt{n_1^2+n_3^2}}$ and $f_3 = \frac{|n_1|}{\sqrt{n_1^2+n_3^2}}$. Hence $\xi = \frac{|n_3|}{\sqrt{n_1^2+n_3^2}}\tau_1 + \frac{|n_1|}{\sqrt{n_1^2+n_3^2}}\tau_3$.
- Similarly, if $f_1 = 0$, then $\xi = \frac{|n_3|}{\sqrt{n_2^2+n_3^2}}\tau_2 + \frac{|n_2|}{\sqrt{n_2^2+n_3^2}}\tau_3$.

By Lemma 5.1, $\mathcal{Q}(\mathbf{n}, \xi)$ consists a supersymmetric quantum mechanics under the above cases for ξ .

A supersymmetric quantum mechanics $\{\mathcal{H}, Q^2, Q, \Gamma\}$ is said to have a *supersymmetric state* if $\ker Q \neq \{0\}$. Furthermore, a supersymmetric quantum mechanics is said to be *supersymmetry (SUSY) breaking* if it has no supersymmetric states.

Lemma 5.3 ([2]). Let us

$$V_0 = \begin{pmatrix} 0 & W_0^* \\ W_0 & 0 \end{pmatrix},$$

where W_0 is a bounded operator on some Hilbert space. Then we have $\sigma(V_0^2) = \sigma(W_0W_0^*)$. In particular, $\|V_0\| = \|W_0\|$.

We recall Lemma 2.2 and Remark 2.3 that for $\mathbf{n} = (n_1, n_2, n_3) \in \mathcal{N}$,

$$H^2 = -\Delta + m^2 + \underbrace{\begin{pmatrix} 0 & W^* \\ W & 0 \end{pmatrix}}_{=V},$$

where $W = -m \sum_{j=1}^3 ((D_1 n_j) + i(D_2 n_j)) \otimes \tau_j$. If $\mathbf{n} = \mathbf{n}(F)$ is the hedgehog-ansatz with $F \in \mathcal{F}$ and $\limsup_{r \rightarrow 0^+} \left| \frac{\sin F(r)}{r} \right| < \infty$, then

$$\widetilde{W} := (U \otimes 1_2)W(U \otimes 1_2)^{-1} = -m \begin{pmatrix} \widetilde{R}_1 & \widetilde{R}_2 \\ \widetilde{R}_3 & -\widetilde{R}_1 \end{pmatrix},$$

where

$$\begin{aligned} \widetilde{R}_1 &= -F'(r) \sin F(r) e^{i\varphi}, \\ \widetilde{R}_2 &= \left(F'(r) \cos F(r) + \frac{N}{r} \sin F(r) \right) e^{-i(N-1)\varphi}, \\ \widetilde{R}_3 &= \left(F'(r) \cos F(r) - \frac{N}{r} \sin F(r) \right) e^{i(N+1)\varphi}. \end{aligned}$$

One can see that

$$(5.1) \quad \|\widetilde{W}\| = m \sup_{r>0} \sqrt{\left| F'(r)^2 - \left(\frac{N}{r} \sin F(r) \right)^2 \right|}.$$

Theorem 5.4 (Theorem 1.4). Let us consider $\mathbf{n} = \mathbf{n}(F) \in \mathcal{N}$ with $\limsup_{r \rightarrow 0^+} \left| \frac{\sin F(r)}{r} \right| < \infty$. If

$$(5.2) \quad m > \sup_{r>0} \sqrt{\left| F'(r)^2 - \left(\frac{N}{r} \sin F(r) \right)^2 \right|},$$

then $\mathcal{Q}(\mathbf{n}, \xi)$ is SUSY breaking for any unitary self-adjoint operators ξ with $\{\xi, \mathbf{n} \cdot \boldsymbol{\tau}\} = 0$.

Proof. Fix an arbitrary unitary self-adjoint operator ξ such that $\{\xi, \mathbf{n} \cdot \boldsymbol{\tau}\} = 0$. Then

$$\begin{aligned} E_0(H^2) &= \inf_{\|\Phi\|=1} \langle \Phi, H^2 \Phi \rangle \\ &\geq m^2 + \inf_{\|\Phi\|=1} \langle \Phi, V \Phi \rangle \\ &\geq m^2 - \|V\| \\ &= m^2 - \|\widetilde{W}\| \quad (\text{by Lemma 5.3 and the unitary invariance}) \\ &= m^2 - m \sup_{r>0} \sqrt{\left| F'(r)^2 - \left(\frac{N}{r} \sin F(r) \right)^2 \right|} \quad (\text{by (5.1)}). \end{aligned}$$

The assumption (5.2) shows that $E_0(H^2) > 0$, and therefore $\text{Ker}H = \{0\}$ as desired. \square

To end this section, we provide an example of the supersymmetric quantum mechanics $\mathcal{Q}(\mathbf{n}(F), \xi)$ in which $H = H(\mathbf{n}(F))$ has a discrete ground state while being the absence of supersymmetry. More precisely, we find $F \in \mathcal{F}$ such that $\limsup_{r \rightarrow 0^+} \left| \frac{\sin F(r)}{r} \right| < \infty$ and two conditions (4.1) and (5.2) hold true. According to Theorems 1.3 and 1.4, such a vector field $\mathbf{n}(F)$ presents the desired supersymmetric quantum mechanics $\mathcal{Q}(\mathbf{n}(F), \xi)$. Before giving it, we give the following mollifier:

$$(5.3) \quad f_0(r) := C r e^{-\frac{1}{1-r^2}} \chi_{\{0 < r < 1\}}(r) \quad \text{and} \quad C := \left(\int_0^1 r^3 e^{-\frac{2}{1-r^2}} dr \right)^{-\frac{1}{2}}$$

where χ_A is the indicator function on $A \subset (0, \infty)$. One can see that the function f_0 is in $C_0^\infty((0, \infty))$ and $\|f_0\|_2 = 1$.

Example 5.5. If we set $N = 1$ and define $F(r) = \frac{\pi}{r+1}$, $r > 0$, then $F \in \mathcal{F}$ and $\limsup_{r \rightarrow 0^+} \left| \frac{\sin F(r)}{r} \right| < \infty$. Consequently, we obtain

$$\sup_{r>0} \sqrt{\left| F'(r)^2 - \left(\frac{N}{r} \sin F(r) \right)^2 \right|} \approx 1.223.$$

To meet the condition (5.2), we choose $m \geq 1.23$.

Next, we choose m such that $\frac{1}{2} < \frac{9\pi}{200}m - \frac{5}{2}$, which implies $m > \frac{200}{3\pi} \approx 21.22$. Additionally, we choose $\ell \in \mathbb{Z}$ such that $c(\ell)^2 < \frac{9\pi}{200}m - \frac{5}{2}$. The existence of such ℓ is guaranteed due to the chosen value of m and the fact that $c(\ell)^2 \geq \frac{1}{2}$ holds for all $\ell \in \mathbb{Z}$. For the function f_0 defined in (5.3),

$$-\Delta_{c(\ell)} f_0(r) = \left\{ \frac{-4(3r^2 - 2)}{(1-r^2)^4} + \frac{c(\ell)^2 - 1}{r^2} \right\} f_0(r),$$

and therefore $\langle f_0, -\Delta_{c(\ell)} f_0 \rangle_2 \leq 15 + 6c(\ell)^2$. Moreover, if we take $t = 1$, then

$$\begin{aligned} \langle f_0, m t F' \sin F f_0 \rangle_2 &= \langle f_0, -\frac{m\pi}{(r+1)^2} \sin\left(\frac{\pi}{r+1}\right) f_0 \rangle_2 \\ &= -m\pi \underbrace{\left(\int_0^1 r^3 e^{-\frac{2}{1-r^2}} dr \right)^{-1}}_{\approx 270.23} \underbrace{\int_0^1 \frac{r^3}{(r+1)^2} \sin\left(\frac{\pi}{r+1}\right) e^{-\frac{2}{1-r^2}} dr}_{\approx 0.00135476} \\ &\leq -\frac{27\pi}{100}m. \end{aligned}$$

With the appropriate choice of values for m and ℓ , we ultimately attain

$$\begin{aligned} \mathcal{E}_0(-\Delta_{c(\ell)} + mtF' \sin F) &\leq \langle f_0, (-\Delta_{c(\ell)} + mtF' \sin F) f_0 \rangle_2 \\ &= 15 + 6c(\ell)^2 - \frac{27\pi}{100}m \\ &< 0. \end{aligned}$$

The conditions (4.1) and (5.2) are satisfied, especially when $m \geq 21.23$.

ACKNOWLEDGEMENT

This work was started when S.O. was affiliated with Department of Mathematics, Tohoku University. D.F. and Y.U. thank Kazuyuki Wada for fruitful discussions.

REFERENCES

- [1] Y. Amari, M. Iida and N. Sawado, Statistical nature of baby-Skyrme models in 2 + 1 dimensions and normalizable fermions, *Theoret. and Math. Phys.* **200** (2019), no. 3, 1253-1268.
- [2] A. Arai, *Analysis on Fock Spaces and Mathematical Theory of Quantum Fields: An Introduction to Mathematical Analysis of Quantum Fields*, World Scientific Pub. Co. Inc. (2018) 862 pp.
- [3] A. Arai, Mathematical analysis of a generalized chiral quark soliton model, *SIGMA Symmetry Integrability Geom. Methods Appl.* **2** (2006), Paper 018, 12 pp.
- [4] A. Arai, K. Hayashi, I. Sasaki, Spectral properties of a Dirac operator in the chiral quark soliton model, *J. Math. Phys.* **46**, 052306.
- [5] F. Gesztesy and M. Waurick, The Callias index formula revisited. *Lecture Notes in Mathematics*, **2157**. Springer, 2016. ix+192 pp.
- [6] A. Yu. Loginov, Fermion scattering on topological solitons in the nonlinear $O(3)$ σ -model, *Phys. Rev. D* **104**, 045011 (2021).
- [7] N. Nagaosa and Y. Tokura, Topological properties and dynamics of magnetic skyrmions, *Nature Nanotechnology*, **8**, 899-911 (2013).
- [8] I. Perapechka and Y. Shnir, Fermion exchange interaction between magnetic skyrmions. *Phys. Rev. D.* **99** (2019), no. 12, 125001, 7 pp.
- [9] M. Reed and B. Simon, *Methods of Modern Mathematical Physics IV*, Analysis of Operators (Academic, New York, 1978).
- [10] M. Reed and B. Simon, *Methods of Modern Mathematical Physics II*, Fourier Analysis, Self-Adjointness (Academic, New York, 1975).

DAIJU FUNAKAWA: DEPARTMENT OF ELECTRONICS AND INFORMATION ENGINEERING, HOKKAI-GAKUEN UNIVERSITY, SAPPORO, HOKKAIDO 062-8605, JAPAN

Email address: funakawa@hgu.jp

SATOSHI OKUMURA: YOKOHAMA, KANAGAWA, JAPAN

Email address: s.okumura.tohoku@gmail.com

YUKI UEDA: DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY OF EDUCATION, 9 HOKUMON-CHO, ASAHIKAWA, HOKKAIDO 070-8621, JAPAN

Email address: ueda.yuki@a.hokkyodai.ac.jp